

Real Analysis HW #6

1 a.) $b-a$ is length of segment. set $\delta := \min \{N, b-a\}$
 Where $\frac{b-a}{N} < \delta$ (by AP) then a partition with lengths $\frac{b-a}{n}$,
 the $\|P\|$ will be less than δ

b.) Suppose $\int_a^b f_1 \neq \int_a^b f_2$ are the two RI values
 of f . f is RI, so $\|P\| < \delta_1$, $|S(P, f) - \int_a^b f_1| < \epsilon/2$
 and $\|P\| < \delta_2$, $|S(P, f) - \int_a^b f_2| < \epsilon/2$

Set $\delta := \min \{\delta_1, \delta_2\}$ And $|\int_a^b f_1 - \int_a^b f_2| \neq 0$

$$\|P\| < \delta, \text{ then } |\int_a^b f_1 - \int_a^b f_2| = |\int_a^b f_1 - S(f, P) + S(f, P) - \int_a^b f_2|$$

$$\leq |S(f, P) - \int_a^b f_1| + |S(f, P) - \int_a^b f_2| < \frac{\epsilon}{2} + \frac{\epsilon}{2} < \epsilon$$

$$\text{Set } \epsilon = \frac{|\int_a^b f_1 - \int_a^b f_2|}{2}$$

$$|\int_a^b f_1 - \int_a^b f_2| < \frac{|\int_a^b f_1 - \int_a^b f_2|}{2} \quad 1 < \frac{1}{2} \quad 2 < 1 \quad \#$$

$\int_a^b f$ is unique

$$2 \text{ a.) } \|P\| < \delta_1 \rightarrow |S(f, P) - \int_a^b f| < \epsilon/2 \quad \|P\| < \delta_2 \rightarrow |S(g, P) - \int_a^b g| < \epsilon/2$$

$$\text{set } \delta = \min \{\delta_1, \delta_2\} \text{ then } \|P\| < \delta \quad |S(f+g, P) - \int_a^b f - \int_a^b g| < \epsilon$$

$$S(f+g, P) = \sum_{i=1}^n (f+g)(c_i)(x_i - x_{i-1}) = \sum_{i=1}^n f(c_i)(x_i - x_{i-1}) + \sum_{i=1}^n g(c_i)(x_i - x_{i-1})$$

number so distributive

$$S(f+g, P) = S(f, P) + S(g, P) \rightarrow |S(f, P) - \int_a^b f + S(g, P) - \int_a^b g| < \frac{\epsilon}{2} + \frac{\epsilon}{2} < \epsilon$$

want

$$2 \text{ b)} f(x) \leq g(x) \quad \int_a^b f \leq \int_a^b g$$

$$S(f, P) \leq S(g, P) \rightarrow \sum_{i=1}^n f(c_i)(x_i - x_{i-1}) \leq \sum_{i=1}^n g(c_i)(x_i - x_{i-1}) \text{ because } f \leq g$$

$$\text{Let } \varepsilon = \frac{\int_a^b f - \int_a^b g}{2} \quad \|P\| < \delta_1 \rightarrow |S(f, P) - \int_a^b f| < \frac{\int_a^b f - \int_a^b g}{2}$$

$$\|P\| < \delta_2 \rightarrow |S(g, P) - \int_a^b g| < \frac{\int_a^b f - \int_a^b g}{2}$$

$$\text{Take } \delta := \min \{ \delta_1, \delta_2 \}$$

$$\text{then } -\frac{\int_a^b f - \int_a^b g}{2} < S(f, P) - \int_a^b f < \frac{\int_a^b f - \int_a^b g}{2}$$

$$\text{then } -\frac{\int_a^b f - \int_a^b g}{2} < S(g, P) - \int_a^b g < \frac{\int_a^b f - \int_a^b g}{2}$$

$$\frac{\int_a^b f - \int_a^b g}{2} < S(f, P) \quad S(g, P) < \frac{\int_a^b f - \int_a^b g}{2}$$

$$\text{so } S(g, P) < \frac{\int_a^b f - \int_a^b g}{2} < S(f, P)$$

Which is a contradiction of $S(f, P) \leq S(g, P)$ #

3 We know that $\int_a^b k = k(b-a)$ for some constant k . If we set $M=k$ then $\int_a^b M = M(b-a)$.

We know that $|f(x)| \leq M$ so, from #2b, we

know that $\int_a^b f \leq \int_a^b M = M(b-a)$ so

$$\int_a^b f \leq M(b-a)$$

$$4 \quad \lim_{n \rightarrow \infty} \|P_n\| \rightarrow 0 \quad \|P_n\| - 0 < \varepsilon \quad \forall \varepsilon > 0, \exists N \text{ s.t. } n \geq N \text{ then } \|P_n\| < \varepsilon$$

$$n \geq N \text{ then } |S(f, P_n) - \int_a^b f| < \varepsilon \quad S(f, P_n) = \sum_{i=1}^n f(c_i)(x_i - x_{i-1})$$

$$(x_i - x_{i-1}) \rightarrow 0 \text{ so } 0 < |0 - \int_a^b f| < \varepsilon \quad \int_a^b f < \varepsilon$$

6 a.) Let P be tagged partition with $\|P\| < \delta$.

$$S(f, P) = \sum_{i=1}^N f(c_i)(x_i - x_{i-1}) \quad f(c_i) \quad \forall c_i \in c_1, \dots, c_N = k$$

$$S(f, P) = \sum_{i=1}^N k(x_i - x_{i-1}) \quad \text{since } P \text{ is tagged partition,}$$

$$\sum_{i=1}^N (x_i - x_{i-1}) = (b-a) \quad \sum_{i=1}^N k(x_i - x_{i-1}) = k \sum_{i=1}^N (x_i - x_{i-1}) = k(b-a)$$

$$\text{So } |S(f, P) - \int_a^b f| < \epsilon \quad \text{for } \epsilon > 0 \quad |S(f, P) - k(b-a)| = |k(b-a) - k(b-a)|$$

$$= 0 < \epsilon \quad \text{so } |S(f, P) - \int_a^b f| < \epsilon \quad \text{so } f \text{ is RI on } [a, b]$$

$$\text{and } \int_a^b f = k(b-a)$$

b.) $\int_a^b \sin^2(x) \quad \sin^2(x) = \frac{1 - \cos(2x)}{2} = \frac{1}{2} - \frac{1}{2}\cos(2x)$

$$\int_a^b \frac{1}{2} - \frac{1}{2}\cos(2x) \rightarrow \int_a^b \frac{1}{2} + \int_a^b -\frac{1}{2}\cos(2x) \quad 2 \in R \text{ so RI}$$

$$\int_a^b \frac{1}{2} = \frac{1}{2} \int_a^b 1 \quad \text{so } \int_a^b \sin^2(2x) \text{ is RI on } [a, b]$$

7 Let P be t.p. of $[0, 1]$ s.t. $\|P\| < \delta$.

$$\text{Say } \int_a^b f = \frac{1}{2} \text{ then } |S(f, P) - \frac{1}{2}| < \epsilon$$

$$S(f, P) = \sum_{i=1}^N f(c_i)(x_i - x_{i-1}) \quad \text{If } c_i \geq \frac{1}{2}, \text{ then } f(c_i) = 0 \text{ and}$$

$$\text{if } c_i < \frac{1}{2}, \text{ then } f(c_i) = 1 \text{ so } S(f, P) = \sum_{i=1}^N (x_i - x_{i-1}) = \frac{1}{2} - 0$$

$$= \frac{1}{2} \quad \text{so } |\frac{1}{2} - \frac{1}{2}| < \epsilon \quad 0 < \epsilon$$

ϵ was arbitrary, f is RI on $[0, 1]$.

8 Say $\int_a^b f = 0$ b.c. more R than N. P be t.p. $[0, 1]$ s.t.

$$\|P\| < \delta \text{ then } |S(f, P)| < \epsilon \quad S(f, P) = \sum_{i=1}^N f(c_i)(x_i - x_{i-1})$$

$$E_\epsilon = \{x \in \mathbb{R} : \frac{1}{N} \geq \frac{\epsilon}{2}\} \text{ For } x \in E_\epsilon \text{ then } N \leq \frac{2}{\epsilon} \text{ there is a } N_0 \text{ s.t. } N_0 = \lceil \frac{2}{\epsilon} \rceil$$

Since set $[0, 1]$ is finite E_ϵ is finite. $P_0 \leq P$ s.t. $P_0 = \{c_i : x_i - x_{i-1} \geq \frac{\epsilon}{2}\}$

$$P = P/P_0 \cup P_0, \quad P_0 \text{ is finite b.c. } E_\epsilon \text{ is.} \quad S(f, P) = S(f, P/P_0) + S(f, P_0)$$

$$S(f, P) = \sum_{i=1}^N f(c_i)(x_i - x_{i-1}) \leq \frac{\delta}{N} \quad S(f, P/P_0) = \sum_{i=1}^N f(c_i)(x_i - x_{i-1}) < \frac{\epsilon}{2}$$

$$\delta = \frac{N\epsilon}{2} \quad |S(f, P)| = S(f, P/P_0) + S(f, P_0) < \frac{\epsilon}{2} + \frac{\delta}{N} = \frac{\epsilon}{2} + \frac{N\epsilon}{2N} = \epsilon \quad \text{so } \int_a^b f = 0$$

4 Let P be a t.p. of $[0, 1]$ s.t. $\|P\| < \delta$. Then
 say $f' = 0$. Then $|S(f, P) - 0| < \epsilon \Rightarrow |S(f, P)| < \epsilon$
 $S(f, P) = \sum_{i=1}^n f(c_i)(x_i - x_{i-1})$. if $x_i > 0$, then $f(c_i) = 0$
 so $\sum_{i=1}^n f(c_i)(0 - 0) = 0$ so $S(f, P) = 0$
 $0 < \epsilon$ ϵ was arbitrary so f is RI on $[0, 1]$

10 $\|P\| = 0.2$ so $\|P_0\| \leq \frac{0.2}{3}$ Have 18 partitions
 from -1 to 2 that are $\frac{0.2}{3}$ units long so that
 $\|P_0\| = \frac{0.2}{3}$.

5 $c = b - \epsilon$ $\|P_1\| < \delta$ $|S(f, P_1) - \int_a^b f| < \epsilon$ P_1 t.p. of $[a, c]$
 P_2 t.p. of $[a, b] = [a, c + \epsilon]$ $S(f, P_2) = S(f, P_1) + f(c)(c + \epsilon - c)$
 $S(f, P_2) = S(f, P_1) + f(c)\epsilon$
 $|S(f, P_2) - \int_a^b f| < \epsilon$ $|S(f, P_1) + \epsilon f(c) - \int_a^b f| \leq |S(f, P_1) - \int_a^b f| + \epsilon f(c)$
 $|S(f, P_1) - \int_a^b f| < \epsilon(1 + f(c))$
 Set ϵ in beginning to $\frac{\epsilon}{1 + f(c)}$
 ϵ arbitrary so f is RI on $[a, b]$.