


Chapter 16


Vector Calculus

16.1 Vector Fields

Examples.



Map retrieved from <http://www.pacioos.hawaii.edu/currents/model-oahu/>




(b) Airflow past an inclined airfoil

Vector Fields in 2D.

1 Definition Let D be a set in \mathbb{R}^2 (a plane region). A **vector field on \mathbb{R}^2** is a function \mathbf{F} that assigns to each point (x, y) in D a two-dimensional vector $\mathbf{F}(x, y)$.

Representation.



Component Functions


$$\vec{F}(x, y) = P(x, y)\vec{i} + Q(x, y)\vec{j}$$

- P : x -component of \vec{F}
- Q : y -component of \vec{F}

Remark:

- P and Q are real-valued functions.
- \vec{F} is continuous if P and Q are continuous.

EXAMPLE 1 A vector field on \mathbb{R}^2 is defined by $\mathbf{F}(x, y) = -y\mathbf{i} + x\mathbf{j}$. Describe \mathbf{F} by sketching some of the vectors $\mathbf{F}(x, y)$ as in Figure 3.



$$\vec{i} = \langle 1, 0 \rangle$$

$$\vec{j} = \langle 0, 1 \rangle$$

2	$\langle -2, 0 \rangle$	$\langle -2, 1 \rangle$	$\langle -2, 2 \rangle$
1	$\langle -1, 0 \rangle$	$\langle -1, 1 \rangle$	$\langle -1, 2 \rangle$
0	$\langle 0, 0 \rangle$	$\langle 0, 1 \rangle$	$\langle 0, 2 \rangle$
y/x	0	1	2


$$\vec{F}(0, 0) = \langle -y, x \rangle = \langle 0, 0 \rangle$$

$$\vec{F}(1, 0) = \langle 0, 1 \rangle$$

Vector Fields in 3D.

2 Definition Let E be a subset of \mathbb{R}^3 . A **vector field on \mathbb{R}^3** is a function \mathbf{F} that assigns to each point (x, y, z) in E a three-dimensional vector $\mathbf{F}(x, y, z)$.

Representation.




Component Functions. $\langle P, Q, R \rangle$

$$P(x, y, z)\vec{i} + Q(x, y, z)\vec{j} + R(x, y, z)\vec{k}$$

- P : x -component of \vec{F}
- Q : y -component of \vec{F}
- R : z -component of \vec{F}


\vec{F} is continuous if P, Q, R are continuous.

EXAMPLE 2 Sketch the vector field on \mathbb{R}^3 given by $\mathbf{F}(x, y, z) = z \mathbf{k} = \langle 0, 0, z \rangle$



on the plane $z = k$
 $\hookrightarrow \mathbf{F}(x, y, k) = \langle 0, 0, k \rangle$

(x, y, z)	$\vec{F}(x, y, z)$
$(0, 0, 1)$	$\langle 0, 0, 1 \rangle$
$(0, 1, 0)$	$\langle 0, 0, 0 \rangle$
$(1, 1, 1)$	$\langle 0, 0, 1 \rangle$
$(1, 0, 1)$	$\langle 0, 0, 1 \rangle$
$(1, 0, 2)$	$\langle 0, 0, 2 \rangle$
\vdots	\vdots



Remark:

A vector field is continuous if each of its component function (that is P, Q, R) are continuous.

EXAMPLE 4 Newton's Law of Gravitation tells you that the magnitude of the force of attraction between two objects of mass m and M is

$$F = \frac{mMG}{r^2}$$

where G is the gravitational constant, and r is the distance between the two objects. Find the vector field describing the gravitational field.

More Examples:

- Force field around an electric charge Q :

$$\vec{F}(\vec{x}) = \frac{\varepsilon_0 q Q}{\|\vec{x}\|^3} \vec{x}$$

- Electric Field around the charge Q :

$$\vec{E}(\vec{x}) = \frac{\vec{F}(\vec{x})}{q} = \frac{\varepsilon_0 Q}{\|\vec{x}\|^3} \vec{x}$$

Gradient Fields.

2D

$$\vec{\nabla} f(x, y) = f_x(x, y)\vec{i} + f_y(x, y)\vec{j}$$

3D

$$\vec{\nabla} f(x, y, z) = f_x(x, y, z)\vec{i} + f_y(x, y, z)\vec{j} + f_z(x, y, z)\vec{k}$$

EXAMPLE 6 Find the gradient vector field of $f(x, y) = x^2y - y^3$. Plot the gradient vector field together with a contour map of f . How are they related?

Conservative Vector Fields.

- A vector field \vec{F} is conservative if there is a scalar-valued function f such that

$$\vec{F} = \vec{\nabla} f$$

- The function f is called the potential function of \vec{F} .

EXAMPLE. Show that the Gravitational field is conservative.