

MATH 644

CHAPTER 2

SECTION 2.4: ANALYTIC FUNCTIONS

CONTENTS

Definition	2
Where Is A Power Series Analytic?	3
Uniqueness Of Power Series Expansion	5
Consequences On the Zeros	7

DEFINITION

We consider Ω to be an open subset of \mathbb{C} , meaning that

$$\forall z \in \Omega, \text{ there is an } r > 0 \text{ such that } \{w : |w - z| < r\} \subset \Omega.$$

DEFINITION 1. Let $f : \Omega \rightarrow \mathbb{C}$.

- f is **analytic** at $z_0 \in \Omega$ if there is an $r > 0$ and a power series $\sum_{n=0}^{\infty} a_n(z - z_0)^n$ converging in $B = \{z : |z - z_0| < r\}$ such that

$$f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n \quad (\forall z \in B).$$

- f is analytic on Ω if f is analytic at each $z_0 \in \Omega$.

Notes:

- The power series is not necessarily the same for each $z_0 \in \Omega$.
- A function f is analytic on a set E (not necessarily open), if there is an open set $\Omega \supset E$ and an analytic function g on Ω such that $f = g$.

THEOREM 2. If f is analytic in Ω , then f is continuous on Ω .

Proof.

Fact: (f_n) conv. unif. to some f & f_n is cont.
 \Downarrow
 f is cont.

Let $z_0 \in \Omega$. $\exists r > 0$ nt.

$$f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n \quad \forall z \in B.$$

From the root test, the power series conv. uniformly in a small enough disk centered at z_0 .
 So, partial sums converge uniformly to f , so f is continuous on $\{z : |z - z_0| \leq r/2\}$. \square

THEOREM 3. If $f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n$ converges on $\{z : |z - z_0| < r\}$, then f is analytic on $\{z : |z - z_0| < r\}$.

Proof. Fix z_1 st. $|z_1 - z_0| < r$. By the

Binomial Theorem:

$$(z - z_1 + z_1 - z_0)^n = \sum_{k=0}^n \binom{n}{k} (z - z_1)^k (z_1 - z_0)^{n-k}.$$

So,

$$f(z) = \sum_{n=0}^{\infty} a_n \left[\sum_{k=0}^n \binom{n}{k} (z - z_1)^k (z_1 - z_0)^{n-k} \right].$$

For now, suppose we can interchange the order of \sum . Then

$$f(z) = \sum_{k=0}^{\infty} \left[\sum_{n=k}^{\infty} a_n \binom{n}{k} (z_1 - z_0)^{n-k} \right] (z - z_1)^k.$$

From the root test, we know that

$$\sum |a_n| |w - z_0|^n \text{ conv. if } |w - z_0| < r$$

$$\text{Set } w := |z - z_1| + |z_1 - z_0| + z_0$$

$$\text{if } |z - z_0| < r - |z_1 - z_0|, \text{ then}$$

$$|w - z_0| = |z - z_1| + |z_1 - z_0| < r$$

&

$$\begin{aligned}
 \sum_{n=0}^{\infty} |a_n| |w - z_0|^n &= \sum_{n=0}^{\infty} |a_n| (|z - z_1| + |z_1 - z_0|)^n \\
 &= \sum_{n=0}^{\infty} |a_n| \left(\sum_{k=0}^n \binom{n}{k} |z - z_1|^k |z_1 - z_0|^{n-k} \right)
 \end{aligned}$$

Since the LHS is conv., then

$$\sum_{n=0}^{\infty} a_n \left(\sum_{k=0}^n \binom{n}{k} (z - z_1)^k (z_1 - z_0)^{n-k} \right) \text{ is abs. convergent. } \square$$

THEOREM 4. Suppose

$$\sum_{n=0}^{\infty} a_n(z - z_0)^n = \sum_{n=0}^{\infty} b_n(z - z_0)^n$$

for all complex numbers in $\{z : |z - z_0| < r\}$. Then $a_n = b_n$ for all $n \geq 0$.

Proof.

Write $c_n = b_n - a_n$. Then

$$\sum_{n=0}^{\infty} c_n(z - z_0)^n = 0 \quad (|z - z_0| < r).$$

Let c_m be the first non-zero coef ($m \geq 1$).

If $0 < |z - z_0| < r$, then

$$\begin{aligned} (z - z_0)^{-m} \sum_{n=m}^{\infty} c_n(z - z_0)^n &= \sum_{n=0}^{\infty} c_{n+m}(z - z_0)^n \\ &\equiv F(z). \end{aligned}$$

the series defining F is convergent on $\{z : 0 < |z - z_0| < r\}$. By the root test, the series for F converges in a disk containing z_0 .

Note:

- The proof actually shows also that if f is analytic at z_0 , then for some $\delta > 0$, either
 - $f(z) \neq 0$ for any $0 < |z - z_0| < \delta$;
 - or $f(z) = 0$ for any $|z - z_0| < \delta$.
- The proof also shows that if f is analytic at z_0 , then there is a $r > 0$, an integer $m \geq 1$ and an analytic function g at z_0 such that
 - $g(z) \neq 0$ for any z such that $|z - z_0| < r$;
 - $f(z) - f(z_0) = (z - z_0)^m g(z)$.

Consequences On the Zeros

- A set $\Omega \subset \mathbb{C}$ is called a **region** if it is
 - open;
 - connected, meaning that we can't write $\Omega = U \cup V$, where U and V are open sets in \mathbb{C} such that $V \cap U = \emptyset$.

Fact: Ω is connected if and only if Ω and \emptyset are the only open and closed subsets of Ω .

- A zero a of a function $f : \Omega \rightarrow \mathbb{C}$ is called **isolated** if there is an open disk B centered at a such that $f(z) \neq 0$ for any $z \in B \setminus \{a\}$.

COROLLARY 5. If f is analytic on a region Ω , then either $f \equiv 0$ or the zeros of f are isolated.

Proof.

Note:

- A consequence of the last Corollary is the **Identity principle**: If f and g are two analytic functions in a region Ω that agree on a set with an accumulation point in Ω , then they must be identical (see Problem 18).
- The last Corollary is not true for continuous functions: $f(x) = x \sin(1/x)$ is an example.