

MATH 307


CHAPTER 1

SECTION 1.1: SYSTEMS OF LINEAR EQUATIONS

| |
|----------|
| CONTENTS |
|----------|

| | |
|--|-----------|
| Why do we care about Systems of Linear Equations? | 2 |
| Terminology | 3 |
| Linear Equation | 3 |
| Systems of Linear Equations | 3 |
| Solution of a System of Linear Equations | 3 |
| How do we solve systems of linear equations? | 4 |
| Systems of two linear equations with two variables | 4 |
| Gauss-Jordan Elimination | 4 |
| Augmented Matrix | 6 |
| Elementary operations revisited | 6 |
| More Terminology | 9 |
| Reduced row-echelon form | 9 |
| Consistent Systems vs Inconsistent Systems | 9 |
| Homogeneous System | 9 |
| Gaussian Elimination | 10 |

WHY DO WE CARE ABOUT SYSTEMS OF LINEAR EQUATIONS?



Ohm's Law

- Voltage drop at a resistor is given by $V = IR$.

Kirchhoff's Laws

- Junction: Current flowing into a junction must flow out of it.
- Path: Sum of IR terms in any direction around a closed path is equal to the total voltage in the path in that direction.

Goal: Find the values of I_1, I_2, I_3

$$\begin{array}{l} J1) \quad I_1 + I_2 = I_3 \\ J2) \quad I_3 = I_1 + I_2 \end{array} \quad \bigg| \rightarrow \quad I_1 - I_2 - I_3 = 0$$

PATH. $P1) \quad 2I_1 + I_3 + 2I_1 = 8 \quad \rightarrow \quad 4I_1 + I_3 = 8$

$P2) \quad 4I_2 + I_3 = 16$

To find I_1, I_2, I_3 , we must solve the system of lin. eqs.:

$$\begin{cases} I_1 + I_2 - I_3 = 0 \\ 4I_1 + I_3 = 8 \\ 4I_2 + I_3 = 16 \end{cases}$$

Linear Equation

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = b$$

where

- a_1, a_2, \dots, a_n are constants.
- n is the number of variables.
- x_1, x_2, \dots, x_n are the variables (unknowns).
- b is the right-hand side constant term.

Systems of Linear Equations

$$\begin{array}{lcl}
 \text{no 1)} & a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\
 \text{no 2)} & a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\
 & \vdots \\
 \text{no m)} & a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m
 \end{array}$$

last example
 $n=3$
 $m=3$

where

- m is the number of linear equations.
- n is the number of variables.
- a_{11}, \dots, a_{mn} are constants.
- b_1, b_2, \dots, b_m are the right-hand side constant terms.
- x_1, \dots, x_n are the variables (unknowns).

Solution of a System of Linear Equations

A list $(x_1^*, x_2^*, \dots, x_n^*)$ is a solution to a system of linear equations if it satisfies each equation of the system.

Going back to our previous example

$(1, 3, 4)$ is a solution to our system in the last example.
 $\uparrow \quad \uparrow \quad \uparrow$
 $I_1 \quad I_2 \quad I_3$

$$\begin{array}{rclcl}
 1 & + & 3 & - & 4 & = & 0 & \checkmark \\
 4 \cdot 1 & + & 4 & & & = & 8 & \checkmark \\
 4 \cdot 3 & + & 4 & & & = & 16 & \checkmark
 \end{array}$$

Systems of two linear equations with two variables

$$\begin{aligned}x_1 + x_2 &= 0 \\ 2x_1 + x_2 &= 1.\end{aligned}$$

Method 1 (Isolate)

$$1) \quad x_2 = 1 - 2x_1$$

$$2) \quad x_1 + 1 - 2x_1 = 0$$

$$\rightarrow 1 - x_1 = 0$$

$$\rightarrow 1 = x_1$$

$$3) \quad x_2 = 1 - 2 \cdot 1 = -1$$

Solution: $\begin{pmatrix} 1 & -1 \end{pmatrix}$
 $\begin{matrix} \uparrow & \uparrow \\ x_1 & x_2 \end{matrix}$

Method 2 (Operations)

$$\begin{aligned}1) \quad x_1 + x_2 &= 0 & E_1 \\ 2x_1 + x_2 &= 1 & E_2\end{aligned}$$

$$\begin{aligned}2) \quad E_1 - E_2 &\rightarrow E_1 & \begin{array}{l} x_1 + x_2 = 0 \\ - (2x_1 + x_2 = 1) \\ \hline -x_1 + x_2 = -1 \end{array} \\ \begin{cases} -x_1 = -1 & E_1 \\ 2x_1 + x_2 = 1 & E_2 \end{cases} & \rightarrow \begin{array}{l} -x_1 = -1 \\ \rightarrow -x_1 = -1 \end{array}\end{aligned}$$

$$3) \quad -E_1 \rightarrow E_1$$

$$-(-x_1) = -1 \rightarrow x_1 = 1$$

$$\begin{cases} x_1 = 1 & E_1 \\ 2x_1 + x_2 = 1 & E_2 \end{cases}$$

$$4) \quad 2E_1 - E_2 \rightarrow E_2$$

$$\begin{array}{rcl} 2x_1 & = & 2 \\ - (2x_1 + x_2 = 1) & & \\ \hline -x_2 & = & 1 \end{array}$$

$$\begin{cases} x_1 = 1 & E_1 \\ -x_2 = 1 & E_2 \end{cases}$$

$$\begin{cases} x_1 = 1 \\ x_2 = -1 \end{cases}$$

Gauss-Jordan Elimination

Based on three *elementary operations* on the equations:

- Interchange two equations in the system.
- Replace an equation by a multiple of itself.
- Replace an equation by itself plus a multiple of another equation.

Main GOAL: transform our system into

$$\begin{aligned}x + 0y + 0z &= \tilde{b}_1 \\ 0x + y + 0z &= \tilde{b}_2 \\ 0x + 0y + z &= \tilde{b}_3.\end{aligned}$$

EXAMPLE 1. Find the solution(s) to the following system of linear equations:

$$\begin{aligned} 1x - y + z &= 0 \\ 2x - 3y + 4z &= -2 \\ -2x - y + z &= 7. \end{aligned}$$

E_1
 E_2
 E_3

$$\begin{array}{r} 1) \quad -2E_1 + E_2 \rightarrow E_2 \\ -2x + 2y - 2z = 0 \\ + \quad 2x - 3y + 4z = -2 \\ \hline -y + 2z = -2 \end{array}$$

$$\begin{array}{r} 2E_1 + E_3 \rightarrow E_3 \\ 2x - 2y + 2z = 0 \\ + \quad -2x - y + z = 7 \\ \hline -3y + 3z = 7 \end{array}$$

$$\begin{cases} x - y + z = 0 & E_1 \\ -y + 2z = -2 & E_2 \\ -3y + 3z = 7 & E_3 \end{cases}$$

$$\begin{array}{r} 2) \quad E_1 - E_2 \rightarrow E_1 \\ x - y + z = 0 \\ - \quad -y + 2z = -2 \\ \hline x - z = 2 \end{array}$$

$$\begin{array}{r} -3E_2 + E_3 \rightarrow E_3 \\ 3y - 6z = 6 \\ + \quad -3y + 3z = 7 \\ \hline -3z = 13 \end{array}$$

$$\begin{cases} x - z = 2 & E_1 \\ -y + 2z = -2 & E_2 \\ -3z = 13 & E_3 \end{cases}$$

$$\begin{array}{r} 3) \quad 2E_3 + 3E_2 \rightarrow E_2 \\ -6z = 26 \\ + \quad -3y + 6z = -6 \\ \hline -3y = 20 \end{array}$$

$$\begin{array}{r} 3E_1 - E_3 \rightarrow E_1 \\ 3x - 3z = 6 \\ - \quad -3z = 13 \\ \hline 3x = -7 \end{array}$$

$$\begin{cases} 3x = -7 \\ -3y = 20 \\ -3z = 13 \end{cases}$$

$$4) \quad \frac{3x}{3} = \frac{-7}{3} \rightarrow$$

$$\frac{-3y}{-3} = \frac{20}{-3} \rightarrow$$

$$\frac{-3z}{-3} = \frac{13}{-3} \rightarrow$$

$$\boxed{\begin{aligned} x &= -\frac{7}{3} \\ y &= -\frac{20}{3} \\ z &= -\frac{13}{3} \end{aligned}}$$

Augmented Matrix

More efficient way: transform the system in an **augmented matrix**.

$$\begin{array}{rcl}
 a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n & = & b_1 \\
 a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n & = & b_2 \\
 & \vdots & \\
 a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n & = & b_m
 \end{array}
 \Rightarrow
 \begin{array}{c}
 x_1 \quad x_2 \quad \cdots \quad x_n \quad b \\
 \left[\begin{array}{ccccc}
 a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\
 a_{21} & a_{22} & \cdots & a_{2n} & b_2 \\
 \vdots & \vdots & \vdots & \vdots & \vdots \\
 a_{m1} & a_{m2} & \cdots & a_{mn} & b_m
 \end{array} \right]
 \end{array}$$

EXAMPLE 2. Find the augmented matrix of the system of Example 1.

$$\begin{array}{rcl}
 1x - y + z & = & 0 \\
 2x - 3y + 4z & = & -2 \\
 -2x - y + z & = & 7
 \end{array}
 \Rightarrow
 \begin{array}{c}
 x \quad y \quad z \quad b \\
 \left[\begin{array}{cccc}
 1 & -1 & 1 & 0 \\
 2 & -3 & 4 & -2 \\
 -2 & -1 & 1 & 7
 \end{array} \right]
 \end{array}$$

Elementary operations revisited

Elementary operations on linear equations become elementary operations on the rows of the augmented matrix:

- Interchange two rows.
- Replace a row by a multiple of itself.
- Replace a row by itself plus a multiple of another row.

EXAMPLE 3. Solve the system:

$$2x + 3y - z = 3$$

$$-x - y + 3z = 0$$

$$x + 2y + 2z = 3$$

$$y + 5z = 3.$$

$$\begin{bmatrix} 2 & 3 & -1 & 3 \\ -1 & -1 & 3 & 0 \\ 1 & 2 & 2 & 3 \\ 0 & 1 & 5 & 3 \end{bmatrix} \sim \begin{bmatrix} 2 & 3 & -1 & 3 \\ \boxed{0} & 1 & 5 & 3 \\ 0 & -1 & -5 & -3 \\ 0 & 1 & 5 & 3 \end{bmatrix} \begin{array}{l} R_1 + 2R_2 \rightarrow R_2 \\ R_1 - 2R_3 \rightarrow R_3 \end{array}$$
$$\sim \begin{bmatrix} 2 & 0 & -16 & -6 \\ 0 & 1 & 5 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{array}{l} R_1 - 3R_2 \rightarrow R_1 \\ R_2 + R_3 \rightarrow R_3 \\ R_2 - R_4 \rightarrow R_4 \end{array}$$

EXAMPLE 4. Solve the system:

$$4x_1 - 8x_2 - x_3 + x_4 + 3x_5 = 0$$

$$5x_1 - 10x_2 - x_3 + 2x_4 + 3x_5 = 0$$

$$3x_1 - 6x_2 - x_3 + x_4 + 2x_5 = 0.$$

Reduced row-echelon form

Transformed augmented matrix after row operations:

- Any rows of zero (called zero rows) appear at the bottom.
- The first nonzero entry of a nonzero row is 1 (called a leading 1).
- The leading 1 of a nonzero row appears to the right of the leading 1 of any preceding row.
- All the other entries of a column containing a leading 1 are zero.

Consistent Systems vs Inconsistent Systems

- Consistent: means the system of equations has at least one solution.
 - How to recognize that a system is consistent?

(1)
(2)

- Inconsistent: means the system of equations has no solution.
 - How to recognize that a system is inconsistent?

(1)

Homogeneous System

$$\begin{aligned}
 a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n &= 0 \\
 a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n &= 0 \\
 &\vdots \\
 a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n &= 0
 \end{aligned}$$

- Trivial solution: $x_1 = x_2 = \cdots = x_n = 0$.

THEOREM 5. A homogeneous system of m linear equations in n variables has

- infinitely many nontrivial solutions if $m < n$;
- exactly one (trivial solution) if $m = n$;
- no solution if $m > n$.

GAUSSIAN ELIMINATION

Goal. Transform the augmented matrix into a new augmented matrix with the following properties:

- any zero rows appear at the bottom.
- The first nonzero entry of a nonzero row is 1.
- The leading 1 of a nonzero row appears to the right of the leading 1 of any preceding row.

EXAMPLE 6. Determine the values of a , b , and c so that the system

$$\begin{aligned}x - y + 2z &= a \\ 2x + y - z &= b \\ x + 2y - 3z &= c\end{aligned}$$

has solutions.