

Example 4

Does the function


$$f(x) = \begin{cases} 1/x & 0 < x \leq 1 \\ 0 & x = 0 \end{cases}$$

↓
↖ 2

have a maximum?

x	$f(x)$
0.5	2
0.25	4
0.10	10
0.01	100

x	$f(x)$
1	1
0.5	2
0.25	4
0.1	10
0.01	100



- One at $x = 1 \rightarrow f(x) = 1 \leq \frac{1}{x} \quad \forall x \leq 1, (x \neq 0)$
- Another one at $x = 0 \rightarrow f(0) = 0 \leq 1 \leq \frac{1}{x} \quad \forall x$

Example 7

Let $f(x) = x^3$. What is $f'(0)$? Is $f(0)$ a local maximum or local minimum?

$$f'(x) = 3x^2$$

The solution to $f'(x) = 0$

$$\Leftrightarrow 3x^2 = 0$$

$$\Leftrightarrow x = 0$$


It seems that $x=0$ is a local max or local min.

x	$f(x)$
2	8
1	1
0	0
-1	-1
-2	-8

x is not a maximum and not a minimum!

Example 8

Let $f(x) = |x|$. Is $f(0)$ a local maximum, global maximum, local minimum, or global minimum?



the $f'(0)$ \nexists

But $f(0) = 0$ is an absolute minimum.

x	$f(x)$
-2	2
-1	1
0	0
1	1
2	2

Example 10

Find the critical numbers of $f(x) = x^{3/5}(4 - x)$.

1) Find the derivative

$$f'(x) = (x^{3/5})'(4-x) + x^{3/5} \cdot (-1)$$

$$= \frac{3}{5} x^{-2/5} (4-x) - x^{3/5}$$

$$= \frac{3 \cdot 4}{5} x^{-2/5} - \frac{3}{5} x^{3/5} - x^{3/5}$$

$$= \frac{12}{5} x^{-2/5} - x^{3/5} \left(\frac{8}{5} \right)$$

$$\frac{12}{5x^{2/5}} - \frac{8x^{3/5}}{5} \leftarrow$$

$$= \frac{12 - 8x}{5x^{2/5}}$$

$$= \frac{4(3-2x)}{5x^{2/5}}$$

2) Solve $f'(x) = 0$

$$f'(x) = 0 \iff \frac{4(3-2x)}{5x^{2/5}} = 0$$

$$\iff x = \frac{3}{2}$$

3) Find $f'(x)$ when \nexists
 $f'(x) \nexists$ when $x = 0$

Answer
critical numbers
are $\frac{3}{2}$ & 0

Example 11

Find the absolute maximum and minimum values of the function

$f(x) = \underline{x^3 - 3x^2 + 1}$ on the interval $\underline{[-1/2, 4]}$.

1) Find the critical numbers.

$$f'(x) = 3x^2 - 6x$$

Solve $f'(x) = 0$ (for x).

$$\Leftrightarrow 3x^2 - 6x = 0$$

$$\Leftrightarrow (3x - 6)x = 0$$

$$\Leftrightarrow 3x - 6 = 0 \quad \text{or} \quad x = 0$$

$$\Leftrightarrow x = 2 \quad \text{or} \quad x = 0$$

We have $f(0) = 1$ & $f(2) = -3$.

2) Compute the values of f at endpoints.

$$a = -\frac{1}{2} \quad \& \quad b = 4$$

$$f(-1/2) = 1/8$$

$$f(4) = 17.$$

3) Take max & min

$$\max \{1, -3, 1/8, 17\} = 17$$

$$\min \{1, -3, 1/8, 17\} = -3$$

Answers: abs maximum is 17

abs minimum is -3

Example 12

Show that the equation $x^3 + x - 1 = 0$ has at least one root.

Here $f(x) = x^3 + x - 1$.

$$f(0) = 0 + 0 - 1 = -1$$

$$a = 0$$

$$f(1) = 1$$

$$b = 1$$

So, $f(0) < 0$ and $f(1) > 0$.

So, $\exists c \in (0, 1)$ s.t. $f(c) = 0$.

In other words, $c^3 + c - 1 = 0$.

Example 13

$$f(x) = x^3 + x - 1.$$

Show that the equation $x^3 + x - 1 = 0$ has exactly one root.

Suppose that there are two different roots a & b ($a \neq b$).

$$\text{So, } f(a) = 0 = f(b).$$

- f is cont. on $[a, b]$
- f is diff. on (a, b) .

So, by Rolle's theorem, $f'(c) = 0$ for some $c \in (a, b)$.
 \uparrow contradiction.

$$f'(x) = 3x^2 + 1 \geq 1, \text{ so } f'(x) \neq 0.$$

So, we must conclude that there is only one root.

Example 14

Consider $f(x) = x^2$.

$$x^2 = 0^2$$

$$x^2 = 2^2$$

- Find the slope of the secant line passing through $Q = (0, 0)$ and $P = (2, 4)$.
- Can you find a tangent line to $y = x^2$ with the same slope?

- the slope of PQ

$$m_{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - 0}{2 - 0} = 2$$

- Here $f(x) = x^2$. We have $f'(x) = 2x$

So,

$$\begin{aligned} f'(x) &= 2 & \Leftrightarrow & 2x = 2 \\ & & \Leftrightarrow & \boxed{x = 1} \end{aligned}$$

Example 15

Find the numbers $c \in [0, 2]$ such that the average of the function $f(x) = x^3 - x$ on the interval $[0, 2]$ is attained by $f'(c)$.

Here, $a = 0$, $b = 2$.

$$f(x) = x^3 - x \rightarrow$$

- continuous on $[0, 2]$
- differentiable on $(0, 2)$.

So, $\exists c \in (0, 2)$ s.t.

$$f'(c) = \frac{f(2) - f(0)}{2 - 0}$$

$$f'(x) = 3x^2 - 1$$

$$\frac{f(2) - f(0)}{2 - 0} = \frac{6 - 0}{2} = 3$$

So,

$$3c^2 - 1 = 3$$

$$\Rightarrow 3c^2 = 4$$

$$\Rightarrow \sqrt{c^2} = \sqrt{4/3}$$

$$\Rightarrow |c| = \sqrt{4/3}$$

$$\Rightarrow c = +\sqrt{4/3} \text{ or } -\sqrt{4/3}$$

The slope of the tangent line is the same as the slope of the secant if

$$c = 2/\sqrt{3}$$

or

$$c =$$

$$-2/\sqrt{3}$$

does not
to $[0, 2]$.