

MATH 644

CHAPTER 2

SECTION 2.5: ELEMENTARY OPERATIONS

CONTENTS

Linear combinaisons	2
Multiplication	3
Composition	4
Complex Derivatives	6

THEOREM 1. Let f and g be analytic at z_0 . Then,

- a) $f + g$ is analytic at z_0 ;
- b) $f - g$ is analytic at z_0 ;
- c) cf is analytic at z_0 , for any $c \in \mathbb{C}$.

Proof.

$$(a) \text{ Analytic at } z_0 \Rightarrow \begin{aligned} f(z) &= \sum_{n=0}^{\infty} a_n (z - z_0)^n, \quad |z - z_0| < r_1 \\ &\& g(z) = \sum_{n=0}^{\infty} b_n (z - z_0)^n, \quad |z - z_0| < r_2 \end{aligned}$$

So, for $r := \min \{r_1, r_2\}$, we have

$$f(z) + g(z) = \sum_{n=0}^{\infty} (a_n + b_n) (z - z_0)^n, \quad |z - z_0| < r.$$

So, $f + g$ is analytic at z_0 .

Repeat for (b) & (c).

□

- If $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are absolutely convergent series with sums A and B respectively, then their Cauchy Product

$$\left(\sum_{n=1}^{\infty} a_n \right) \left(\sum_{n=1}^{\infty} b_n \right) = \sum_{n=1}^{\infty} \left(\sum_{m=1}^{n-1} a_m b_{n-m} \right)$$

converges absolutely to AB . [See Problem]

THEOREM 2. Let f and g be two analytic functions at z_0 . Then, the function $h = fg$ is analytic at z_0 .

Proof.

Write, for $0 < r < \infty$ small enough,

$$f(z) = \sum_{n=0}^{\infty} a_n (z-z_0)^n \quad \& \quad g(z) = \sum_{n=0}^{\infty} b_n (z-z_0)^n$$

on $\{z: |z-z_0| < r\}$.

Since the power series of f & g converge absolutely on $\{z: |z-z_0| < r\}$, we have

$$\begin{aligned} f(z)g(z) &= \sum_{n=0}^{\infty} \sum_{m=0}^n a_m b_{n-m} (z-z_0)^n \\ &= \sum_{n=0}^{\infty} \left(\sum_{m=0}^n a_m b_{n-m} \right) (z-z_0)^n \\ &= \sum_{n=0}^{\infty} c_n (z-z_0)^n \end{aligned}$$

converges in $\{z: |z-z_0| < r\}$. So $h = fg$ is analytic at z_0 . □

THEOREM 3. If f is analytic at z_0 and g is analytic at $a_0 = f(z_0)$, then the function $h = g \circ f$ is analytic at z_0 .

Proof.

$$\text{Suppose } f(z) = \sum_{n=0}^{\infty} a_n (z-z_0)^n, \quad |z-z_0| < r$$

$$\& \quad g(z) = \sum_{n=0}^{\infty} b_n (z-a_0)^n, \quad |z-a_0| < \rho.$$

The series

$$(*) \quad \sum_{m=1}^{\infty} |a_m| |z-z_0|^{m-1}$$

converges in $\{z: 0 < |z-z_0| < r\}$. By the

root test, the power series in $(*)$ conv.

uniformly on $\{z: |z-z_0| \leq r_1\}$, $r_1 < r$.

Therefore, it is bounded:

$$\exists M > 0, \quad \sum_{m=1}^{\infty} |a_m| |z-z_0|^m \leq M |z-z_0|$$

for $|z-z_0| \leq r_1$. Therefore,

$$\sum_{n=0}^{\infty} |b_n| \left(\sum_{m=1}^{\infty} |a_m| |z-z_0|^m \right)^n \leq \sum_{n=0}^{\infty} |b_n| M^n |z-z_0|^n$$

and the RHS converges if

$$|z-z_0| < \min \left\{ \frac{r_1}{M}, \frac{\rho}{M} \right\}.$$

So,

$$\sum_{n=0}^{\infty} b_n \left(\sum_{m=0}^{\infty} a_m (z-z_0)^m - a_0 \right)$$

conv. abs. in $|z-z_0| < \min \left\{ \frac{r_1}{M}, \frac{\rho}{M} \right\}$.

We can therefore rearrange the doubly indexed series so that

$$h(z) = \sum_{n=0}^{\infty} c_n (z-z_0)^n, \quad |z-z_0| < \min \left\{ \frac{r_1}{M}, \frac{\rho}{M} \right\}.$$

□

Consequences:

- If f is analytic at z_0 with $f(z_0) \neq 0$, then $1/f$ is analytic at z_0 .
- If $r = p/q$ is a rational function, then r is analytic on $\{z : q(z) \neq 0\}$.

DEFINITION 4. If f is defined in a disk (neighborhood) of z , then

$$f'(z) := \lim_{w \rightarrow z} \frac{f(w) - f(z)}{w - z}$$

is called the **(complex) derivative** of f , provided the limit exists.

Note:

- The function $f(z) = \bar{z}$ does not have a complex derivative.
- If n is a non-negative integer, then

$$(z^n)' = nz^{n-1}.$$

THEOREM 5. If $f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n$ converges in $B = \{z : |z - z_0| < r\}$, then

a) $f'(z)$ exists for all $z \in B$ and

$$f'(z) = \sum_{n=1}^{\infty} na_n(z - z_0)^{n-1} = \sum_{n=0}^{\infty} (n+1)a_{n+1}(z - z_0)^n \quad (\forall z \in B).$$

b) Moreover, the series for f' based at z_0 has the same radius of convergence as the series for f .

Proof.

Note: The rules of differentiation hold:

- $(f + g)'(z) = f'(z) + g'(z);$
- $(cf)'(z) = cf'(z);$
- $(fg)'(z) = f'(z)g(z) + f(z)g'(z);$
- $(\frac{f}{g})'(z) = (f'(z)g(z) - f(z)g'(z))/(g(z))^2;$
- $(g \circ f)(z) = g'(f(z))f'(z).$

COROLLARY 6. An analytic function f has derivatives of all orders. Moreover, if f is equal to a convergent power series on $B = \{z : |z - z_0| < r\}$, then the power series is given by

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n \quad (\forall z \in B).$$

Proof.

Consequences:

- If f is analytic in a region Ω with $f'(z) = 0$ for all z in a neighborhood of $z_0 \in \Omega$, then f is constant in Ω .
- If f and g are analytic in a region Ω with $f' = g'$, then $f - g$ is constant.
- If $f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n$ converges in $B = \{z : |z - z_0| < r\}$, then the power series

$$F(z) := \sum_{n=0}^{\infty} \frac{a_n}{n+1} (z - z_0)^{n+1}$$

converges in B and satisfies $F'(z) = f(z)$ for all $z \in B$.

COROLLARY 7. If $f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n$ converges in $B = \{z : |z - z_0| < r\}$, then

$$f'(z_0) = \lim_{z, w \rightarrow z_0} \frac{f(z) - f(w)}{z - w}.$$

Proof.