


15.6 Triple Integrals.



3 Definition The **triple integral** of f over the box B is

$$\iiint_B f(x, y, z) dV = \lim_{l, m, n \rightarrow \infty} \sum_{i=1}^l \sum_{j=1}^m \sum_{k=1}^n f(x_{ijk}^*, y_{ijk}^*, z_{ijk}^*) \Delta V$$

if this limit exists.

4 Fubini's Theorem for Triple Integrals If f is continuous on the rectangular box $B = [a, b] \times [c, d] \times [r, s]$, then

$$\iiint_B f(x, y, z) dV = \int_r^s \int_c^d \int_a^b f(x, y, z) dx dy dz$$


EXAMPLE 1 Evaluate the triple integral $\iiint_B xyz^2 dV$, where B is the rectangular box given by

$$B = \{(x, y, z) \mid 0 \leq x \leq 1, -1 \leq y \leq 2, 0 \leq z \leq 3\}$$

General Domains.


$$\iiint_E f(x, y, z) dV = \iiint_B F(x, y, z) dV$$

Domain of type 1.




$$\iiint_E f(x, y, z) dV = \iint_D \left[\int_{u_1(x, y)}^{u_2(x, y)} f(x, y, z) dz \right] dA$$

Projection D is of Type I.



$$\iiint_E f(x, y, z) dV = \int_a^b \int_{g_1(x)}^{g_2(x)} \int_{u_1(x, y)}^{u_2(x, y)} f(x, y, z) dz dy dx$$


Projection D is of Type II.



$$\iiint_E f(x, y, z) dV = \int_c^d \int_{h_1(y)}^{h_2(y)} \int_{u_1(x, y)}^{u_2(x, y)} f(x, y, z) dz dx dy$$

EXAMPLE 2 Evaluate $\iiint_E z \, dV$, where E is the solid tetrahedron bounded by the four planes $x = 0$, $y = 0$, $z = 0$, and $x + y + z = 1$.

Domains of type 2.



$$\iiint_E f(x, y, z) dV = \iint_D \left[\int_{u_1(y, z)}^{u_2(y, z)} f(x, y, z) dx \right] dA$$


Projection D is of Type I.

Projection D is of Type II.

$$\iiint_E f(x, y, z) dV = \int_a^b \int_{z_1(y)}^{z_2(y)} \int_{u_1(y, z)}^{u_2(y, z)} f(x, y, z) dx dz dy$$

$$\iiint_E f(x, y, z) dV = \int_a^b \int_{y_1(z)}^{y_2(z)} \int_{u_1(y, z)}^{u_2(y, z)} f(x, y, z) dx dy dz$$

Domains of type 3.



$$\iiint_E f(x, y, z) dV = \iint_D \left[\int_{u_1(x, z)}^{u_2(x, z)} f(x, y, z) dy \right] dA$$

Projection D is of type I.

Projection D is of type II.

$$\iiint_E f(x, y, z) dV = \int_a^b \int_{z_1(x)}^{z_2(x)} \int_{u_1(x, z)}^{u_2(x, z)} f(x, y, z) dy dz dx$$

$$\iiint_E f(x, y, z) dV = \int_a^b \int_{x_1(z)}^{x_2(z)} \int_{u_1(x, z)}^{u_2(x, z)} f(x, y, z) dy dx dz$$

EXAMPLE 3 Evaluate $\iiint_E \sqrt{x^2 + z^2} \, dV$, where E is the region bounded by the paraboloid $y = x^2 + z^2$ and the plane $y = 4$.

EXAMPLE 4 Express the iterated integral $\int_0^1 \int_0^{x^2} \int_0^y f(x, y, z) \, dz \, dy \, dx$ as a triple integral and then rewrite it as an iterated integral in a different order, integrating first with respect to x , then z , and then y .

Application: computing volumes of solids.

$$V(E) = \iiint_E dV$$

EXAMPLE 5 Use a triple integral to find the volume of the tetrahedron T bounded by the planes $x + 2y + z = 2$, $x = 2y$, $x = 0$, and $z = 0$.

Other applications.

Mass.

Moments.

Moments of Inertia.

EXAMPLE 6 Find the center of mass of a solid of constant density that is bounded by the parabolic cylinder $x = y^2$ and the planes $x = z$, $z = 0$, and $x = 1$.