
Chapter 1 Functions and Limits 1.8 Continuity

Continuity

Example. What are the main difference(s) between the two following curves?

Illustration: https://www.desmos.com/calculator/hflxgbsemz



Example. Now, what are the differences between the two following functions?

(a)
$$f(x) = \begin{cases} 2 - x & \text{if } -2 \le x < 1 \\ 0 & \text{if } 1 \le x \le 2 \end{cases}$$

(a)
$$f(x) = \begin{cases} 2-x & \text{if } -2 \le x < 1 \\ 0 & \text{if } 1 \le x \le 2 \end{cases}$$
 (b) $g(x) = \begin{cases} \frac{4}{9}(1-x)^2 & \text{if } -2 \le x < 1 \\ 0 & \text{if } 1 \le x \le 2 \end{cases}$

1 Definition A function f is **continuous at a number** a if


$$\lim_{x \to a} f(x) = f(a)$$

Three things to verify to show a function is continuous:

- a) The function is defined at x = a.
- b) The limit of the function exists at x = a.
- c) The limit of the function at x = a equals the value of the function at x = a.

Discontinuity:

EXAMPLE 1 Figure 2 shows the graph of a function f. At which numbers is f discontinuous? Why?




Example. Check if the functions in the first example are continuous at x=1 using the formulas.


(a)
$$f(x) = \frac{x^2 - x - 2}{x - 2}$$


(a)
$$f(x) = \frac{x^2 - x - 2}{x - 2}$$
 (b) $f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$ (c) $f(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x \leq 0 \end{cases}$

(c)
$$f(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x \le 0 \end{cases}$$

3 kinds of discontinuity.







(c) Jump discontinuity.

Definition A function f is **continuous from the right at a number a** if

$$\lim_{x \to a^+} f(x) = f(a)$$

and f is **continuous from the left at** a if

$$\lim_{x \to a^{-}} f(x) = f(a)$$

EXAMPLE. Is the function

$$f(x) = \begin{cases} 1 & \text{, if } x > 0 \\ 0 & \text{, if } x \le 0 \end{cases}$$

(a) continuous from the right at x = 0 (b) continuous from the left at x = 0.

Properties of Continuous Functions.

4 Theorem If f and g are continuous at a and if c is a constant, then the following functions are also continuous at a:

1.
$$f + g$$

2.
$$f - g$$

5.
$$\frac{f}{g}$$
 if $g(a) \neq 0$

Consequences:

7 Theorem The following types of functions are continuous at every number in their domains:

- polynomials
- rational functions
- root functions
- trigonometric functions

Substitution Rule Revisited.

EXAMPLE 5 Find $\lim_{x \to -2} \frac{x^3 + 2x^2 - 1}{5 - 3x}$.

EXAMPLE 7 Evaluate $\lim_{x \to \pi} \frac{\sin x}{2 + \cos x}$.

Theorem If f is continuous at b and $\lim_{x \to a} g(x) = b$, then $\lim_{x \to a} f(g(x)) = f(b)$. In other words,

$$\lim_{x \to a} f(g(x)) = f\left(\lim_{x \to a} g(x)\right)$$

9 Theorem If g is continuous at a and f is continuous at g(a), then the composition f(g(x)) is continuous at a.

Example. Find the value of


$$\lim_{x \to 1/2} \sin(\pi - \pi x^2)$$

EXAMPLE. Suppose we have a function


$$f(x) = x^2 - 1.$$

Does the graph of the function f cross the horizontal line y = 3?

10 The Intermediate Value Theorem Suppose that f is continuous on the closed interval [a, b] and let N be any number between f(a) and f(b), where $f(a) \neq f(b)$. Then there exists a number c in (a, b) such that f(c) = N.



(a) Find one number c



(b) Find multiple numbers c

EXAMPLE 9 Show that there is a root of the equation

$$4x^3 - 6x^2 + 3x - 2 = 0$$

between 1 and 2.