

Chapter 2

Derivatives

2.3 Differentiation Formulas

Constant Function.

$$f(x) = c \Rightarrow \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{c - c}{h} = 0$$

Derivative of a Constant Function

$$\frac{d}{dx}(c) = 0$$

Power Functions.

$n = 1.$ $f(x) = x^1$

$$\lim_{h \rightarrow 0} \frac{x+h - x}{h} = 1 \Rightarrow \frac{d}{dx}(x) = 1$$

$n = 2.$ $f(x) = x^2$

$$\lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h} = \lim_{h \rightarrow 0} \frac{2x\cancel{h} + \cancel{h}^2}{\cancel{h}} = 2x$$

$$\Rightarrow \frac{d}{dx}(x^2) = 2x$$

The Power Rule If n is a positive integer, then

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

Derivatives rules: Constant multiple, Sum and Difference

EXAMPLE. Find the derivative of $y = x^4 - 6x^2 + 4$.

Multiplication by a constant.

The Constant Multiple Rule If c is a constant and f is a differentiable function, then

$$\frac{d}{dx}[cf(x)] = c \frac{d}{dx}f(x)$$

Sum.

The Sum Rule If f and g are both differentiable, then

$$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x)$$

Difference.

The Difference Rule If f and g are both differentiable, then

$$\frac{d}{dx}[f(x) - g(x)] = \frac{d}{dx}f(x) - \frac{d}{dx}g(x)$$

Product.

The Product Rule If f and g are both differentiable, then

$$\frac{d}{dx}[f(x)g(x)] = f(x) \frac{d}{dx}[g(x)] + g(x) \frac{d}{dx}[f(x)]$$

Caution!!!

$$\frac{d}{dx}(fg) \neq \frac{d}{dx}(f) \frac{d}{dx}(g).$$

Example.



Example. Find the derivative of the function $f(x) = (5x^2 - 2)(x^3 + 3x)$.

Quotient.

The Quotient Rule If f and g are differentiable, then

$$\frac{d}{dx} \left[\frac{f(x)}{g(x)} \right] = \frac{g(x) \frac{d}{dx} [f(x)] - f(x) \frac{d}{dx} [g(x)]}{[g(x)]^2}$$

Caution !!

$$\frac{d}{dx} \left(\frac{f}{g} \right) \neq \frac{\frac{d}{dx}(f)}{\frac{d}{dx}(g)}$$

Example.

EXAMPLE 8 Let $y = \frac{x^2 + x - 2}{x^3 + 6}$. Compute the derivative.

General Power rule.

The Power Rule (General Version) If n is any real number, then

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

Case $n = 0$:

Example. Find the derivative of the function $f(x) = x^{2/3}$.

EXAMPLE 13 At what points on the hyperbola $xy = 12$ is the tangent line parallel to the line $3x + y = 0$?

Summary of Differentiation Formulas.

Table of Differentiation Formulas

$\frac{d}{dx}(c) = 0$	$\frac{d}{dx}(x^n) = nx^{n-1}$	
$(cf)' = cf'$	$(f + g)' = f' + g'$	$(f - g)' = f' - g'$
$(fg)' = fg' + gf'$	$\left(\frac{f}{g}\right)' = \frac{gf' - fg'}{g^2}$	