

MATH 644

CHAPTER 2

SECTION 2.3: POWER SERIES

CONTENTS

A First Example	2
Definition	3
Tests	4

A FIRST EXAMPLE

More complicated functions are found by taking limits of polynomials.

EXAMPLE 1. Study the series $\sum_{n=0}^{\infty} z^n$ for a fixed $z \in \mathbb{C}$.

Fix $z \in \mathbb{C}$.

1) If $|z| \geq 1$, then $z^n \not\rightarrow 0$ and
therefore $\sum_{n=0}^{\infty} z^n$ diverges.

2) If $|z| < 1$. We have

$$S_n = \sum_{k=0}^n z^k = \frac{1 - z^{n+1}}{1 - z}$$

$$\Rightarrow \left| S_n - \frac{1}{1-z} \right| = \frac{|z|^{n+1}}{|1-z|} \rightarrow 0 \quad (n \rightarrow \infty)$$

So,

$$\sum z^n = \frac{1}{1-z} \quad (|z| < 1)$$

Power series $(|z| < 1)$

function defined on $\mathbb{C} \setminus \{1\}$.

DEFINITION

A formal power series

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n = a_0 + a_1(z - z_0) + a_2(z - z_0)^2 + \dots$$

is called a **convergent power series centered (or based) at z_0** if

$$\exists r > 0 \text{ s.t. } \sum_{n=0}^{\infty} a_n (z - z_0)^n \text{ conv. in } \{z: |z - z_0| < r\}.$$

Convention: $f(z_0) = a_0$ (to avoid 0^0).

EXAMPLE 2. Find a convergent power series centered at $z_0 \neq a$ representing $\frac{1}{z-a}$.

Write


$$\frac{1}{z-a} = \frac{1}{z-z_0 + z_0-a} = \frac{1}{-(a-z_0)} \cdot \frac{1}{1 - \frac{z-z_0}{a-z_0}} = f(z)$$

$$\text{If } \left| \frac{z-z_0}{a-z_0} \right| < 1, \quad \rightarrow \quad |z-z_0| < |a-z_0|$$

$$\frac{1}{1 - \frac{z-z_0}{a-z_0}} = \sum_{n=0}^{\infty} \frac{(z-z_0)^n}{(a-z_0)^n}$$

$$\Rightarrow f(z) = \sum_{n=0}^{\infty} \frac{-(z-z_0)^n}{(a-z_0)^{n+1}} \quad (*)$$

Note



- (*) conv. in $|z-z_0| < |a-z_0|$
Biggest disk in domain of $\frac{1}{z-a}$
- (*) div. in $|z-z_0| \geq |a-z_0|$.

THEOREM 3. Let $r > 0$ and suppose that

a) $|a_n(z - z_0)^n| \leq M_n$ for every z such that $|z - z_0| \leq r$;

b) $\sum_{n=0}^{\infty} M_n < \infty$.

Then $f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n$ converges uniformly and absolutely in the region

$$D_r := \{z : |z - z_0| \leq r\}.$$

Proof.

Abs conv. Let $z \in D_r$. & $A_N = \sum_{k=0}^N |a_k| |z - z_0|^k$.

Then,

$$A_N \leq \sum_{k=0}^N M_k \leq \sum_{k=0}^{\infty} M_k < \infty$$

$$\Rightarrow (A_N)_{N=0}^{\infty} \text{ converges.}$$

Unif conv. Let $S_N = \sum_{k=0}^N a_k (z - z_0)^k$

We will show that (S_N) is unif. Cauchy.

For $z \in D_r$, & $N > M$,

$$|S_N - S_M| = \left| \sum_{k=M+1}^N a_k (z - z_0)^k \right| \leq \sum_{k=M+1}^N M_k.$$

$$\exists K = K(\varepsilon) \text{ s.t. } \forall N > M \geq K, \sum_{j=M+1}^N M_j < \varepsilon. \text{ So,}$$

(S_N) is uniformly Cauchy in D_r . \square

Note:

Convergence depends only on the tail of the series. So it is sufficient to satisfy the hypothesis only for $n \geq n_0$, for some non-negative integer n_0 .


THEOREM 4. Let $\sum_{n=0}^{\infty} a_n(z - z_0)^n$ be a formal power series. Let

$$R := \liminf_{n \rightarrow \infty} |a_n|^{-1/n} = \frac{1}{\limsup_{n \rightarrow \infty} |a_n|^{1/n}} \in [0, \infty].$$

Then, the power series

- a) converges abs. in $\{z : |z - z_0| < R\}$;
- b) converges uniformly in $\{z : |z - z_0| \leq r\}$, for any $r < R$;
- c) diverges in $\{z : |z - z_0| > R\}$.

Notes:



- R is called the radius of convergence of the power series.
- Biggest open disk where the power series converges is $\{z : |z - z_0| < R\}$.
- Information on the decay rate of a_n : for any $S < R$, there is an $n_0 \geq 0$ such that $|a_n| \leq S^{-n}$.

Proof.

Note: Root test does not give any information on the convergence of the power series on the circle

$$\{z : |z - z_0| = R\}.$$

EXAMPLE 5. Find the Radius of convergence R of the following power series and study their behavior on the boundary of the disk of radius R .

A) $\sum_{n=1}^{\infty} \frac{z^n}{n};$

C) $\sum_{n=1}^{\infty} n z^n;$

B) $\sum_{n=1}^{\infty} \frac{z^n}{n^2};$

D) $\sum_{n=1}^{\infty} 2^{n^2} z^n.$

EXAMPLE 6. Let $(a_n)_{n=0}^{\infty}$ be defined by

$$a_n = \begin{cases} 3^{-n} & , \text{ if } n \text{ is even} \\ 4^n & , \text{ if } n \text{ is odd.} \end{cases}$$

What is the radius of convergence of the power series $\sum_{n=0}^{\infty} a_n(z - z_0)^n$.