

Next Silicon: CM Home Assignment

Djordje Zivanovic

LinScale

April 24, 2025

Contents

1	Introduction	2
2	The Existing Implementation: Code Analysis and Documentation	2
2.1	Code Drawbacks	2
2.2	Numerical and Implementation Drawbacks	2
2.3	Mathematical Analysis	4
2.4	Accuracy and Correctness Failures	4
2.5	Experiments	5
3	Additional Algorithms	5

1 Introduction

This report ¹ contains the responses to the tasks stated in the home project pdf. It is accompanied with the repository ² that contains reproducible solutions with the installation instructions alongside: experiments and tests according to the task requirements.

2 The Existing Implementation: Code Analysis and Documentation

In this section we analyze the existing code shown in Algorithm 1.

2.1 Code Drawbacks

The code is written for C and follows the following bad practices:

1. Reusing the variable multiple times, `(float)M_PI` and `2.0f * (float)M_PI`, (lines 3, 4, 6, 6, 5, 7 of Algorithm 1);
2. Not using `auto` in order to automatically deduce types since the results of all the statements are known;
3. Cleaning if loop to be more understandable: `fmodf` returns the result in the range $(-2\pi, 2\pi)$. Then, now it is obvious that one checks whether the number is outside of the range $[-\pi, \pi]$, and then updates `x` for 2π period, so the further code operates with the number in the range $[-\pi, \pi]$;
4. Adding more verbosity `()`;
5. Reusing variable names -> more verbose names should be used in order to improve readability of the code. The c ompiler will optimize for the least number of variables/registers to be used;
6. Renaming function names and migrating these functions to the corresponding headers and sources that would contain the custom maths functions.

2.2 Numerical and Implementation Drawbacks

Here, I will give state several main drawbacks in terms of the implementation and numerical accuracy. The division by In the next subsection, I will list the drawbacks related to the method itself.

¹report

²next-silicon-maths

- 1: **Input:** A float (IEEE-754) number
- 2: **Output:** A float (IEEE-754) sine value of this number computed using Taylor Series.
- 3: **Steps:**

```
1 float fp32_custom_sine(float x)
2 {
3     x = fmodf(x, 2.0f * (float)M_PI);
4     if (x > (float)M_PI)
5         x -= 2.0f * (float)M_PI;
6     else if (x < -(float)M_PI)
7         x += 2.0f * (float)M_PI;
8     float result = 0.0f;
9     float term = x;
10    float x_squared = x * x;
11    int sign = 1;
12    for (int n = 1; n <= 7; n += 2)
13    {
14        result += sign * term;
15        sign = -sign;
16        term = term * x_squared;
17        term = term / (float)(n + 1);
18        term = term / (float)(n + 2);
19    }
20    return result;
21 }
```

Algorithm 1: Algorithm with Code Listing

2.3 Mathematical Analysis

Let us state the general Taylor series formula that is implemented in Algorithm 1.

Theorem 2.1 (Theorem 5.19 from [1, p. 113]): Let f be a function having finite n -th derivative $f^{(n)}$ everywhere in an open interval (a, b) and assume that $f^{(n-1)}$ is continuous on the closed interval $[a, b]$. Then, for every x in $[a, b]$, $x \neq c$, there exists a point x_1 interior to the interval joining x and c such that

$$f(x) = f(c) + \sum_{k=1}^{n-1} \frac{f^{(k)}(c)(x-c)^{(k)}}{k!} + \frac{f^{(n)}(x_1)}{n!}(x-c)^n.$$

A corollary of Theorem 2.1 when we set $c = 0$ is a Maclaurin Series.

Corollary 2.2 (Maclaurin Series): Let f be a function having finite n -th derivative $f^{(n)}$ everywhere in an open interval (a, b) and assume that $f^{(n-1)}$ is continuous on the closed interval $[a, b]$. Assume that $c \in [a, b]$. Then, for every x in $[a, b]$, $x \neq 0$, there exists a point x_1 interior to the interval joining x and 0 such that

$$f(x) = f(0) + \sum_{k=1}^{n-1} \frac{f^{(k)}(0)(x)^{(k)}}{k!} + \frac{f^{(n)}(x_1)}{n!}x^n.$$

Let us now set $a = -\pi$, $b = \pi$ and $f(x) = \sin(x)$. The Maclaurin Series becomes for sin function:

Corollary 2.3: For $x \in \mathbb{R}$, $x \neq 0$ and $n \in \mathbb{N}$, and $0 < |x_1| < |x|$ the approximation of a degree n is

$$\sin(x) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^i \frac{x^{2i+1}}{(2i+1)!} + L(x, n).$$

where

$$L(x, n) = \begin{cases} \frac{f^{(n+2)}(x_1)}{(n+2)!}x^{n+2}, & \text{if } n \bmod 2 = 0, \\ \frac{f^{(n+1)}(x_1)}{(n+1)!}x^{n+1}, & \text{if } n \bmod 2 = 1. \end{cases}$$

Thus, the method stated in Algorithm 1 has an algorithmic error $L(x, 7)$ which is smaller than $\frac{x^8}{8!}$. Now, we obtain

2.4 Accuracy and Correctness Failures

Based on Corollary 2.3, we can notice that Algorithm 1 is incorrect for $x = 0$. Similar, the farther the x is from 0, the error is larger since x^8 grows exponentially. There are several ways to solve these problems:

2.5 Experiments

The test plan will cover the following experiments:

1. Different number distributions
 - (a) Equally distanced numbers between certain multiplicands of $\frac{\pi}{2}$; We will vary the distance and provide statistical analysis with plots for each of such distance and multiplicand.
 - (b) Normally distributed numbers around certain multiplicands of $\frac{\pi}{2}$; We will vary the variance and provide statistical analysis with plots for each of the multiplicands and variances.
2. Edge cases
 - (a) the multiplicands of $\frac{\pi}{2}$;
 - (b) large numbers (close to the absolute minimum and maximum for the float numbers);
 - (c) normal distributions around the multiplicands of $\frac{\pi}{2}$;
 - (d) `nans`.
3. Numbers

For each of these experiments we would compute the relative error in comparison to the value computed by the `sin x` provided in `cmath`.

3 Additional Algorithms

For the problems stated in Subsection 2.4 there are several ways to approach:

1. Have another Taylor series expansion for numbers around π and $-\pi$. In this case we would manually calculate the `sin x` for π and $-\pi$.
2. Add more terms in Taylor series expansion. It is important to find the optimal degree for the balance of accuracy and performance.
3. Implement one or more of the alternative methods: Minimax Polynomial Approximation, Chebyshev Polynomial Expansion, Lookup Table with Linear Interpolation and Lookup Table with Spline or Cubic Interpolation.

References

- [1] Tom M Apostol. Mathematical analysis. Narosa Publishing House Pvt. Ltd., 1985.