← Back **Week Four Quiz**
Graded Assignment

Due Mar 20, 11:59 PM CET

1. Quota sampling requires quota controls. 1 point

- (●) True
- ( ) False

2. Response bias is an example of: 1 point

- ( ) sampling error
- (●) non-sampling error

3. In probability sampling: 1 point

- (●) there is a known, non-zero probability of a population element being selected
- ( ) some population elements have a zero probability of being selected, while others have an unknown probability of being selected

4. In systematic sampling, N/n, is referred to as: 1 point

- (●) the sampling interval
- ( ) the sampling fraction

5. In stratified random sampling, elements between strata should be: 1 point

- (●) as heterogeneous as possible
- ( ) as homogeneous as possible

6. A sampling distribution is the: 1 point

- ( ) probability (frequency) distribution of a parameter
- (●) probability (frequency) distribution of a statistic

7. An unbiased estimator has its expected value equal to the: 1 point

- (●) parameter being estimated
- ( ) statistic used as the estimator

8. An x% confidence interval covers the unknown parameter 1 point

- (●) with x% probability over repeated samples
- ( ) with a probability of x%

9. Which of the following does **not** influence the size of the margin of error when considering the confidence interval for a population mean? 1 point

- ( ) the sample size, n
- (●) the sample mean, $\bar{x}$
- ( ) the standard deviation, σ or s

10. Other things equal, as the sample size n increases the margin of error: 1 point

- (●) decreases
- ( ) stays the same
- ( ) increases