

Biggest Maths-3 Buzz Now or Never Offer !

**Inspiring
the Future**

Fulfilling aspirations,
raising barriers, surpassing
Expectations

ધુવીલ બોરડ : ૯૬૮૭૦ ૦૫૫૭૫
શ્રીનાથજી એન્જીનીયરીંગ ઝોન

All Sem All Branch Admission Open

For All Engineering Mathematics

Diploma & Degree Engineering



Maths-1 (CALCULUS)

Maths-2 Maths -4

Maths-3

**Understand
Concepts**

Enjoy Learning

Enjoy learning Of Easy Mathematics
By Highly Quality Faculties

CRASH COURSE 100% RESULT

Maths-3 ના આગામી Crash Course માં ખુબજ સારો Response બદલ

આપ સૌનો ખુબ આભાર. તમામ સ્ટુડન્ટસ ને આપેલ **Question Bank** માંથી

45-50 Marks નુ GTU માં Questions બદલ્યા વગર પુછતા

શ્રીનાથજી એન્જીનીયરીંગ ઝોન પ્રત્યે સ્ટુડન્ટસ ઓ ઓ ખુબ સારો વિશ્વાસ વ્યક્ત કર્યો છે.

2 Hours NONSTOP MATHS-3 FOR Perfect Results

Resgistration Contact : 96870 05575

Crash Course Available At New Ranip

Branch-1 F.F-30 Aryavilla Residency, Anand Partyplot, G.S.T Crossing Road, New Ranip, Ahmedabad

Branch-2 A/21 Krushnanagar Soc. Indiacolony Road, Bapunagar, Ahmedabad

100 % Result !! Attention !! Grab Your Good Result Now

Highest Results in GTU 2017-18 & Exclusive IMP By Proff, D.G BORAD
Last Two Years 1000 % Results Proven With Over 300+ Students Clear Maths
After Join SHREENATHJI ENGINEERING ZONE & Witness Most Successful Engineering Class

Crash Course Special 100% Results

Once again See the Power of **Dhruvil Borad**

40-45 marks example asked in **Maths - 3**

IMP Given by Dhruvil Borad..

Imp posted in @Ametogtuvada

@Gujju_comedy

@anokhogujju

More then 1500 students save the post

Wich is Given By Dhruvil Borad..



For All Engineering Tutition

Shreenathji Education

DhruvilBorad(9687005576)

BRANCH 1

NEW RANIPET

BRANCH 2

INDIA COLONY

BRANCH 2

AANAND

Thanks all the students Trust on Shreenathji Education.

Wish You all Bright future Ahead...

More then 500 students send DM Dhruvil Borad on Instagram.

Wish you all Best results in your exam..

Thanks for Trust on us...

Thanks for being a part of

Shreenathji Education

Feel proud to help future Engineer...

Thank You

15/9 (1)

* Solve, $y' = 2xy$ by power series method.

Sol. Given eq: $y' - 2xy = 0$

$$y' = \sum_{k=0}^{\infty} a_k \cdot k \cdot x^{k-1}$$

$$y'' + P(x)y' + Q(x)y = 0$$

which is finite at $x=0$.

$\therefore x=0$ is a ordinary point.

$$\text{Let } y = \sum_{k=0}^{\infty} a_k x^k$$

$$y' = \sum_{k=0}^{\infty} a_k \cdot k \cdot x^{k-1}$$

Substitute in given equation we get,

$$\sum_{k=0}^{\infty} a_k \cdot k \cdot x^{k-1} - 2x \sum_{k=0}^{\infty} a_k x^k = 0$$

Power-Series

* Marks.

Rule-1. Limit series method.

$$\sum_{k=0}^{\infty} a_k \cdot k \cdot x^{k-1} - \sum_{k=2}^{\infty} 2 a_{k-2} x^{k-1} = 0$$

Rule-2

(Limit series method)

(Limit series method)

$$a_1 + \sum_{k=2}^{\infty} a_k \cdot k \cdot x^{k-1} - \sum_{k=2}^{\infty} 2 a_{k-2} x^{k-1} = 0$$

(Common term)

(Common term)

Homework: $y'' = y' \therefore y'' - y' = 0$

$$\text{Imp. } (1+x^2)y'' + xy' - 9y = 0$$

* most Imp (170) 110%

$$a_1 = 0 \text{ \& } \frac{a_k \cdot k - 2a_{k-2}}{k} = 0 \therefore a_k \cdot k = 2a_{k-2}$$

$$\Rightarrow \text{for } k=2; a_2 = \frac{2a_0}{2} = a_0$$

$$k=3; a_3 = \frac{2a_1}{3} = 0$$

$$k=4; a_4 = \frac{2a_2}{4} = \frac{2 \times a_0}{4} = \frac{a_0}{2}$$

$$k=5; a_5 = \frac{2a_3}{5} = \frac{2 \times 0}{5} = 0$$

$$k=6; a_6 = \frac{2a_4}{6} = \frac{2 \times \frac{a_0}{2}}{6} = \frac{a_0}{6}$$

Sol.:-

$$y = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + a_4 x^4 + a_5 x^5 + a_6 x^6 + \dots$$

$$y = a_0 + a_0 x^2 + \frac{a_0}{2} x^4 + \frac{a_0}{6} x^6 + \dots$$

Rule 1 Power series.

Rule 2 Limit series.

Rule 3. Power Series.

|| 9th ||

$$a_2 = \frac{9}{2} a_0, \quad a_3 = \frac{4}{3} a_1, \quad a_4 = \frac{5 a_2}{12} = \frac{5}{12} \cdot \frac{9}{2} a_0 = \frac{15}{24} a_0$$

$$y = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + a_4 x^4 + a_5 x^5 + \dots$$

$$y = a_0 + a_1 x + \frac{9 a_0}{2} x^2 + \frac{4}{3} a_1 x^3 + \frac{15 a_0}{24} x^4 + \dots$$

* find the series solution of

$$(1+x^2)y'' + xy' - 9y = 0 \quad (\text{GTO, 2013, 2014, 2015, 2016})$$

$$\therefore y'' + \frac{x}{1+x^2} y' - \frac{9}{1+x^2} y = 0$$

$$(1+x^2) \sum_{k=0}^{\infty} a_k k(k-1) x^{k-2} + x \sum_{k=0}^{\infty} a_k k x^{k-1} - 9 \sum_{k=0}^{\infty} a_k x^k = 0$$

20183
0123
0126

$$P(x) = \frac{x}{1+x^2} \text{ \& \; } Q(x) = \frac{-9}{1+x^2}$$

$$y'' + P(x)y' + Q(x)y = 0$$

$\Rightarrow x=0$ is an ordinary point.

$$\text{Let } y = \sum_{k=0}^{\infty} a_k x^k$$

$$y' = \sum_{k=0}^{\infty} a_k k x^{k-1}$$

$$y'' = \sum_{k=0}^{\infty} a_k k(k-1) x^{k-2}$$

$$= \sum_{k=0}^{\infty} a_k k(k-1) x^{k-2} + \sum_{k=0}^{\infty} a_k k(k-1) x^{k-1} + \sum_{k=0}^{\infty} a_k k x^k - \sum_{k=0}^{\infty} 9 a_k x^k = 0$$

$$= \sum_{k=0}^{\infty} a_k k(k-1) x^{k-2} + \sum_{k=2}^{\infty} a_{k-2} (k-2)(k-3) x^{k-2} + \sum_{k=2}^{\infty} a_{k-2} (k-2) x^{k-2} - \sum_{k=2}^{\infty} 9 a_{k-2} x^{k-2} = 0$$

$$a_k k(k-1) + a_{k-2} [(k-2)(k-3) + (k-2) - 9] = 0$$

$$a_k = - \frac{a_{k-2} [(k-2)(k-3) + (k-2) - 9]}{k(k-1)}$$