

Graduate Seminar 2

26121301

Ongonwou Fred

**Phd Candidate
2343003**

June 24, 2026

Graduate Seminar 2

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Section 1 - Solving for $\psi(x, t)$

- We solve $\psi(x, t)$ by a numerical resolution of the Time-Dependent Schrödinger Equation (TDSE), i.e

$$i \frac{\partial}{\partial t} \psi(x, t) = \left(\hat{H}_a + \hat{V}_{\text{ext}}(t) \right) \psi(x, t), \quad \psi(x, 0) = \psi_0(x), \quad (1)$$

$$\hat{H}_a = -\frac{1}{2} \frac{d^2}{dx^2} - \kappa \delta(x), \quad \hat{V}_{\text{ext}}(t) = F(t)x. \quad (2)$$

$$E_0 = -\kappa^2/2, \quad \psi_0(x) = \kappa^{1/2} e^{-\kappa|x|}, \quad \kappa = 1, \quad (3)$$

$$E_k = k^2/2, \quad \psi_k^{(\pm)}(x) = e^{ikx} + R(\pm|k|)e^{\pm i|kx|}, \quad R(k) = \frac{i\kappa}{k - i\kappa}. \quad (4)$$

- The parameters of the laser we will be using are given by the relations

$$F(t) = F_0 e^{-(2t'/\tau)^2} \cos(\omega_0 t'), \quad 0 \leq t \leq T, \quad t' = t - T/2, \quad (5)$$

$$\tau = 2\pi n_{oc}/\omega_0, \quad n_{oc} = 5. \quad (6)$$

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Density, survival probability and PEMD

- The density probability of the wavefunction, at t is given by

$$\mathcal{D}(x, t) = |\psi(x, t)|^2 = \int \psi^*(x, t) \psi(x, t) dx. \quad (7)$$

- The coefficients of the projection of ψ are then, at t , and by definition, expressed as

$$a_0 = e^{iE_0 t} \int \psi_0(x) \psi(x, t) dx, \quad a_k = e^{iE_k t} \int \psi_k^{(-)*}(x) \psi(x, t) dx. \quad (8)$$

- Then,

$$P_0 = |a_0|^2, \quad \text{and} \quad P(k) = |a_k|^2. \quad (9)$$

- The ionization probability can then be found many different ways that should be consistent with each others

$$P_{\text{ion}} = \int P(k) \frac{dk}{2\pi} = 1 - P_0. \quad (10)$$

- $P(k)$ is the probability for the electron being ionized with the energy E_k .

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Observables

- For an arbitrary operator $\mathcal{O}$, we have

$$\mathcal{O}(t) = \int \psi^*(x, t) \mathcal{O} \psi(x, t) dx \quad (11)$$

- The the Fourier transform is obtained using

$$\mathcal{O}(\omega) = \int \mathcal{O}(t) e^{i\omega t} dt. \quad (12)$$

- Following Bransden and Joachain, "Quantum Mechanics (2000)" (see 11.145), we define the transition amplitude $\mathcal{O}_{ba}$ and probability P_{ba} between two states a and b such as

$$\mathcal{O}_{ba} = \int dx \psi_b^*(x) \mathcal{O} \psi_a(x), \quad P_{ba} = |\mathcal{O}_{ba}|^2. \quad (13)$$

When continuum states between q and $q + dq$ are involved, the quantity of interest is

$$P_{ba} dq = \left| \int dx \psi_b^*(x) \mathcal{O} \psi_a(x) \right|^2 dq. \quad (14)$$

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Length, velocity and acceleration

- We want to evaluate the quantities

$$d(t) = -\langle \hat{x} \rangle = -\int \psi^*(x, t) \hat{x} \psi(x, t) dx, \quad \text{and} \quad \dot{x}(t) = \frac{d}{dt} x(t), \quad (15)$$

$$p(t) = \langle \hat{p} \rangle = -i \int \psi^*(x, t) \frac{\partial}{\partial x} \psi(x, t) dx. \quad (16)$$

- Direct numerical differentiation can sometimes be avoided with the help of the Ehrenfest theorem and the commutator algebra

$$[\hat{p}^2, \hat{x}] = -2i\hat{p} \implies \hat{p} = i[\hat{H}, \hat{x}], \quad (17)$$

$$\frac{d}{dt} \langle \hat{x} \rangle = \langle \hat{p} \rangle, \quad \frac{d}{dt} \langle \hat{p} \rangle = \left\langle -\frac{\partial V_{\text{ext}}}{\partial x} \right\rangle. \quad (18)$$

and some others (see Bransden and Joachain, Quantum Mechanics (2000), p.531-532)

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

High Harmonic Generation (HHG) - (semi-classical approx.)

- In the classical approx., the spectrum is obtained from the following (Eq.(80))

$$Q_{\nu}^{(c)}(\omega) = \alpha^2 \omega^2 |\mathbf{A}_{\nu} \mathbf{d}(\omega)|^2, \quad (19)$$

Let us apply the following simplifications

$$\nu \longrightarrow \omega, \quad \mathbf{A}_{\nu} = \sqrt{\frac{2\pi c^2}{\omega L^3}} \mathbf{e}_{\nu} \longrightarrow A_{\omega}, \quad \text{and} \quad \alpha A_{\omega} = 1. \quad (20)$$

- So the observable we are considering here is

$$Q^{(c)}(\omega) = \omega^2 |d(\omega)|^2. \quad (21)$$

Additionally let us consider Eq. (104)

$$Q^{(c)}(\omega) = |p(\omega)|^2. \quad (22)$$

and, by virtue of $\mathbf{p}(t) = -\dot{\mathbf{d}}(t)$, which reads

$$Q^{(c)}(\omega) = \left| -\dot{\mathbf{d}}(\omega) \right|^2. \quad (23)$$

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Additional note from Ehrenfest theorem

- Using the integration by parts formula, we get

$$p(\omega) = \int_{-T/2}^{T/2} e^{i\omega t} \frac{dx(t)}{dt} dt, \quad u(t) = x(t), \quad v(t) = e^{i\omega t}, \quad (24)$$

thus

$$p(\omega) = \int_{-T/2}^{T/2} \frac{d}{dt} \left(e^{i\omega t} x(t) \right) dt - i\omega \int_{-T/2}^{T/2} e^{i\omega t} x(t) dt = \left[e^{i\omega t} x(t) \right]_{-T/2}^{T/2} - i\omega \int_{-T/2}^{T/2} e^{i\omega t} x(t) dt, \quad (25)$$

which reduces to (21) up to a constant.

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Transition amplitude

- From Yangaliev *et al.*, we have

$$p_{k0} = \int dx \psi_k^{(-)*}(x) \hat{p} \psi_0(x) = \frac{2k\mathcal{K}^{3/2}}{\mathcal{K}^2 + k^2}, \quad (26)$$

$$p_{kk'} = \int dx \psi_k^{(-)*}(x) \hat{p} \psi_{k'}^{(-)}(x) = 2\pi k \delta(k - k') - \frac{2\mathcal{K}}{k^2 - k'^2 + i\epsilon} \left(\frac{k'|k|}{|k| - i\mathcal{K}} - \frac{k|k'|}{|k'| + i\mathcal{K}} \right). \quad (27)$$

- Using (17), we can rewrite those expressions as

$$p_{k0} = -i(E_k - E_0) d_{k0}, \quad (28)$$

$$p_{kk'} = -i(E_k - E_{k'}) d_{kk'}. \quad (29)$$

- Those quantities are another way to study convergence, at the structural level

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Part 2 - Solving $\phi_\omega(x, t)$

- We are now interested in solving the following 1D problem with a ZRP

$$\begin{cases} i \frac{\partial}{\partial t} \psi(x, t) &= \left(\hat{H}_a + \hat{V}_{\text{ext}}(t) \right) \psi(x, t), \\ i \frac{\partial}{\partial t} \phi_\omega(x, t) &= \left(\hat{H}_a + \hat{V}_{\text{ext}}(t) \right) \phi_\omega(x, t) + e^{i\omega t} S(x, t) \end{cases} \quad (30)$$

where

$$\hat{H}_a = -\frac{1}{2} \frac{d^2}{dx^2} - \kappa \delta(x), \quad \hat{V}_{\text{ext}}(t) = F(t)x, \quad S(x, t) = \hat{p} \psi(x, t) \quad (31)$$

$$\phi_\omega(x, 0) = \frac{-i\kappa^{3/2}}{\omega} \text{sgn}(x) \left(e^{-\kappa|x|} - e^{-\kappa(\omega)|x|} \right), \quad \psi(x, 0) = \kappa^{1/2} e^{-\kappa|x|}, \quad (32)$$

$$\hat{p} = -id/dx, \quad \text{and} \quad \kappa(\omega) = \sqrt{\kappa^2 + 2\omega}. \quad (33)$$

$$\phi_\omega(x, t) = U(t, t_0) \phi_\omega(x, t_0) - i \int_{t_0}^t U(t - \tau) e^{i\omega \tau} S(x, \tau) d\tau. \quad (34)$$

High-order exponential splitting (composition methods) (Phys. Rev. A 101, 013410 (2020))

Yoshida (1990) - 4th-order symmetric decomposition

$$\phi_\omega(x, t + \Delta t) = U_4(t + \Delta t, t)\phi_\omega(x, t) - i \int_t^{t+\Delta t} U_4(t + \Delta t - \tau) e^{i\omega\tau} S(x, \tau) d\tau. \quad (34)$$

- From the following Strang operator $U(h) = e^{-iE_n h/2} e^{-iV_{\text{ext}}(t+h/2)h} e^{-iE_n h/2}$, we compose

$$U_4(h) = U(c_1 h) U(c_2 h) U(c_1 h), \quad \text{with} \quad c_1 = \frac{1}{2 - 2^{1/3}}, \quad c_2 = -\frac{2^{1/3}}{2 - 2^{1/3}}. \quad (35)$$

- More precisely, we can observe the following

Theorem 8.6 (Hairer, Solving ODE II p.554)

If Ψ_h represent a one-step method that is symmetric, of order $p = 2k$ and defined in the domain, then the relations

$$2c_1 + c_2 = 1, \quad 2c_1^{p+1} + c_2^{p+1} = 0, \quad (36)$$

imply that the composition (35) is of order $p + 2$.

If you start from the Strang method, then in the above $k = 1$.

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Midpoint Duhamel - 2nd order approximation

$$\phi_{\omega}(x, t + \Delta t) = U(t + \Delta t, t)\phi_{\omega}(x, t) - i \int_t^{t+\Delta t} U(t + \Delta t - \tau) e^{i\omega\tau} S(x, \tau) d\tau. \quad (34)$$

- We can show that the above can be expressed as

$$\phi_{n+1}^{\omega} = U(\Delta t)\phi_n^{\omega} - i\Delta t e^{i\omega t_{n+1/2}} S_{n+1/2}, \quad (37)$$

where the source is evaluated at $t_{n+1/2} = t + \Delta t/2$, i.e and $S_{n+1/2} = U(\Delta t/2)S_n$, with $S_n = S(x, t_n)$

- This is the midpoint Duhamel formula, of order 2.
- It's what will allow us to get familiar with the numerical intricacies of our problem
- The driver shall be regarded as decent if the expected order is met

Geometric propagators

Simpson Duhamel - 4th-order quadrature

$$\phi_\omega(x, t + \Delta t) = U_4(t + \Delta t, t)\phi_\omega(x, t) - i \int_t^{t+\Delta t} U_4(t + \Delta t - \tau) e^{i\omega\tau} S(x, \tau) d\tau. \quad (34)$$

- The Simpson quadrature consists in writing

$$\phi_{n+1}^\omega = U_4(\Delta t)\phi_n^\omega - \frac{i\Delta t}{6} \left(U_4(\Delta t)e^{i\omega t_n} S_n + 4U_4(\Delta t/2)e^{i\omega t_{n+\frac{1}{2}}} S_{n+\frac{1}{2}} + U_4(0)e^{i\omega t_{n+1}} S_{n+1} \right) \quad (38)$$

where the source is evaluated at t_n , $t_{n+\frac{1}{2}}$ and t_{n+1}

- This is the Simpson Duhamel formula, of order 4.
- It requires a 4th order evolution operator to work
- Simpson-Duhamel multiplies the number of operations required by Midpoint-Duhamel by 3.

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Observables

$$b_{0\omega}(T) = e^{iE_0 T} \int \psi_0(x) \phi_\omega(x, T) dx, \quad (39)$$

$$b_{k\omega}(T) = e^{iE_k T} \int \psi_k^{(-)*}(x) \phi_\omega(x, T) dx, \quad (40)$$

$$b_{0\omega} = b_{0\omega}(T) + \int \frac{e^{i(E_0 + \omega - E_{k'})T} p_{0k'} a_{k'}}{E_0 + \omega - E_{k'} + i0} \frac{dk'}{2\pi} \quad (41)$$

$$b_{k\omega} = b_{k\omega}(T) + \frac{e^{i(E_k + \omega - E_0)T} p_{k0} a_0}{E_k + \omega - E_0} + \int \frac{e^{i(E_k + \omega - E_{k'})T} p_{kk'} a_{k'}}{E_k + \omega - E_{k'} + i0} \frac{dk'}{2\pi} \quad (42)$$

$$Q(\omega) = |b_{0\omega}|^2 + \int |b_{k\omega}|^2 \frac{dk}{2\pi} \quad (43)$$

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Midpoint Duhamel - Self conv. test results for $S(x, t) = \hat{p}\psi(x, t)$ - ZRP

$$\phi_{n+1}^\omega = U(\Delta t)\phi_n^\omega - i\Delta t e^{i\omega t_{n+\frac{1}{2}}} S_{n+\frac{1}{2}} \tag{38}$$

t	$\Re \psi(1)$	$\Im \psi(1)$	$\ \psi\ $	$\Re \phi(2)$	$\Im \phi(2)$	$\ \phi\ $
-1102.3132	1.0000000	0.0000000	1.0000000	0.0000000	1.4733146(-4)	0.6074294
-1080.2669	1.0000000	-1.5299150(-5)	1.0000000	9.0161878(-8)	1.4733146(-4)	2.9977570
-1058.2207	1.0000000	-3.0598299(-5)	1.0000000	1.8032375(-7)	1.4733145(-4)	4.2519438
			...			
-22.0463	-4.6878735(-7)	1.1266426(-6)	1.0000000	-3.4864304(-3)	-2.0577179(-3)	0.7712524
-0.0000	4.8643339(-1)	-9.9881227(-1)	1.0000000	-1.3514523	-8.0617898(-4)	3.3039060
22.0463	2.8495297(-7)	1.2366110(-6)	1.0000000	-3.7992762(-3)	1.3930728(-3)	4.2371529
			...			
1058.2207	-9.9526450(-1)	-9.7202282(-2)	1.0000000	1.5396368(-5)	-1.4659389(-4)	1.1029687
1080.2669	-9.9526599(-1)	-9.7187055(-2)	1.0000000	1.5389349(-5)	-1.4659255(-4)	3.5763755
1102.3132	-9.9526748(-1)	-9.7171829(-2)	1.0000000	1.5382713(-5)	-1.4659538(-4)	4.1762517

Level	Δt (relative)	E_ψ	E_ϕ	P_ψ/P_ϕ
1	Δt_0	0.1185025(-1)	0.1632911(+0)	—
2	$\Delta t_0/2$	0.2953778(-2)	0.3951342(-1)	2.004 / 2.048
3	$\Delta t_0/4$	0.7378959(-3)	0.9798974(-2)	2.001 / 2.011
Observed order				2.0011/2.0116

Ne = 100, Np = 10
-xmin = xrange/2 = Np Ns/2
 $\tau = 2\pi n_{oc}/\omega_0$, $T = 8\tau$

Initial conditions and parameters

$$S(x, t) = -i\partial_x \psi(x, t)$$
$$F(t) = F_0 e^{-(2t'/\tau)^2} \cos(\omega_0 t), \quad t' = t - T/2$$
$$F_0 = 0.03, \quad \omega_0 = 0.114, \quad \omega = 0.5$$
$$\Delta t_0 = 2.204, \quad n_{oc} = 5$$

■ Results of the time propagation for the zero range potential with and without source

$$p_k^{\text{self}} = (\text{err}_{k-1}/\text{err}_k) / \log(2)$$

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Simpson Duhamel - Self conv. test results for $S(x, t) = \hat{p}\psi(x, t)$ - ZRP

$$\phi_{n+1}^\omega = U_4(\Delta t)\phi_n^\omega - \frac{i\Delta t}{6} \left(U_4(\Delta t)e^{i\omega t_n} S_n + 4U_4(\Delta t/2)e^{i\omega t_{n+\frac{1}{2}}} S_{n+\frac{1}{2}} + U_4(0)e^{i\omega t_{n+1}} S_{n+1} \right) \quad (38)$$

$$U_4(t) = U(c_1 h)U(c_2 h)U(c_1 h), \quad (35)$$

t	$\Re \psi(1)$	$\Im \psi(1)$	$\ \psi\ $	$\Re \phi(2)$	$\Im \phi(2)$	$\ \phi\ $
-1102.3132	1.0000000	0.0000000	1.0000000	0.0000000	1.4733146(-4)	0.6074294
-1080.2669	1.0000000	-1.5299150(-5)	1.0000000	9.0161878(-8)	1.4733146(-4)	2.9977570
-1058.2207	1.0000000	-3.0598299(-5)	1.0000000	1.8032375(-7)	1.4733145(-4)	4.2519438
			...			
-22.0463	-4.6878735(-7)	1.1266426(-6)	1.0000000	-3.4864304(-3)	-2.0577179(-3)	0.7712524
-0.0000	4.8643339(-1)	-9.9881227(-1)	1.0000000	-1.3514523	-8.0617898(-4)	3.3039060
22.0463	2.8495297(-7)	1.2366110(-6)	1.0000000	-3.7992762(-3)	1.3930728(-3)	4.2371529
			...			
1058.2207	-9.9526450(-1)	-9.7202282(-2)	1.0000000	1.5396368(-5)	-1.4659389(-4)	1.1029687
1080.2669	-9.9526599(-1)	-9.7187055(-2)	1.0000000	1.5389349(-5)	-1.4659255(-4)	3.5763755
1102.3132	-9.9526748(-1)	-9.7171829(-2)	1.0000000	1.5382713(-5)	-1.4659538(-4)	4.1762517

Level	Δt (relative)	E_ψ	E_ϕ	p_ψ/p_ϕ
1	Δt_0	0.5564105(-3)	0.1529880(-1)	—
2	$\Delta t_0/2$	0.3480484(-4)	0.1093299(-2)	3.999 / 3.807
3	$\Delta t_0/4$	0.2175758(-5)	0.7047939(-4)	4.000 / 3.955
Observed order		3.9997/3.9553		

Ne = 100, Np = 10

-xmin = xrange/2 = Np Ns/2

$\tau = 2\pi n_{oc}/\omega_0$, $T = 8\tau$

Initial conditions and parameters

$S(x, t) = -i\partial_x \psi(x, t)$

$F(t) = F_0 e^{-(2t'/\tau)^2} \cos(\omega_0 t)$, $t' = t - T/2$

$F_0 = 0.03$, $\omega_0 = 0.114$, $\omega = 0.5$


$\Delta t_0 = 2.204$, $n_{oc} = 5$

■ Results of the time propagation for the zero range potential with and without source

$$p_k^{\text{self}} = (\text{err}_{k-1}/\text{err}_k) / \log(2)$$

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Transition amplitudes - gauge-related convergence



$N_p = 20, \quad N_s = 100$
 $x_{\max} \sim N_p N_s / 2$
 $\omega_0 = 0.114 \text{ a.u.}$
 $F_0 = 0.03 \text{ a.u.}, \quad T = 8\tau$



$$p_{k0} = \int dx \psi_k^{(-)*}(x) \hat{p} \psi_0(x) \quad (44)$$

$$\sum_{k'} |p_{kk'} a_{k'}| dk = \sum_{k'} \left| \int dx \psi_k^{(-)*}(x) \hat{p} \psi_k^{(-)}(x) a_{k'} \right| dk \quad (45)$$

U: $N_p = 20, N_s = 100, x_{\max} = 1000$
 D: $N_p = 40, N_s = 200, x_{\max} = 4000$

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Convergence region for ϕ_ω in x , $T = 5\tau$, with $F_0 = 0.03$ a.u. and $\omega_0 = .114$ a.u.



- The convergence region for ψ is $x_{\text{range}} \sim 2000$
- The convergence region for ϕ is $x_{\text{range}} \sim 4500$
- The plateau for ϕ seems to be $1e - 8$, above the double precision limit
- The result is confirmed by 2 different methods and has been tested against an analytical result

$$N_s = 200$$

$$x_{\text{max}} \sim (N_p N_s + q) / 2$$

$$\omega_0 = 0.114 \text{ a.u.}$$

$$q = 1/4$$


$$F_0 = 0.03 \text{ a.u.}, \quad T = 5\tau$$

$$n_{\text{steps}} = 20000, \quad \Delta t = 0.06889$$



Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Study of the behavior of ϕ_ω against the equation (homogeneous vs inhomogeneous)



$$b_{0\omega}(T) = e^{iE_0 T} \int \psi_0(x) \phi_\omega(x, T) dx,$$

$$b_{k\omega}(T) = e^{iE_k T} \int \psi_k^{(-)*}(x) \phi_\omega(x, T) dx,$$

$$Q(\omega) = |b_{0\omega}|^2 + \int |b_{k\omega}|^2 \frac{dk}{2\pi}$$

$$b_{0\omega} = b_{0\omega}(T) + \int \frac{e^{i(E_0 + \omega - E_{k'})T} p_{0k'} a_{k'}}{E_0 + \omega - E_{k'} + i0} \frac{dk'}{2\pi}$$

$$b_{k\omega} = b_{k\omega}(T) + \frac{e^{i(E_k + \omega - E_0)T} p_{k0} a_0}{E_k + \omega - E_0} + \int \frac{e^{i(E_k + \omega - E_{k'})T} p_{kk'} a_{k'}}{E_k + \omega - E_{k'} + i0} \frac{dk'}{2\pi}$$

$$i\partial_t \phi_\omega(t) = (\hat{H}_a + \hat{V}_{\text{ext}}(t)) \phi_\omega(t)$$

$$i\partial_t \phi_\omega(t) = (\hat{H}_a + \hat{V}_{\text{ext}}(t)) \phi_\omega(t) - ie^{i\omega t} \partial_x \psi(x, t)$$

$$N_p = 20, \quad N_s = 100$$

$$x_{\text{max}} \sim (N_p N_s + q) / 2$$

$$\omega_0 = 0.114 \text{ a.u.}$$


$$q = 1/4$$

$$F_0 = 0.03 \text{ a.u.}, \quad T = 5\tau$$

$$n_{\text{steps}} = 20000, \quad \Delta t = 0.06889$$

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Convergence of the vector, non integral transition terms $p_{k0}a_0$



$$C_{k0}^\omega p_{k0}a_0 = \frac{e^{i(E_k + \omega - E_0)T}}{E_k + \omega - E_0} p_{k0}a_0$$

$$Q(\omega) = |b_{0\omega}|^2 + \int |b_{k\omega}|^2 \frac{dk}{2\pi}$$

$$N_p = 20(\text{up}), 30(\text{down}), \quad N_s = 200$$

$$x_{\text{max}} \sim (N_p N_s + q) / 2$$

$$\omega_0 = 0.114 \text{ a.u.}$$

$$q = 1/10$$

$$F_0 = 0.03 \text{ a.u.}, \quad T = 5\tau$$


$$n_{\text{steps}} = 20000, \quad \Delta t = 0.06889$$

$$b_{0\omega} = b_{0\omega}(T) + \int \frac{e^{i(E_0 + \omega - E_{k'})T} p_{0k'} a_{k'}}{E_0 + \omega - E_{k'} + i0} \frac{dk'}{2\pi}$$

$$b_{k\omega} = b_{k\omega}(T) + \frac{e^{i(E_k + \omega - E_0)T} p_{k0} a_0}{E_k + \omega - E_0} + \int \frac{e^{i(E_k + \omega - E_{k'})T} p_{kk'} a_{k'}}{E_k + \omega - E_{k'} + i0} \frac{dk'}{2\pi}$$

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Convergence of the integral transition terms $p_{0k'} a_{k'}$, and $p_{kk'} a_{k'}$



$$\int C_{0k}^{\omega} p_{0k} a_k dk = \int dk \frac{e^{i(E_0 + \omega - E_k)T}}{E_0 + \omega - E_k} p_{0k} a_k \frac{dk}{2\pi}$$

$$\int C_{kk'}^{\omega} p_{kk'} a_{k'} dk = \int dk' \frac{e^{i(E_k + \omega - E_{k'})T}}{E_k + \omega - E_{k'}} p_{kk'} a_{k'} \frac{dk'}{2\pi}$$

$$Q(\omega) = |b_{0\omega}|^2 + \int |b_{k\omega}|^2 \frac{dk}{2\pi}$$

$$N_p = 20, \quad N_s = 200$$

$$x_{\max} \sim (N_p N_s + q) / 2$$

$$\omega_0 = 0.114 \text{ a.u.}$$

$$q = 1/10$$

$$F_0 = 0.03 \text{ a.u.}, \quad T = 5\tau$$


$$n_{\text{steps}} = 20000, \quad \Delta t = 0.06889$$

$$b_{0\omega} = b_{0\omega}(T) + \int \frac{e^{i(E_0 + \omega - E_{k'})T}}{E_0 + \omega - E_{k'} + i0} p_{0k'} a_{k'} \frac{dk'}{2\pi}$$

$$b_{k\omega} = b_{k\omega}(T) + \frac{e^{i(E_k + \omega - E_0)T} p_{k0} a_0}{E_k + \omega - E_0} + \int \frac{e^{i(E_k + \omega - E_{k'})T} p_{kk'} a_{k'}}{E_k + \omega - E_{k'} + i0} \frac{dk'}{2\pi}$$

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Convergence of the integral transition terms $p_{0k'} a_{k'}$, and $p_{kk'} a_{k'}$



$$\int C_{0k}^{\omega} p_{0k} a_k dk = \int dk \frac{e^{i(E_0 + \omega - E_k)T}}{E_0 + \omega - E_k} p_{0k} a_k \frac{dk}{2\pi}$$

$$\int C_{kk'}^{\omega} p_{kk'} a_{k'} = \int dk' \frac{e^{i(E_k + \omega - E_{k'})T}}{E_k + \omega - E_{k'}} p_{kk'} a_{k'} \frac{dk'}{2\pi}$$

$$Q(\omega) = |b_{0\omega}|^2 + \int |b_{k\omega}|^2 \frac{dk}{2\pi}$$

$$N_p = 20, \quad N_s = 200$$

$$x_{\max} \sim (N_p N_s + q) / 2$$

$$\omega_0 = 0.114 \text{ a.u.}$$

$$q = 1/10$$

$$F_0 = 0.03 \text{ a.u.}, \quad T = 5\tau$$


$$n_{\text{steps}} = 20000, \quad \Delta t = 0.06889$$

$$b_{0\omega} = b_{0\omega}(T) + \int \frac{e^{i(E_0 + \omega - E_{k'})T}}{E_0 + \omega - E_{k'} + i0} p_{0k'} a_{k'} \frac{dk'}{2\pi}$$

$$b_{k\omega} = b_{k\omega}(T) + \frac{e^{i(E_k + \omega - E_0)T} p_{k0} a_0}{E_k + \omega - E_0} + \int \frac{e^{i(E_k + \omega - E_{k'})T} p_{kk'} a_{k'}}{E_k + \omega - E_{k'} + i0} \frac{dk'}{2\pi}$$

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Study of the the coefficients $b_{k\omega}(T)$, norm and PEMD



$$b_{0\omega}(T) = e^{iE_0 T} \int \psi_0(x) \phi_{\omega}(x, T) dx,$$

$$b_{k\omega}(T) = e^{iE_k T} \int \psi_k^{(-)*}(x) \phi_{\omega}(x, T) dx,$$

$$Q(\omega) = |b_{0\omega}|^2 + \int |b_{k\omega}|^2 \frac{dk}{2\pi}$$

$$N_p = 20(\text{up}), 30(\text{down}), \quad N_s = 200$$

$$x_{\text{max}} \sim (N_p N_s + q) / 2$$

$$\omega_0 = 0.114 \text{ a.u.}$$

$$q = 1/10$$

$$F_0 = 0.03 \text{ a.u.}, \quad T = 5\tau$$

$$n_{\text{steps}} = 20000, \quad \Delta t = 0.06889$$

$$b_{0\omega} = b_{0\omega}(T) + \int \frac{e^{i(E_0 + \omega - E_{k'})T} p_{0k'} a_{k'}}{E_0 + \omega - E_{k'} + i0} \frac{dk'}{2\pi}$$

$$b_{k\omega} = b_{k\omega}(T) + \frac{e^{i(E_k + \omega - E_0)T} p_{k0} a_0}{E_k + \omega - E_0} + \int \frac{e^{i(E_k + \omega - E_{k'})T} p_{kk'} a_{k'}}{E_k + \omega - E_{k'} + i0} \frac{dk'}{2\pi}$$

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Transition amplitude convergence

- From the two precedent slides, we can infer for p_{k0} and $p_{kk'}$, that
 - their associated values only depends on the structure
 - their converged values has been evaluated (see table next slide)
- We can then reliably reduce the convergence of $Q(\omega)$ to $b_{0\omega}(T)$ and $b_{k\omega}(T)$
 - ~~Convergence of $b_{k\omega}(T)$ is strongly influenced by the norm of ϕ_ω at the end of the interaction. The norm of ϕ_ω strongly oscillates and its values dictate the amplitudes $b_{0\omega}$ and $b_{k\omega}$~~
 - ϕ_ω is not normed at t_0 and converges toward a value a priori different from $|\phi_\omega(x, 0)|$
 - We have been unsuccessful predicting self value to guide our debugging and the validity of our procedure

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Quantum High Harmonic Generation, for ϕ_ω with $\omega = 0.5$ a.u. (old file)

N_p	N_s	q	$x_{\max}$	$b_0(\omega)$	$Q(\omega)$
30	200	1/2	9000	12.7406754	6.3649490×10^9
40	200	0	4000	12.7400795	6.2368512×10^3
40	200	1/2	12000	12.7412440	2.6370941×10^7
50	200	1/2	15000	12.7976569	1.2622004×10^7

For this laser intensity and frequency, the reference result is

$$Q(\omega) \sim 1e - 5$$

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Quantum High Harmonic Generation, for ϕ_ω with $\omega = 0.5$ a.u. (old file)

Case	N_p	$b_0(\omega)$	$Q(\omega)$
Homogeneous	20	1.4297231×10^{-5}	1.0152861×10^2
Homogeneous	30	1.4296998×10^{-5}	9.3657583
Inhomogeneous	20	12.7625624	1.1429193×10^2
Inhomogeneous	30	12.7625623	2.2128765×10^1
Inhomogeneous	40	12.7625622	2.0121887×10^1

For this laser intensity and frequency, the reference result is

$$Q(\omega) \sim 1e-5$$

Quantum theory of radiation by nonstationary systems with application to high-order harmonic generation (Phys. Rev. A 101, 013410 (2020))

Quantum High Harmonic Generation, for ϕ_ω with $\omega = 0.5$ a.u.

Run	nsteps	Re $b_0(\omega)$	Im $b_0(\omega)$	$Q(\omega)$
Homogeneous	20000	$-0.6830301(-5)$	$0.1928069(-5)$	$3.2058559(-3)$
Driven	15000	$-0.1770619(-2)$	$-0.3836150(-3)$	$1.7834170(-3)$
Driven	20000	$-0.1770615(-2)$	$-0.3836362(-3)$	$1.7828446(-3)$

For this laser intensity and frequency, the reference result is $Q(\omega) \sim 1e-5$