



Expected Value & Standard Error

What does the sum look like?



Review



Review

Box Model

... is like drawing _____ times from the box _____ with replacement and summing the draws.



Review

Box Model

... *is like* drawing _____ times from the box _____ with replacement and summing the draws.

Example #1

Rolling a die 40 times and summing the numbers shown

Review

Box Model

... is like drawing _____ times from the box _____ with replacement and summing the draws.

Example #1

Rolling a die 40 times and summing the numbers shown
...is like...

Review

Box Model

... is like drawing _____ times from the box _____ with replacement and summing the draws.

Example #1

Rolling a die 40 times and summing the numbers shown
...is like...

drawing _____ times from the box _____ with replacement and summing the draws.

Review

Box Model

... *is like* drawing _____ times from the box _____ with replacement and summing the draws.

Example #1

Rolling a die 40 times and summing the numbers shown
...*is like*...

drawing 40 times from the box _____ with replacement and summing the draws.

Review

Box Model

... is like drawing _____ times from the box _____ with replacement and summing the draws.

Example #1

Rolling a die 40 times and summing the numbers shown
...is like...

drawing 40 times from the box 1, 2, 3, 4, 5, 6 with replacement and summing the draws.

Review

Box Model

... is like drawing _____ times from the box _____ with replacement and summing the draws.

Example #2

Rolling a die 40 times and counting the number of aces

...is like...

drawing _____ times from the box _____ with replacement and summing the draws.

Review

Box Model

... is like drawing _____ times from the box _____ with replacement and summing the draws.

Example #2

Rolling a die 40 times and counting the number of aces

...is like...

drawing 40 times from the box _____ with replacement and summing the draws.

Review

Box Model

... is like drawing _____ times from the box _____ with replacement and summing the draws.

Example #2

Rolling a die 40 times and counting the number of aces

...is like...

drawing 40 times from the box 0, 0, 0, 0, 0, 1 with replacement and summing the draws.

Fact

**If we execute the box model,
the result is a sum.**

Fact

**If we execute the box model,
the result is a sum.**

Box Model

***... is like* drawing _____ times from the
box _____ with replacement and
summing the draws.**

Fact

If we execute the box model,
the result is a sum.

Box Model

... *is like* drawing _____ times from the
box _____ with replacement and
summing the draws.

Fact

If we execute the box model,
the result is a sum.

Box Model

... *is like* drawing _____ times from the
box _____ with replacement and
summing the draws.

Question

Fact

If we execute the box model,
the result is a sum.

Box Model

... *is like* drawing _____ times from the
box _____ with replacement and
summing the draws.

Question

What can we say about this
(yet to be produced) sum?

Question

**What can we say about this
(yet to be produced) sum?**

Question

**What can we say about this
(yet to be produced) sum?**

The sum will be about _____ give or take _____ or so.

Question

What can we say about this
(yet to be produced) sum?

the expected value



The sum will be about _____ give or take _____ or so.

Question

What can we say about this
(yet to be produced) sum?

The sum will be about give or take or so.

the expected value ↓

the standard error ↓

Question

What can we say about this
(yet to be produced) sum?

the expected value

the standard error

The sum will be about _____ give or take _____ or so.







Suppose an actual list of numbers.



Suppose an actual list of numbers.

The entries in that list are about [the average] give or take [the SD] or so.



Suppose an actual list of numbers.

The entries in that list are about [the average] give or take [the SD] or so.

Suppose a hypothetical list of numbers that we generate by executing the box model an infinite number of times.



Suppose an actual list of numbers.

The entries in that list are about [the average] give or take [the SD] or so.

Suppose a hypothetical list of numbers that we generate by executing the box model an infinite number of times.

The (hypothetical) average is the expected value.



Suppose an actual list of numbers.

The entries in that list are about [the average] give or take [the SD] or so.

Suppose a hypothetical list of numbers that we generate by executing the box model an infinite number of times.

The (hypothetical) average is the expected value.

The (hypothetical) SD is the standard error.



Suppose an actual list of numbers.

The entries in that list are about [the average] give or take [the SD] or so.

Suppose a hypothetical list of numbers that we generate by executing the box model an infinite number of times.

The (hypothetical) average is the expected value.

The (hypothetical) SD is the standard error.

We can think of the expected value and standard error as a “long-run” average and SD of a chance process.



Suppose an actual list of numbers.

The entries in that list are about [the average] give or take [the SD] or so.

Suppose a hypothetical list of numbers that we generate by executing the box model an infinite number of times.

The (hypothetical) average is the expected value.

The (hypothetical) SD is the standard error.

We can think of the expected value and standard error as a “long-run” average and SD of a chance process.

	actual list of numbers	chance process
typical value	average	expected value
give or take	SD	standard error

Equations

Equations

expected value for sum = (number of draws) \times (average of box)

Equations

expected value for sum = (number of draws) \times (average of box)

SE for sum = $\sqrt{\text{number of draws}}$ \times (SD of box)

Equations

expected value for sum = (number of draws) \times (average of box)

SE for sum = $\sqrt{\text{number of draws}} \times (\text{SD of box})$

Helpful Hints

Equations

expected value for sum = (number of draws) \times (average of box)

SE for sum = $\sqrt{\text{number of draws}} \times (\text{SD of box})$

Helpful Hints

Suppose the box is a “big-small box” that only has big numbers B and small numbers S (e.g., the box 2, 2, 2, 2, 14, 14), then

Equations

expected value for sum = (number of draws) \times (average of box)

SE for sum = $\sqrt{\text{number of draws}} \times (\text{SD of box})$

Helpful Hints

Suppose the box is a “big-small box” that only has big numbers B and small numbers S (e.g., the box 2, 2, 2, 2, 14, 14), then

SD of big-small box = $(B - S) \times \sqrt{(\text{fraction that are B}) \times (\text{fraction that are S})}$

Equations

expected value for sum = (number of draws) \times (average of box)

SE for sum = $\sqrt{\text{number of draws}} \times (\text{SD of box})$

Helpful Hints

Suppose the box is a “big-small box” that only has big numbers B and small numbers S (e.g., the box 2, 2, 2, 2, 14, 14), then

SD of big-small box = $(B - S) \times \sqrt{(\text{fraction that are B}) \times (\text{fraction that are S})}$

Suppose the box is a “0-1 box” that only has 0s and 1s (e.g., the box 0, 0, 0, 1), then

Equations

expected value for sum = (number of draws) \times (average of box)

SE for sum = $\sqrt{\text{number of draws} \times (\text{SD of box})}$

Helpful Hints

Suppose the box is a “big-small box” that only has big numbers B and small numbers S (e.g., the box 2, 2, 2, 2, 14, 14), then

SD of big-small box = $(B - S) \times \sqrt{(\text{fraction that are B}) \times (\text{fraction that are S})}$

Suppose the box is a “0-1 box” that only has 0s and 1s (e.g., the box 0, 0, 0, 1), then

SD of 0-1 box = $\sqrt{(\text{fraction that are 0}) \times (\text{fraction that are 1})}$

Example #1

Rolling a die 40 times and summing the numbers shown
...is like...

drawing 40 times from the box 1, 2, 3, 4, 5, 6 with
replacement and summing the draws.

Example #1

Rolling a die 40 times and summing the numbers shown
...is like...

drawing 40 times from the box 1, 2, 3, 4, 5, 6 with
replacement and summing the draws.

The sum will be about _____ give or take _____ or so.

Example #1

Rolling a die 40 times and summing the numbers shown
...is like...

drawing 40 times from the box 1, 2, 3, 4, 5, 6 with
replacement and summing the draws.

the expected value



The sum will be about _____ give or take _____ or so.

Example #1

Rolling a die 40 times and summing the numbers shown
...is like...

drawing 40 times from the box 1, 2, 3, 4, 5, 6 with
replacement and summing the draws.

the expected value
the standard error

The sum will be about _____ give or take _____ or so.

Example #1

Rolling a die 40 times and summing the numbers shown
...is like...

drawing 40 times from the box 1, 2, 3, 4, 5, 6 with replacement and summing the draws.

the expected value
the standard error

The sum will be about _____ give or take _____ or so.

$$\begin{aligned}\text{expected value for sum} &= (\text{number of draws}) \times (\text{average of box}) \\ &= 40 \times 3.5 \\ &= 140\end{aligned}$$

Example #1

Rolling a die 40 times and summing the numbers shown
...is like...

drawing 40 times from the box 1, 2, 3, 4, 5, 6 with replacement and summing the draws.

the expected value ↓
the standard error ↓
The sum will be about 140 give or take _____ or so.

$$\begin{aligned}\text{expected value for sum} &= (\text{number of draws}) \times (\text{average of box}) \\ &= 40 \times 3.5 \\ &= 140\end{aligned}$$

Example #1

Rolling a die 40 times and summing the numbers shown
...is like...

drawing 40 times from the box 1, 2, 3, 4, 5, 6 with replacement and summing the draws.

the expected value
the standard error

The sum will be about 140 give or take _____ or so.

Example #1

Rolling a die 40 times and summing the numbers shown
...is like...

drawing 40 times from the box 1, 2, 3, 4, 5, 6 with replacement and summing the draws.

the expected value

the standard error

The sum will be about 140 give or take _____ or so.

SE for sum = $\sqrt{\text{number of draws}} \times (\text{SD of box})$

= $\sqrt{40} \times ???$

= **6.32** × **1.71**

= **10.81**

```
> x <- c(1, 2, 3, 4, 5, 6)
> sqrt(mean((x - mean(x))^2))
[1] 1.707825
```


Example #1

Rolling a die 40 times and summing the numbers shown
...is like...

drawing 40 times from the box 1, 2, 3, 4, 5, 6 with replacement and summing the draws.

the expected value
the standard error

The sum will be about 140 give or take 11 or so.

SE for sum = $\sqrt{\text{number of draws}} \times (\text{SD of box})$


= $\sqrt{40} \times ???$

= **6.32** × **1.71**

= **10.81**


```
> x <- c(1, 2, 3, 4, 5, 6)
> sqrt(mean((x - mean(x))^2))
[1] 1.707825
```

I rolled a die 40 times and got a sum of 141. That's right in line with our claim that the sum will be about 140 give or take 11 or so.



I rolled a die 40 times and got a sum of 141. That's right in line with our claim that the sum will be about 140 give or take 11 or so.

I did it nine more times and got 116, 137, 143, 146, 130, 126, 136, 128, and 138. Again, that's right in line with our claim that the sum will be about 140 give or take 11 or so.



Highlights

- If we have a chance process, we can sometimes describe it with a **box model**.
- If we have a box model, then we can say compute the **expected value** and the **standard error**.

expected value for sum = (number of draws) \times (average of box)

SE for sum = $\sqrt{\text{number of draws}}$ \times (SD of box)

- With the expected value and standard error, we can fill in the following: **The sum will be about ____ give or take ____ or so.**