1

Probability Assignment -III

Posa Harsha vardhan(EE22BTECH11214)*

Question: Let X represent the difference between the number of heads and the number of tails obtained when a coin is tossed 6 times. What are possible values of X?

Solution:

Let H be a random variable which represents the number of Heads obtained in 6 coin tosses.

And T be a random variable which represents the number of Tails obtained in 6 coin tosses. Then,

 $H \in \{0, 1, 2, 3, 4, 5, 6\}$

Similarly,

 $T \in \{0, 1, 2, 3, 4, 5, 6\}$

$$H + T = 6 \tag{1}$$

$$X = |H - T| \tag{2}$$

$$X = |H - (6 - H)| \tag{3}$$

$$X = |2H - 6| \tag{4}$$

$$F_X(k) = \Pr(X \le 0) = \Pr(|2H - 6| \le k)$$
 (5)

$$= \Pr(2H - 6 \le k) \text{ and } \Pr(2H - 6 \ge -k)$$
 (6)

$$= \Pr\left(H \le \frac{k+6}{2}\right) \text{ and } \Pr\left(H \ge \frac{6-k}{2}\right) \tag{7}$$

$$=\Pr\left(\frac{6-k}{2} \le H \le \frac{k+6}{2}\right) \tag{8}$$

$$= F_H \left(\frac{6+k}{2} \right) - F_H \left(\frac{6-k}{2} - 1 \right) \tag{9}$$

$$=F_H\left(\frac{6+k}{2}\right) - F_H\left(\frac{4-k}{2}\right) \tag{10}$$

$$F_X(0) = F_H(3) - F_H(2) \tag{11}$$

$$=^{6} C_{3} \left(\frac{1}{2}\right)^{6} \tag{12}$$

$$F_X(2) = F_H(4) - F_H(1) \tag{13}$$

$$= {}^{6}C_{2}\left(\frac{1}{2}\right)^{6} + {}^{6}C_{3}\left(\frac{1}{2}\right)^{6} + {}^{6}C_{4}\left(\frac{1}{2}\right)^{6}$$
 (14)

$$=\frac{25}{32}$$
 (15)

$$F_X(4) = F_H(5) - F_H(0) \tag{16}$$

$$= {}^{6}C_{1}\left(\frac{1}{2}\right)^{6} + {}^{6}C_{2}\left(\frac{1}{2}\right)^{6} + {}^{6}C_{3}\left(\frac{1}{2}\right)^{6} \tag{17}$$

$$+^{6} C_{4} \left(\frac{1}{2}\right)^{6} +^{6} C_{5} \left(\frac{1}{2}\right)^{6} \tag{18}$$

$$=\frac{31}{32}$$
 (19)

$$F_X(6) = F_H(6) (20)$$

$$= {}^{6} C_{0} \left(\frac{1}{2}\right)^{6} + {}^{6} C_{1} \left(\frac{1}{2}\right)^{6} + {}^{6} C_{2} \left(\frac{1}{2}\right)^{6} + {}^{6} C_{3} \left(\frac{1}{2}\right)^{6}$$

$$+^{6} C_{4} \left(\frac{1}{2}\right)^{6} +^{6} C_{5} \left(\frac{1}{2}\right)^{6} +^{6} C_{6} \left(\frac{1}{2}\right)^{6}$$
 (21)

$$=\frac{64}{64}=1$$
 (22)