Experiment 7 - Random Sampling and Sampling Distribution

Aim:

To explore random sampling from a population and understand the concept of sampling distribution.

Program:

```
import numpy as np
import matplotlib.pyplot as plt

# Generating a population
population_mean = 50
population_std = 10
population_size = 100000

population = np.random.normal(
    population_mean,
    population_std,
    population_size
)

# Random sampling
sample_sizes = [30, 50, 100]
num_samples = 1000
sample_means = {}

for size in sample_sizes:
    sample_means[size] = []

    for _ in range(num_samples):
        sample = np.random.choice(
```

```
            population,
            size=size,
            replace=False
        )
        sample_means[size].append(np.mean(sample))

# Plotting sampling distributions
plt.figure(figsize=(12, 8))

for i, size in enumerate(sample_sizes):
    plt.subplot(len(sample_sizes), 1, i + 1)


    plt.hist(
        sample_means[size],
        bins=30,
        alpha=0.7,
        label=f'Sample Size {size}'
    )

    plt.axvline(
        np.mean(population),
        color='red',
        linestyle='dashed',
        linewidth=1.5,
        label='Population Mean'
    )

    plt.title(f'Sampling Distribution (Sample Size {size})')
    plt.xlabel('Sample Mean')
    plt.ylabel('Frequency')
    plt.legend()
```

```
plt.tight_layout()
plt.show()
```

Output:



Result:

Thus, random sampling from a population has been understood and explored.