Journal of Psychiatric Research Manuscript Draft

Manuscript Number: JPR5140

Title: Anxiety symptoms in bipolar offspring with the brain-derived neurotrophic factor (BDNF) Val66Met genotype

Article Type: Original Article

Keywords: Risk; Pediatric bipolar; Gene-Environment Interaction; Brain-Derived Neurotrophic Factor; Anxiety

Abstract: Several genetic and environmental factors place youth offspring of parents with bipolar disorder (BD) at high risk for developing mood and anxiety disorders. Recent studies suggest that anxiety symptoms, even at subclinical levels, have been associated with an increased risk for developing BD. The brain-derived neurotrophic factor (BDNF) gene has been implicated in the pathophysiology of both BD and anxiety disorders. We aimed to explore whether anxiety in BD offspring was associated with the BDNF Val66Met polymorphism. 64 BD offspring (mean age: 13.73 (S.D. 3.45) M=30, F=34) and 51 HC (mean age: 13.68 (S.D. 2.68) M=23, F=28) were compared on presence of the BDNF Val66Met polymorphism and on scores from the Multidimensional Anxiety Scale for Children (MASC). To assess family function, we used the Family Adaptability and Cohesio Evaluation Scales (FACES-IV). BD offspring showed higher levels of overall anxiety than did the HC group. BD offspring with the val/val genotype showed higher levels of anxiety than BD offspring with other genotypes. No significant levels of anxiety or its association with BDNF genotype were found in the HC group. BD offspring group showed significantly more family dysfunction when compared with the HC group and the family dysfunction moderated the interaction between the BDNF genotype and anxiety symptoms. This study demonstrated an association between the BDNF genotype and anxiety in BD offspring, moderated by family dysfunction. Longitudinal studies in high-risk offspring will aid in elucidating how these phenomenological, environmental, and genetic factors converge to the development of BD.

Acknowledgement

Acknowledgements

The authors gratefully acknowledge funding from the NIMH (R01 MH077047 and K23 MH064460 to KDC and K23 MH085919 to MKS). We thank the participants who gave their time for this study.

*Conflict of Interest

Conflicts of interest

Dr. Chang is a consultant for GlaxoSmithKline, Bristol-Myers Squibb, Merck, and Sunovion. The other authors have nothing to disclose.

*Contributors

Contributors:

Drs Singh and Chang designed the study, wrote the protocol, and wrote and edited all drafts of the manuscript. Dr. Park conducted the literature search, analyses, and wrote the first draft of the manuscript. Dr. Hong contributed to the statistical analysis. Drs. Park, Chang, Hallmayer, Kim, Hong and Singh, and Ms. Howe contributed to and approved the final manuscript.





April 30, 2014

Alan Schatzberg, MD
Department of of Psychiatry & Behavioral Sciences
Stanford University School of Medicine,
401 Quarry Road, Stanford, CA 94305-5117

Dear Dr. Schatzberg,

We are submitting our manuscript, "Anxiety symptoms in bipolar offspring with the brain-derived neurotrophic factor (BDNF) Val66Met genotype," for consideration for publication in the *Journal of Psychiatric Research*. This manuscript demonstrates an association between the BDNF genotype and anxiety in youth offspring of parents with bipolar disorder, moderated by family dysfunction. We have not previously submitted this or any related manuscript for publication. None of the authors have commercial or any other financial interests in the results reported. All of the authors contributed significantly in this work so as to take public responsibility for this paper's contents. Given their expertise in the clinical at risk population or psychiatric genetics described in our study, we request the following individuals as potential reviewers for this paper:

- 1) Melissa DelBello, M.D., Professor, Division of Bipolar Disorders Research, Department of Psychiatry, University of Cincinnati College of Medicine, 231 Albert Sabin Way, PO Box 670559, Cincinnati, OH 45267-0559, USA. delbelmp@ucmail.uc.edu; Ph: (513) 558-6194; Fax: (513) 558-3399.
- 2) **Jair C. Soares, M.D.**, Professor and Chairman, Department of Psychiatry and Behavioral Sciences, UT Houston Medical School, 1300 Moursund St., Houston, TX 77030; Ph: (713) 500-2507; Fax: (713) 500-2553; Email: jair.c.soares@uth.tmc.edu
- 3) Mani Pavuluri, M.D., Professor and Chair of Child Psychiatry. Director, Pediatric Mood Disorders Clinic, University of Illinois at Chicago, Department of Psychiatry, Center for Cognitive Medicine (MC913), 912 S. Wood St., Chicago IL 60612-7327 USA; Ph: (312) 413-1722; Fax: (312) 413-8837 Email: mpavuluri@psych.uic.edu
- 4) **Fernando Goes, M.D.,** Assistant Professor of Psychiatry and Behavioral Sciences, Johns Hopkins University, Meyer 4-119D, The Johns Hopkins Hospital 600 North Wolfe Street Baltimore, MD 21287 USA; Phone: 443-287-6382; Fax: 410-502-0065; E-mail: fgoes@jhmi.edu

Thank you for your consideration. We look forward to hearing from you.

Sincerely,

Manpreet K. Singh, MD MS

Mang 1 K 8

Assistant Professor of Psychiatry and Behavioral Sciences

Stanford University School of Medicine

*Role of the Funding Source

Role of the funding source:

Funding for this study was provided by the NIMH (R01 MH077047 and K23 MH064460 to KDC and K23 MH085919 to MKS). The NIMH had no further role in study design, in the collection, analysis and interpretation of data, in the writing of the report, or in the decision to submit the paper for publication.

*Highlights (for review)

Highlights

BD offspring group had higher levels of anxiety than the HC group.

BD offspring group had significantly more family dysfunction than the HC group.

BD offspring with the val/val genotype of the BDNF gene had higher anxiety than the other genotypes.

Family dysfunction moderated the association between the BDNF genotype and anxiety.

Text Word Count: 3,609

Abstract Word Count: 246

Tables: 2

Figures: 2

Anxiety symptoms in bipolar offspring with the brain-derived neurotrophic factor (BDNF) Val66Met genotype

Min-Hyeon Park^a, Kiki D. Chang^{b, *}, Joachim Hallmayer^b, Meghan E. Howe^b, Eunjoo Kim^c, Seung Chul Hong^a, Manpreet K. Singh^b

^aDepartment of Psychiatry, The Catholic University of Korea, St. Vincent Hospital, Suwon, Korea

^bDepartment of Psychiatry and Behavioral Sciences, Stanford University School of Medicine, Stanford, CA, USA

^cDepartment of Psychiatry and Institute of Behavioral Science in Medicine Yonsei University College of Medicine, Seoul, Korea

*Corresponding Author: Manpreet K. Singh, M.D. M.S.

Stanford University School of Medicine, Division of Child and Adolescent Psychiatry, 401 Quarry Road, Stanford, CA 94305-5719, USA

Tel: (650) 725-5922, Fax: (650) 723-5531, E-mail: mksingh@stanford.edu

ABSTRACT

Several genetic and environmental factors place youth offspring of parents with bipolar

disorder (BD) at high risk for developing mood and anxiety disorders. Recent studies suggest

that anxiety symptoms, even at subclinical levels, have been associated with an increased risk

for developing BD. The brain-derived neurotrophic factor (BDNF) gene has been implicated

in the pathophysiology of both BD and anxiety disorders. We aimed to explore whether

anxiety in BD offspring was associated with the BDNF Val66Met polymorphism. 64 BD

offspring (mean age: 13.73 (S.D. 3.45) M=30, F=34) and 51 HC (mean age: 13.68 (S.D.

2.68) M=23, F=28) were compared on presence of the BDNF Val66Met polymorphism and

on scores from the Multidimensional Anxiety Scale for Children (MASC). To assess family

function, we used the Family Adaptability and Cohesion Evaluation Scales (FACES-IV).

BD offspring showed higher levels of overall anxiety than did the HC group. BD offspring

with the val/val genotype showed higher levels of anxiety than BD offspring with other

genotypes. No significant levels of anxiety or its association with BDNF genotype were

found in the HC group. BD offspring group showed significantly more family dysfunction

when compared with the HC group and the family dysfunction moderated the interaction

between the BDNF genotype and anxiety symptoms. This study demonstrated an association

between the BDNF genotype and anxiety in BD offspring, moderated by family dysfunction.

Longitudinal studies in high-risk offspring will aid in elucidating how these

phenomenological, environmental, and genetic factors converge to the development of BD.

Keywords: Risk; Pediatric bipolar; Gene-Environment Interaction; Brain-Derived

Neurotrophic Factor; Anxiety

2

1. Introduction

Children of parents with bipolar disorder (BD) (BD offspring) are at higher risk for developing BD (Axelson et al., 2011; Chang et al., 2003) as well as other psychiatric disorders (Lapalme et al., 1997; Singh et al., 2007) than the general population. Anxiety disorders have attracted special attention not only because of their high prevalence in BD offspring but also because the presence of anxiety disorders may serve as a predictor for developing BD (Duffy et al., 2013; Henin et al., 2005) iven subclinical levels of anxiety can predispose individuals to developing BD, suggesting that having early anxiety symptoms may represent an endophenotype of BD (Contreras et al., 2010).

Decrease ain-derived neurotrophic factor (BDNF) levels have been implicated in both bipolar and anxiety disorders. BDNF is a member of the neurotrophin family of growth factors and is concentrated in brain regions involved in learning and memory, such as the hippocampus and amygdala (Hofer et al., 1990; Rakofsky et al., 2012). Reduced (BDNF) levels have been observed in patients with BD during manic and depressive episodes and these levels have normalized with episode recovery (Cunha et al., 2006; de Oliveira et al., 2009; Lin, 2009). Post-mortem studies have also demonstrated decreased hippocampal BDNF in patients with BD (Dunham et al., 2009; Knable et al., 2004). Similarly, reduced BDNF levels have been observed in individuals with anxiety disorders compared to those without anxiety disorders (Suliman et al., 2013). Finally, acute and chronic stress have been associated with decreased BDNF expression in the hippocampus with subsequent enhancement of anxiety-related behaviors (McEwen, 2001; Russo-Neustadt, 2003). Youth offspring of parents with BD are biologically sensitive to stre striguy et al., 2009; Ostiguy et al., 2011) and typically live in disorganized and conflictual family environments (Belardinelli et al., 2008; Chang et al., 2001).

Decreased BDNF levels in humans has also been associated with the met allele of

BDNF Val66Met polymorphism (rs6265) (Chen et al., 2006; Egan et al., 2003). However, studies on the association between the BDNF Val66Met polymorphism and BD or anxiety have been mixed, with a number of studies that have also found that the val/val allele also confers risk for anxiety or B his inconsistency across studies may be due to the heterogenous nature of these disorders or unidentified moderating factors between the BDNF genotype and psychiatric symptoms.

Although, anxiety may serve as a predisposing factor for BD development in BD offspring, no study has investigated the relation between anxiety and the BDNF genotype in cohorts at high-risk for BD, or the moderating effects of family dysfunction on this relation. Therefore, we aimed to compare levels of anxiety, as measured by the Multidimensional Anxiety Scale for Children (MASC) (March et al., 1997), family dysfunction, measured by the Family Adaptability and Cohesion Evaluation Scales, version IV (FACES-IV) (Olson, 2011), and their associations with the BDNF genotype in BD offspring and healthy control (HC) groups. We hypothesized that (1) BD offspring would have higher levels of anxiety than the HC group; (2) levels of anxiety would interact with the BDNF Val66Met genotype in BD offspring but not in healthy controls, such that group status would moderate the relation between the BDNF Val66Met genotype and anxiety; and (3) compared to HC youth, youth offspring of parents with BD would have lower levels of family cohesion, organization, and satisfaction, and that these dysfunctional family characteristics would moderate the effect of the BDNF genotype on anxiety symptoms.

2. Materials and methods

2.1. Participants

The protocol was approved by the Stanford University Panel of Medical Research in Human Subjects. One hundred BD offspring and 60 HC were recruited from an ongoing

study of youth at risk for BD and from the community. After obtaining oral and written informed consent and assent from parents from their offspring, respectively, we conducted semi-structured interviews to assess for BD in parents and psychopathology in their offspring. Parents were diagnosed using the Structured Clinical Interview for DSM-IV Axis I Disorders (SCID) (First et al., 1996), and were interviewed for psychiatric history of firstand second-degree relatives following the Family History-Research Diagnostic Criteria (FH-RDC) (Andreasen et al., 1977). BD offspring were assessed using the Affective Disorders Module of the Washington Schedule for Affective Disorders and Schizophrenia for School-Age Children (WASH-U-KSADS) (Geller et al., 2001). All evaluations were conducted by either a child and adolescent psychiatrist or a trained master's level research assistant with at least two years experience in psychiatric interviewing. Evaluators were aware of parental diagnosis, who were interviewed while they were euthymic at the time of their child's assessment. Inter-rater reliability was established by rating videotaped interviews, observing a trained rater, observing trained rater interviews, and performing interviews with observation by a trained rater. Diagnostic decisions were ultimately made by a board-certified child and adolescent psychiatrist (K.C. or M.S.) based on personal interview, discussion with the research assistant and written notes of parental and offspring responses to interview questions. Current and lifetime diagnoses were established according to DSM-IV criteria (American Psychiatric Association, 1994).

Youths aged 9 to 17 having at least one biological parent with BD were included in the BD offspring group. BD offspring with the following disorders or conditions were excluded: BD I or II, Tourette's disorder, substance use disorder, or a neurological condition (such as a seizure disorder). HC had no personal or parental psychiatric history and no family history of BD, and were excluded with current Young Mania Rating Scale (YMRS) (Young et al., 1978) scores greater than 8 or Childhood Depression Rating Scale (CDRS-R)

(Poznanski et al., 1985) greater than 26.

2.2. Instruments

2.2.1. *Multidimensional Anxiety Scale for Children (MASC):* To assess anxiety symptoms, we used the MASC, a 39-item scale, distributed into four major factors: physical symptoms, social anxiety, harm avoidance, and separation anxiety. The MASC has been shown to distinguish anxiety from depression, and differentiate between various anxiety disorders (Dierker et al., 1999).

2.2.2. Family Adaptability and Cohesion Evaluation Scales, version IV (FACES-IV): To assess family function, we used the FACES-IV, a questionnaire completed by parents participating in this study. The FACES-IV includes 42 items loading on six subscales: balanced cohesion, balanced flexibility, disengaged, enmeshed, rigid, and chaotic. The family communication and family satisfaction scales are additional scales and they indicate how healthy the communication with family is and how satisfied individual's are in regard to life with their family. High family satisfaction is usually related to higher levels of balanced cohesion and flexibility and lower levels of the four unbalanced scales (disengaged, enmeshed, rigid, and chaotic).

2.3. Genotyping

DNA was obtained from peripheral venous blood (20ml). All genotyping assays were conducted within the laboratory of Dr. Joachim Hallmayer at Stanford by the same technician blinded to all clinical data. BDNF val66met genotyping was conducted as per the laboratory's standard protocol, utilizing the following primers for G196A in the BDNF gene: forward 5'-ATC CGA GGA CAA GGT GGC-3' and reverse 5'-CCT CAT GGA CAT GTT

TGC AG-3'. This generated 300 bp of polymerized chain reaction (PCR) products, subsequently digested by Pml I (New England Biolabs, Ipswich, MA) to yield either met allele (undigested, 300 bp) or val allele (digested to 180 bp b 120 bp bands), which were visualized on 7% polyacrylamide gel using a 50 bp marker.

2.4. Statistical analysis

Since participants with the met/met were rare, the BDNF Val66Met genotype was dichotomized according to whether or not they possessed the rare met allele (Kaufman et al., 2006; Kim et al., 2007). To compare categorical variables, we conducted χ^2 tests. To compare continuous variables, we conducted independent t test for parametric data or Mann-Witney U test for non-parametric data. The statistical significance for all tests was set at p < 0.05. We corrected significance levels for multiple comparisons using a Bonferroni correction for MASC subscales(0.05/5 subscales = 0.01) and for FACES-IV subscales (0.05/8 subscales = 0.00625). We used Baron & Kenny's method to examine the moderating effect between the independent and dependent variables (Baron and Kenny, 1986).

3. Results

Among 100 BD offspring and 60 HC, 36 BD offspring and 9 HC were excluded due to incomplete MASC data, leaving 64 BD offspring (mean age: 13.73 (3.45) M=30, F=34) and 51 HC (mean age: 13.68 (2.68) M=23, F= 28) to be included for statistical analysis.

Participant demographic and clinical characteristics are provided in Table 1. The BD offspring group showed significantly higher anxiety scores than the HC group in the physical subscale (p=0.001), social anxiety scale (p=0.037), and in the overall MASC score (p=0.018). Results for the physical subscale and MASC total scores remained significant after adjusting for multiple comparisons (Table 1, Fig. 1-a).

Parent reports on FACES-IV data was available for 22 BD offspring and 28 HC. There were no differences in demographic and clinical characteristics between those who provided and those who didn't provide FACES-IV data both in the BD offspring and HC groups. The BD offspring group showed significantly more family dysfunction when compared with the HC group with lower levels of cohesion (p=0.006), higher levels of enmeshment (p=0.002) and chaos (p=0.049), and lower levels of family satisfaction (p=0.003) and communication (p=0.001). Results for all FACES subscales remained significant after adjusting for multiple comparisons.

The val/val genotype of the BDNF Val66Met polymorphism was observed in 34 subjects (53.1 %), the val/met genotype in 27 subjects (42.2%), and the met/met genotype in 3 subjects (4.7 %) in the BD offspring group and the val/val genotype was observed in 34 subjects (66.7 %), the val/met genotype in 15 subjects (29.4%), and the met/met genotype in 2 subjects (3.9 %) in the HC group. The distributions of genotypes for the Val66Met polymorphism were consistent with expected values based on Hardy-Weinberg equilibrium in both the BD offspring (p<0.164) and HC groups (p<0.997).

When all subjects were dichotomized according to the presence or absence of the met allele (val/val vs. val/met + met/met), no significant difference was observed in the MASC scores for each BDNF genotype. There were also no significant differences in the MASC total or subscales by BDNF genotype within the HC group. However, within the BD offspring group, subjects with the val/val genotype showed significantly higher anxiety scores than subjects with the val/met or met/met genotype in the harm avoidance subscale (p=0.005), social anxiety subscale (p=0.019) and total MASC scores (p=0.010). Results for the harm avoidance subscale and MASC total scores remained significant after adjusting for multiple comparisons (Table 2, Fig. 1-b).

We examined the moderating effect of being BD offspring on anxiety symptoms by BDNF genotype using the Baron & Kenny method. In step 1, we conducted a regression analysis in which we analyzed the effect of the BDNF genotype and BD offspring status on the MASC total score, social anxiety subscale and harm avoidance subscale. Then, in step 2, we input the interaction variable between the BDNF genotype and BD offspring into the variables used in the step 1 analysis. The step 1 analysis revealed that the BDNF genotype did not have an effect on the harm avoidance subscale (p>0.05), social anxiety subscale (p>0.05) or total MASC scores (p>0.05). However, in step 2, we found a moderating effect of being BD offspring, in which BDNF genotype had an effect on the harm avoidance subscale (B=-4.147, p<0.01), social anxiety subscale (B=-4.582, p<0.05) and total MASC scores (B=-12.500, p<0.05) (Fig. 2). Thus, the BDNF genotype itself did not affect anxiety scores. However, when the BDNF genotype was combined with BD offspring status, the BDNF genotype had an effect on anxiety scores, suggesting that BD offspring status moderated the interaction between the BDNF genotype and anxiety.

We also examined the moderating effect of the FACES-IV scores on social anxiety symptoms by BDNF genotype using the Baron & Kenny method. The step 1 analysis revealed that the BDNF genotype did not have an effect on social anxiety subscale However, in step 2, we found a moderating effect of enmeshed, communication or satisfaction scores, in which BDNF genotype had an effect on social anxiety subscale (B=2.091, B=1.530 or B=1.090 respectively, p<0.05). Thus, the BDNF genotype itself did not affect social anxiety score. However, when the BDNF genotype was combined with the FACES-IV scores, the BDNF genotype had an effect on social anxiety score, suggesting that the FACES-IV scores moderated the interaction between the BDNF genotype and social anxiety.

Our main findings were not affected by the current usage of anti-anxiety or stimulant medications (Table 2). When subsequent analyses were performed excluding the subjects

who were taking anti-anxiety medication and stimulants respectively, similar results were found.

4. Discussion

In this study, we demonstrated that as hypothesized, BD offspring have higher levels of anxiety than typically developing healthy controls, and anxiety symptoms interacted with BDNF genotype in BD offspring but not in healthy controls. Specifically, BD offspring with the val/val genotype showed higher anxiety symptoms than BD offspring with other genotypes. Further, we found BD offspring group showed significantly more family dysfunction when compared with the HC group and the family dysfunction moderated the interaction between the BDNF genotype and anxiety symptoms. These findings have implications for early identification of certain genetic and environmental vulnerabilities for developing BD in youth at risk for BD.

Among the MASC subscales, the physical subscale showed the most prominent difference between the BD offspring and HC groups. The physical subscale score can be affected by elevated autonomic nervous system activity, release of cortisol and epinephrine, and amygdala excitability (McTeague and Lang, 2012). Further, the physical subscale has been shown to predict the future development of panic disorder and agoraphobia (van Gastel and Ferdinand, 2008). These findings suggest that youth offspring of BD parents may have an increased biological predisposition for anxiety disorders that manifests most prominently by physical symptoms of anxiety.

Interestingly, the harm avoidance and social anxiety subscales showed the most prominent difference between BDNF Val66Met genotype groups in the BD offspring group. The harm avoidance and social anxiety subscales have previously been related to subclinical anxiety symptoms and traits, but neither has been fully accurate in predicting a DSM-

diagnosis of an anxiety disorder (van Gastel and Ferdinand, 2008). The harm avoidance subscale has also been related to perfectionism and anxiety coping, whereas the social anxiety subscale has been related to humiliation and public performance fears (March et al., 1997). Therefore, it may be assumed that high scores on those subscales are related to subclinical range of anxiety symptoms or anxiety trait rather than an anxiety disorder.

Prior studies on the association between the BDNF genotype and anxiety have shown inconsistent results. Interestingly, most studies have not demonstrated an increased incidence of anxiety disorders by BDNF genotype, including studies of obsessive-compulsive disorder, panic disorder, generalized- anxiety disorders, or phobias (Enoch et al., 2008; Shimizu et al., 2005; Surtees et al., 2007; Wendland et al., 2007; Zhang et al., 2006). A meta-anaylsis relating anxiety to BDNF genotype in adults with anxiety or anxiety-related personality traits was also negative (Frustaci et al., 2008) An association between higher anxiety trait and the val/val genotype has been found in healthy individuals and in animals (Hünnerkopf et al., 2007; Lang et al., 2004). Others have demonstrated an association between higher anxiety trait and the met allele (Colzato et al., 2011; Jiang et al., 2005; Montag et al., 2010; Soliman et al., 2010), while yet others have shown no relation between anxiety and the BDNF genotype (Baekken et al., 2008; Tochigi et al., 2006). These inconsistencies may result from significant symptom heterogeneity among their participants and environmental factors such as family functioning.

Prior studies have also been inconsistent about the relation between BD and the Val66Met polymorphism of BDNF gene. Some family-based association studies reported significant over-transmission of the val allele in patients with BD (Geller et al., 2004; Neves-Pereira et al., 2002; Sklar et al., 2002; Strauss et al., 2005; Tang et al., 2008) whereas others have found no relation between BD and the Val66Met polymorphism of BDNF gene (Green et al., 2006; Hong et al., 2003; Kunugi et al., 2004; Nakata et al., 2003; Oswald et al., 2004;

Skibinska et al., 2004). Some studies have found that the val allele is associated with early age of onset of BD (Geller et al., 2004; Strauss et al., 2005; Tang et al., 2008), rapid cycling (Green et al., 2006; Müller et al., 2006) and suicide attempts (Vincze et al., 2008). However, other studies have linked the met allele to early age of BD onset (Skibinska et al., 2004), suicide attempts (Kim et al., 2008), and structural or functional abnormalities in areas important for the regulation of mood and anxiety, such as the hippocampus and amygdala (Bueller et al., 2006; Egan et al., 2003; Frodl et al., 2007; Hariri et al., 2003; Montag et al., 2009; Nemoto et al., 2006; Soliman et al., 2010). Thus, while BDNF genotype appears to be relevant to mood and anxiety status, the exact relationship is not clear, and it may be that environmental factors moderate the effects of BDNF genotype on behavioral outcome.

Perhaps the most significant environmental factor in BD offspring is the family environment, as families with a BD parent have been noted to have high levels of conflict and lower levels of cohesion (Barron et al., 2014; Chang et al., 2001; Romero et al., 2005). These studies found that families with a bipolar parent are chaotic, have lower family satisfaction, and lower family communication. As familial influences on defense and coping mechanisms have been identified as important mediators of psychopathology (Thienemann et al., 1998), gene by environment interaction effects may cause epigenetic modulation in BD offspring that might explain our findings in the context of exposure to family stress. For example, Gatt et al. (2009) showed a significant interaction between Val66Met polymorphism and early life stress. Animal studies have also shown that early life traumas can affect the epigenetic regulation of BDNF genes (Bazak et al., 2009; Roth et al., 2009). Such stressors have been found to cause reduced expression of BDNF, which is mediated by stress-related hormones such as catecholamines and glucocorticoids (Carbone and Handa, 2013; Duman, 2002; Hartmann et al., 2001; Rasmusson et al., 2002).

Finally, even though not many BD offspring with the val/val genotype showed clinical

levels of anxiety symptoms as defined by structured clinical interview, many of BD offspring with the val/val genotype showed subclinical anxiety symptoms. It is possible that subclinical anxiety symptoms may represent early prodromal symptoms of BD, as well as potential early forms of anxiety disorders. Future longitudinal studies would clarify the relation between these early symptoms and the development of BD.

There are some limitations in our study that are worth considering. First, this study is a cross-sectional study that will require longitudinal follow up to determine whether anxiety traits predict conversion to BD in youth at familial risk. Second, the number of subjects is relatively small for genetic analysis, and our sample contains multiple ethnic groups that could account for variability in our sample. However, our results did not change when we re-analyzed our data after excluding non-Caucasian participants.

It is important to identify comprehensive risk factors for BD development in these high-risk offspring. We found that our BD offspring group showed significantly more family dysfunction when compared with the HC group, and that the family dysfunction moderated the interaction between the BDNF genotype and anxiety symptoms. Early identification of specific environmental and genetic risk factors will enable clinicians to perform earlier interventions to prevent the onset of mood problems in high-risk youth.

References

- American Psychiatric Association. Diagnostic and statistical manual of mental disorders:

 DSM-IV. 4th ed. Washington, DC: American Psychiatric Association; 1994.
- Andreasen NC, Endicott J, Spitzer RL, Winokur G. The family history method using diagnostic criteria. Reliability and validity. Arch Gen Psychiatry 1977;34:1229-35.
- Axelson DA, Birmaher B, Strober MA, Goldstein BI, Ha W, Gill MK, et al. Course of subthreshold bipolar disorder in youth: diagnostic progression from bipolar disorder not otherwise specified. Journal of the American Academy of Child and Adolescent Psychiatry 2011;50:1001-16 e3.
- Baekken PM, Skorpen F, Stordal E, Zwart JA, Hagen K. Depression and anxiety in relation to catechol-O-methyltransferase Val158Met genotype in the general population: the Nord-Trøndelag Health Study (HUNT). BMC Psychiatry 2008;8:48.
- Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. J Pers Soc Psychol 1986;51:1173-82.
- Barron E, Sharma A, Le Couteur J, Rushton S, Close A, Kelly T, et al. Family environment of bipolar families: a UK study. Journal of Affective Disorders 2014;152-154:522-5.
- Bazak N, Kozlovsky N, Kaplan Z, Matar M, Golan H, Zohar J, et al. Pre-pubertal stress exposure affects adult behavioral response in association with changes in circulating corticosterone and brain-derived neurotrophic factor. Psychoneuroendocrinology 2009;34:844-58.
- Belardinelli C, Hatch JP, Olvera RL, Fonseca M, Caetano SC, Nicoletti M, et al. Family environment patterns in families with bipolar children. J Affect Disord 2008;107:299-305.
- Bueller JA, Aftab M, Sen S, Gomez-Hassan D, Burmeister M, Zubieta JK. BDNF Val66Met

- allele is associated with reduced hippocampal volume in healthy subjects. Biol Psychiatry 2006;59:812-5.
- Carbone DL, Handa RJ. Sex and stress hormone influences on the expression and activity of brain-derived neurotrophic factor. Neuroscience 2013;239:295-303.
- Chang K, Steiner H, Ketter T. Studies of offspring of parents with bipolar disorder. Am J Med Genet C Semin Med Genet 2003;123C:26-35.
- Chang KD, Blasey C, Ketter TA, Steiner H. Family environment of children and adolescents with bipolar parents. Bipolar Disord 2001;3:73-8.
- Chen ZY, Jing D, Bath KG, Ieraci A, Khan T, Siao CJ, et al. Genetic variant BDNF (Val66Met) polymorphism alters anxiety-related behavior. Science 2006;314:140-3.
- Colzato LS, Van der Does AJ, Kouwenhoven C, Elzinga BM, Hommel B. BDNF Val66Met polymorphism is associated with higher anticipatory cortisol stress response, anxiety, and alcohol consumption in healthy adults. Psychoneuroendocrinology 2011;36:1562-9.
- Contreras J, Hare E, Pacheco A, Escamilla M, Raventos H. Is subclinical anxiety an endophenotype for bipolar I patients? A study from a Costa Rican sample. J Affect Disord 2010;122:267-72.
- Cunha AB, Frey BN, Andreazza AC, Goi JD, Rosa AR, Gonçalves CA, et al. Serum brainderived neurotrophic factor is decreased in bipolar disorder during depressive and manic episodes. Neurosci Lett 2006;398:215-9.
- de Oliveira GS, Ceresér KM, Fernandes BS, Kauer-Sant'Anna M, Fries GR, Stertz L, et al.

 Decreased brain-derived neurotrophic factor in medicated and drug-free bipolar patients. J Psychiatr Res 2009;43:1171-4.
- Dierker LC, Merikangas KR, Szatmari P. Influence of parental concordance for psychiatric disorders on psychopathology in offspring. J Am Acad Child Adolesc Psychiatry

- 1999;38:280-8.
- Duffy A, Horrocks J, Doucette S, Keown-Stoneman C, McCloskey S, Grof P. Childhood anxiety: an early predictor of mood disorders in offspring of bipolar parents. Journal of Affective Disorders 2013;150:363-9.
- Duman RS. Synaptic plasticity and mood disorders. Mol Psychiatry 2002;7 Suppl 1:S29-34.
- Dunham JS, Deakin JF, Miyajima F, Payton A, Toro CT. Expression of hippocampal brainderived neurotrophic factor and its receptors in Stanley consortium brains. J Psychiatr Res 2009;43:1175-84.
- Egan MF, Kojima M, Callicott JH, Goldberg TE, Kolachana BS, Bertolino A, et al. The BDNF val66met polymorphism affects activity-dependent secretion of BDNF and human memory and hippocampal function. Cell 2003;112:257-69.
- Enoch MA, White KV, Waheed J, Goldman D. Neurophysiological and genetic distinctions between pure and comorbid anxiety disorders. Depress Anxiety 2008;25:383-92.
- First MB, Spitzer RL, Gibbon M, Williams JBW. Structured Clinical Interview for DSM-IV

 Axis I Disorders Patient Edition (SCID-I/P, Version 2.0). New York: Biometrics

 Research Department, New York State Psychiatric Institute; 1996.
- Frodl T, Schüle C, Schmitt G, Born C, Baghai T, Zill P, et al. Association of the brainderived neurotrophic factor Val66Met polymorphism with reduced hippocampal volumes in major depression. Arch Gen Psychiatry 2007;64:410-6.
- Frustaci A, Pozzi G, Gianfagna F, Manzoli L, Boccia S. Meta-analysis of the brain-derived neurotrophic factor gene (BDNF) Val66Met polymorphism in anxiety disorders and anxiety-related personality traits. Neuropsychobiology 2008;58:163-70.
- Gatt JM, Nemeroff CB, Dobson-Stone C, Paul RH, Bryant RA, Schofield PR, et al.

 Interactions between BDNF Val66Met polymorphism and early life stress predict brain and arousal pathways to syndromal depression and anxiety. Mol Psychiatry

- 2009;14:681-95.
- Geller B, Badner JA, Tillman R, Christian SL, Bolhofner K, Cook EH Jr. Linkage disequilibrium of the brain-derived neurotrophic factor Val66Met polymorphism in children with a prepubertal and early adolescent bipolar disorder phenotype. Am J Psychiatry 2004;161:1698-700.
- Geller B, Zimerman B, Williams M, Bolhofner K, Craney JL, DelBello MP, et al. Reliability of the Washington University in St. Louis Kiddie Schedule for Affective Disorders and Schizophrenia (WASH-U-KSADS) mania and rapid cycling sections. J Am Acad Child Adolesc Psychiatry 2001;40:450-5.
- Green EK, Raybould R, Macgregor S, Hyde S, Young AH, O'Donovan MC, et al. Genetic variation of brain-derived neurotrophic factor (BDNF) in bipolar disorder: case-control study of over 3000 individuals from the UK. Br J Psychiatry 2006;188:21-5.
- Hünnerkopf R, Strobel A, Gutknecht L, Brocke B, Lesch KP. Interaction between BDNF Val66Met and dopamine transporter gene variation influences anxiety-related traits. Neuropsychopharmacology 2007;32:2552-60.
- Hariri AR, Goldberg TE, Mattay VS, Kolachana BS, Callicott JH, Egan MF, et al. Brainderived neurotrophic factor val66met polymorphism affects human memory-related hippocampal activity and predicts memory performance. J Neurosci 2003;23:6690-4.
- Hartmann M, Heumann R, Lessmann V. Synaptic secretion of BDNF after high-frequency stimulation of glutamatergic synapses. EMBO J 2001;20:5887-97.
- Henin A, Biederman J, Mick E, Sachs GS, Hirshfeld-Becker DR, Siegel RS, et al.

 Psychopathology in the offspring of parents with bipolar disorder: a controlled study.

 Biol Psychiatry 2005;58:554-61.
- Hofer M, Pagliusi SR, Hohn A, Leibrock J, Barde YA. Regional distribution of brain-derived neurotrophic factor mRNA in the adult mouse brain. EMBO J 1990;9:2459-64.

- Hong CJ, Huo SJ, Yen FC, Tung CL, Pan GM, Tsai SJ. Association study of a brain-derived neurotrophic-factor genetic polymorphism and mood disorders, age of onset and suicidal behavior. Neuropsychobiology 2003;48:186-9.
- Jiang X, Xu K, Hoberman J, Tian F, Marko AJ, Waheed JF, et al. BDNF variation and mood disorders: a novel functional promoter polymorphism and Val66Met are associated with anxiety but have opposing effects. Neuropsychopharmacology 2005;30:1353-61.
- Kaufman J, Yang BZ, Douglas-Palumberi H, Grasso D, Lipschitz D, Houshyar S, et al.

 Brain-derived neurotrophic factor-5-HTTLPR gene interactions and environmental modifiers of depression in children. Biol Psychiatry 2006;59:673-80.
- Kim B, Kim CY, Hong JP, Kim SY, Lee C, Joo YH. Brain-derived neurotrophic factor Val/Met polymorphism and bipolar disorder. Association of the Met allele with suicidal behavior of bipolar patients. Neuropsychobiology 2008;58:97-103.
- Kim JM, Stewart R, Kim SW, Yang SJ, Shin IS, Kim YH, et al. Interactions between life stressors and susceptibility genes (5-HTTLPR and BDNF) on depression in Korean elders. Biol Psychiatry 2007;62:423-8.
- Knable MB, Barci BM, Webster MJ, Meador-Woodruff J, Torrey EF. Molecular abnormalities of the hippocampus in severe psychiatric illness: postmortem findings from the Stanley Neuropathology Consortium. Mol Psychiatry 2004;9:609-20.
- Kunugi H, Iijima Y, Tatsumi M, Yoshida M, Hashimoto R, Kato T, et al. No association between the Val66Met polymorphism of the brain-derived neurotrophic factor gene and bipolar disorder in a Japanese population: a multicenter study. Biol Psychiatry 2004;56:376-8.
- Lang UE, Hellweg R, Gallinat J. BDNF serum concentrations in healthy volunteers are associated with depression-related personality traits. Neuropsychopharmacology 2004;29:795-8.

- Lapalme M, Hodgins S, LaRoche C. Children of parents with bipolar disorder: a metaanalysis of risk for mental disorders. Can J Psychiatry 1997;42:623-31.
- Lin PY. State-dependent decrease in levels of brain-derived neurotrophic factor in bipolar disorder: a meta-analytic study. Neurosci Lett 2009;466:139-43.
- Müller DJ, de Luca V, Sicard T, King N, Strauss J, Kennedy JL. Brain-derived neurotrophic factor (BDNF) gene and rapid-cycling bipolar disorder: family-based association study. Br J Psychiatry 2006;189:317-23.
- March JS, Parker JD, Sullivan K, Stallings P, Conners CK. The Multidimensional Anxiety Scale for Children (MASC): factor structure, reliability, and validity. J Am Acad Child Adolesc Psychiatry 1997;36:554-65.
- McEwen BS. From molecules to mind. Stress, individual differences, and the social environment. Ann N Y Acad Sci 2001;935:42-9.
- McTeague LM, Lang PJ. The anxiety spectrum and the reflex physiology of defense: from circumscribed fear to broad distress. Depress Anxiety 2012;29:264-81.
- Montag C, Basten U, Stelzel C, Fiebach CJ, Reuter M. The BDNF Val66Met polymorphism and anxiety: support for animal knock-in studies from a genetic association study in humans. Psychiatry Res 2010;179:86-90.
- Montag C, Weber B, Fliessbach K, Elger C, Reuter M. The BDNF Val66Met polymorphism impacts parahippocampal and amygdala volume in healthy humans: incremental support for a genetic risk factor for depression. Psychol Med 2009;39:1831-9.
- Nakata K, Ujike H, Sakai A, Uchida N, Nomura A, Imamura T, et al. Association study of the brain-derived neurotrophic factor (BDNF) gene with bipolar disorder. Neurosci Lett 2003;337:17-20.
- Nemoto K, Ohnishi T, Mori T, Moriguchi Y, Hashimoto R, Asada T, et al. The Val66Met polymorphism of the brain-derived neurotrophic factor gene affects age-related brain

- morphology. Neurosci Lett 2006;397:25-9.
- Neves-Pereira M, Mundo E, Muglia P, King N, Macciardi F, Kennedy JL. The brain-derived neurotrophic factor gene confers susceptibility to bipolar disorder: evidence from a family-based association study. Am J Hum Genet 2002;71:651-5.
- Olson D. FACES IV and the Circumplex Model: validation study. J Marital Fam Ther 2011;37:64-80.
- Ostiguy CS, Ellenbogen MA, Linnen AM, Walker EF, Hammen C, Hodgins S. Chronic stress and stressful life events in the offspring of parents with bipolar disorder. J Affect Disord 2009;114:74-84.
- Ostiguy CS, Ellenbogen MA, Walker CD, Walker EF, Hodgins S. Sensitivity to stress among the offspring of parents with bipolar disorder: a study of daytime cortisol levels.

 Psychol Med 2011;41:2447-57.
- Oswald P, Del-Favero J, Massat I, Souery D, Claes S, Van Broeckhoven C, et al. Non-replication of the brain-derived neurotrophic factor (BDNF) association in bipolar affective disorder: a Belgian patient-control study. Am J Med Genet B Neuropsychiatr Genet 2004;129B:34-5.
- Poznanski EO, Freeman LN, Mokros HB. Children's Depression Rating scale-Revised (September 1984). Psychopharmacology Bulletin 1985;21:979-89.
- Rakofsky JJ, Ressler KJ, Dunlop BW. BDNF function as a potential mediator of bipolar disorder and post-traumatic stress disorder comorbidity. Mol Psychiatry 2012;17:22-35.
- Rasmusson AM, Shi L, Duman R. Downregulation of BDNF mRNA in the hippocampal dentate gyrus after re-exposure to cues previously associated with footshock.

 Neuropsychopharmacology 2002;27:133-42.
- Romero S, Delbello MP, Soutullo CA, Stanford K, Strakowski SM. Family environment in

- families with versus families without parental bipolar disorder: a preliminary comparison study. Bipolar Disord 2005;7:617-22.
- Roth TL, Lubin FD, Funk AJ, Sweatt JD. Lasting epigenetic influence of early-life adversity on the BDNF gene. Biol Psychiatry 2009;65:760-9.
- Russo-Neustadt A. Brain-derived neurotrophic factor, behavior, and new directions for the treatment of mental disorders. Semin Clin Neuropsychiatry 2003;8:109-18.
- Shimizu E, Hashimoto K, Koizumi H, Kobayashi K, Itoh K, Mitsumori M, et al. No association of the brain-derived neurotrophic factor (BDNF) gene polymorphisms with panic disorder. Prog Neuropsychopharmacol Biol Psychiatry 2005;29:708-12.
- Singh MK, DelBello MP, Stanford KE, Soutullo C, McDonough-Ryan P, McElroy SL, et al. Psychopathology in children of bipolar parents. J Affect Disord 2007;102:131-6.
- Skibinska M, Hauser J, Czerski PM, Leszczynska-Rodziewicz A, Kosmowska M, Kapelski P, et al. Association analysis of brain-derived neurotrophic factor (BDNF) gene Val66Met polymorphism in schizophrenia and bipolar affective disorder. World J Biol Psychiatry 2004;5:215-20.
- Sklar P, Gabriel SB, McInnis MG, Bennett P, Lim YM, Tsan G, et al. Family-based association study of 76 candidate genes in bipolar disorder: BDNF is a potential risk locus. Brain-derived neutrophic factor. Mol Psychiatry 2002;7:579-93.
- Soliman F, Glatt CE, Bath KG, Levita L, Jones RM, Pattwell SS, et al. A genetic variant BDNF polymorphism alters extinction learning in both mouse and human. Science 2010;327:863-6.
- Strauss J, Barr CL, George CJ, Devlin B, Vetró A, Kiss E, et al. Brain-derived neurotrophic factor variants are associated with childhood-onset mood disorder: confirmation in a Hungarian sample. Mol Psychiatry 2005;10:861-7.
- Suliman S, Hemmings SM, Seedat S. Brain-Derived Neurotrophic Factor (BDNF) protein

- levels in anxiety disorders: systematic review and meta-regression analysis. Front Integr Neurosci 2013;7:55.
- Surtees PG, Wainwright NW, Willis-Owen SA, Sandhu MS, Luben R, Day NE, et al. No association between the BDNF Val66Met polymorphism and mood status in a non-clinical community sample of 7389 older adults. J Psychiatr Res 2007;41:404-9.
- Tang J, Xiao L, Shu C, Wang G, Liu Z, Wang X, et al. Association of the brain-derived neurotrophic factor gene and bipolar disorder with early age of onset in mainland China. Neurosci Lett 2008;433:98-102.
- Thienemann M, Shaw RJ, Steiner H. Defense style and family environment. Child Psychiatry Hum Dev 1998;28:189-98.
- Tochigi M, Otowa T, Suga M, Rogers M, Minato T, Yamasue H, et al. No evidence for an association between the BDNF Val66Met polymorphism and schizophrenia or personality traits. Schizophr Res 2006;87:45-7.
- van Gastel W, Ferdinand RF. Screening capacity of the Multidimensional Anxiety Scale for Children (MASC) for DSM-IV anxiety disorders. Depress Anxiety 2008;25:1046-52.
- Vincze I, Perroud N, Buresi C, Baud P, Bellivier F, Etain B, et al. Association between brainderived neurotrophic factor gene and a severe form of bipolar disorder, but no interaction with the serotonin transporter gene. Bipolar Disord 2008;10:580-7.
- Wendland JR, Kruse MR, Cromer KR, Murphy DL. A large case-control study of common functional SLC6A4 and BDNF variants in obsessive-compulsive disorder.

 Neuropsychopharmacology 2007;32:2543-51.
- Young RC, Biggs JT, Ziegler VE, Meyer DA. A rating scale for mania: reliability, validity and sensitivity. Br J Psychiatry 1978;133:429-35.
- Zhang H, Ozbay F, Lappalainen J, Kranzler HR, van Dyck CH, Charney DS, et al. Brain derived neurotrophic factor (BDNF) gene variants and Alzheimer's disease, affective

disorders, posttraumatic stress disorder, schizophrenia, and substance dependence.

Am J Med Genet B Neuropsychiatr Genet 2006;141B:387-93.

Table 1 Demographic and clinical characteristics of the participants.

| | | BD offspring | Control | p |
|--------------------|---------------|--------------|--------------|-------|
| | | (N=64) | (N=51) | |
| Age | | 13.73 (3.45) | 13.68 (2.68) | .930 |
| Gender | male (%) | 30(46.88) | 23(45.10) | 1.000 |
| Race | Non-caucasian | 6(9.37) | 20(39.22) | <.000 |
| | (%) | | | |
| YMRS | mean (S.D.) | 6.50(6.90) | 1.10(1.34) | <.000 |
| CDRS | mean (S.D.) | 27.33(11.74) | 18.85(2.24) | <.000 |
| MASC | mean (S.D.) | | | |
| Physical | | 8.05 (7.01) | 4.31(4.06) | .001 |
| Harm avoidance | | 16.41(4.71) | 15.76(4.64) | .466 |
| Social anxiety | | 10.06 (6.14) | 7.75(5.41) | .037 |
| Separation anxiety | | 5.95(5.24) | 5.51(4.35) | .628 |
| MASC Total | | 40.47(17.09) | 33.33(14.24) | .018 |

MASC, Multidimensional Anxiety Scale for Children

YMRS, Young Mania Rating Scale

CDRS, Children's Depression Rating Scale

BD, Bipolar disorder

NOS, not otherwise specified

ADHD, attention-deficit/hyperactivity disorder

ODD, oppositional defiant disorder

GAD, generalized anxiety disorder

Table 2 The demographic and clinical characteristics by BDNF genotype in bipolar offspring.

| | | val/val | val/met | p |
|---------------|----------------|--------------|--------------|-------|
| | | (N=34) | +met/met | |
| | | | (N=30) | |
| Age | | 13.08 (3.34) | 14.51 (3.48) | .099 |
| Gender | Male (%) | 17 (48.6) | 13(44.8) | .481 |
| Race | Non-caucasian | 3 (8.82) | 3 (10.00) | .809 |
| | (%) | | | |
| YMRS | mean (S.D.) | 5.93(6.60) | 7.21(7.33) | .505 |
| CDRS | mean (S.D.) | 26.40(10.28) | 28.50(13.49) | .519 |
| Mood disorder | BD.NOS (%) | 9 (25.7) | 2 (6.9) | .046 |
| | MDD (%) | 10(28.6) | 9(31.0) | 1.000 |
| | Dysthymia (%) | 1(2.9) | 4(13.8) | .167 |
| Comorbidity | | | | |
| | ADHD (%) | 9 (25.7) | 6 (20.7) | .432 |
| | ODD (%) | 1(2.9) | 3(10.3) | .321 |
| | GAD (%) | 8 (22.9) | 5(17.2) | .406 |
| | Anxiety | 0 (0.0) | 3(10.3) | .088 |
| | disorder.NOS | | | |
| | Phobia | 2 (5.7) | 2(6.9) | 1.000 |
| Anti-anxiety | Total (%) | 8 (22.9) | 5 (17.2) | .757 |
| medication | | | | |
| | SSRI (%) | 5 (14.3) | 5 (17.2) | 1.000 |
| | Antidepressant | 5 (14.3) | 4 (13.8) | 1.000 |

| | other | than | | | | | |
|----------------|----------------|------|-------------|-------------|-------|--|--|
| | SSRI(%) | | | | | | |
| | Anxiolytics (% | %) | 2 (5.7) | 2 (6.9) | 1.000 | | |
| MASC | mean (S.D.) | | | | | | |
| Physical | | | 8.74(6.7) | 7.21(7.4) | .388 | | |
| Harm avoidance | | | 17.89(3.9) | 14.62(5.0) | .005 | | |
| Social | | | 11.69(5.7) | 8.10(6.2) | .019 | | |
| anxiety | | | | | | | |
| Separation | | | 7.09(5.6) | 4.59 (4.4) | .057 | | |
| anxiety | | | | | | | |
| MASC Total | | | 45.40(14.7) | 34.51(18.1) | .010 | | |
| | | | | | | | |

YMRS, Young Mania Rating Scale

CDRS, Children's Depression Rating Scale

BD, Bipolar disorder

NOS, not otherwise specified

MDD, major depressive disorder

ADHD, attention-deficit/hyperactivity disorder

ODD, oppositional defiant disorder

GAD, generalized anxiety disorder

SSRI, selective serotonin reuptake inhibitor


MASC, Multidimensional Anxiety Scale for Children

Figure Captions


Fig. 1. Anxiety symptoms differences (a) between the bipolar offspring group and healthy control group (b) by BDNF genotype in bipolar offspring.

Fig. 2. The moderating effect of BD offspring status on anxiety symptoms.

Fig. 1. Anxiety symptoms differences (a) between the bipolar offspring group and healthy control group (b) by BDNF genotype in bipolar offspring.




(a)




(b)


Fig. 2. The moderating effect of BD offspring status on anxiety symptoms.



(a) MASC total



(b) Harm avoidance



(c) Social anxiety