

# COGNITIVE IMPAIRMENT IN BIPOLAR DISORDER: MAGNITUDE, MODERATORS AND PREVENTION STRATEGIES

Journal:	<i>The International Journal of Neuropsychopharmacology</i>
Manuscript ID	IntJNP-16-0191
Manuscript Type:	Review
Date Submitted by the Author:	15-Dec-2016
Complete List of Authors:	<p>Solé, Brisa; Bipolar Disorders Program. Clinical Institute of Neuroscience, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM</p> <p>Jiménez, Esther; Bipolar Disorder Program, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM</p> <p>Torrent, Carla; Clinical Institute of Neurosciences, Hospital Clínic Barcelona, Bipolar Disorders Program</p> <p>Bonnín, Caterina; Clinical Institute of Neurosciences, Hospital Clínic Barcelona, Bipolar Disorders Program</p> <p>Torres, Imma; Bipolar Disorders Program. Clinical Institute of Neuroscience, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM</p> <p>Reinares, Maria; Bipolar Disorders Program, Clinical Institute of Neuroscience, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM</p> <p>Varo, Cristina; Bipolar Disorders Program. Clinical Institute of Neuroscience, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM</p> <p>Grande, Iria; Bipolar Disorders Program, Clinical Institute of Neuroscience, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM</p> <p>Valls, Elia; Bipolar Disorders Program. Clinical Institute of Neuroscience, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM</p> <p>Salagre, Estela; Bipolar Disorders Program. Clinical Institute of Neuroscience, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM</p> <p>Sanchez-Moreno, Jose; Bipolar Disorders Program. Clinical Institute of Neuroscience, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM</p> <p>Martínez-Aran, Anabel; Clinical Institute of Neuroscience. Hospital Clinic of Barcelona, Psychiatry</p> <p>carvalho, af; Federal University of Ceara,</p> <p>Vieta, Eduard; Bipolar Disorders Program. Clinical Institute of Neuroscience, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM,</p> <p>;</p>
Research Focus: Choose one or more (maximum 5) that best describe the research focus of your paper:	Neuropsychology < Clinical
Keywords: Enter up to 5 (minimum 3) keywords that best reflect the research reported in your paper.:	bipolar disorder, cognitive impairment, cognitive remediation, functional remediation

SCHOLARONE™  
Manuscripts

## COGNITIVE IMPAIRMENT IN BIPOLAR DISORDER: MAGNITUDE, MODERATORS AND PREVENTION STRATEGIES

**Category:** review

**Authors:** Brisa Sole<sup>1</sup> PhD, Esther Jimenez<sup>1</sup> PhD, Carla Torrent<sup>1</sup> PhD, Caterina del Mar Bonnin<sup>1</sup> PhD, Imma Torres<sup>1</sup> PsyD, Maria Reinares<sup>1</sup> PhD, Cristina Varo<sup>1</sup> PsyD, Iria Grande<sup>1</sup> MD PhD, Elia Valls<sup>1</sup> PsyD, Estela Salagre<sup>1</sup> MD, Jose Sanchez-Moreno<sup>1</sup> PsyD, Anabel Martinez-Aran A PhD, André F Carvalho<sup>2</sup> MD PhD, Eduard Vieta<sup>1</sup> MD PhD

**Affiliations:** <sup>1</sup>Barcelona Bipolar Disorders Program, Institute of Neurosciences, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM, Barcelona, Catalonia, Spain.

<sup>2</sup> Translational Psychiatry Research Group, Faculty of Medicine, Federal University of Ceará, Fortaleza, CE, Brazil

Correspondence: Eduard Vieta, Hospital Clinic, 170 Villarroel st, 08036 Barcelona, Spain. Tel +34932275477; Fax +34932279228; email: [evieta@clinic.ub.es](mailto:evieta@clinic.ub.es)

**Word count abstract:** 145

**Word count manuscript:** 6392

**Number of references:** 118

**Number of figures:** 1

**Number of tables:** 2

**Abstract:**

Over the last decade there has been a growing appreciation of the importance of identifying and treating cognitive impairment associated with bipolar disorder (BD) since it persists in remission periods, and evidence indicates that neurocognitive dysfunction may significantly influence patients' psychosocial outcomes. An ever increasing body of research seeks to achieve a better understanding of potential moderators contributing to cognitive impairment in BD. This report provides an overview of data from studies examining the magnitude of cognitive dysfunction in bipolar disorder and those potential clinical and demographic factors influencing cognition. Then, we also review currently available treatments for cognitive dysfunction in BD as well as potential novel treatments. All these data encourage additional efforts and further studies to find effective strategies to prevent and treat cognitive impairment associated with bipolar disorder. These efforts may ultimately lead to an improvement of psychosocial functioning in these patients.

**Key words:** cognitive impairment, bipolar disorder, cognitive remediation, functional remediation

## **INTRODUCTION**

The study of neurocognitive impairment, its causes and consequences, as well as the development of new therapeutic strategies to manage, or even to prevent this kind of deficits, is nowadays one of the hottest areas of research in bipolar disorder (BD) (Martinez-Aran and Vieta, 2015). Data from different meta-analyses confirm that most patients with bipolar disorder present with neurocognitive dysfunction, even during euthymia (Robinson et al., 2006; Bourne et al., 2013; Bortolato et al., 2015). Some of these neurocognitive deficits seem to be present not only in the early course of the illness (Torres et al., 2010; Lee et al., 2014; Bora and Pantelis, 2015), but also in premorbid stages before illness onset (Martino et al., 2015). According to the most recent meta-analyses, the most affected domains, with effect sizes ranging from moderate to high, are attention, verbal learning and memory and executive functions, whereas premorbid intelligence appears to be preserved (Kurtz and Gerraty, 2009; (Bourne, et al., 2013). Nevertheless, it is worth to mention that the effect sizes have become smaller since the first meta-analysis was published. Although cognitive abnormalities are present across all illness phases, they are usually more notable during acute episodes (Kurtz and Gerraty, 2009). For instance, patients could present a poorer verbal learning performance during manic episodes, and bipolar depressed patients may exhibit lower performance in verbal fluency tasks.

Due to the fact that BD has a high heritability, it was not surprising that unaffected first-degree relatives and offspring of patients with BD present mild verbal memory and executive dysfunction. In this sense, some authors have suggested that neurocognitive deficits could be considered as putative endophenotypes of BD (Arts et al., 2008; Balanzá-Martínez et al., 2008; Bora et al., 2009). Nonetheless, it is also remarkable that neurocognitive deficits are not specific of BD and they may be considered as a common dimension across disparate psychiatric disorders, a trans-nosological domain (Millan et al., 2015; Vieta, 2016). In this vein,

the Research Domain Criteria (RDoc) is an alternative approach with the purpose to conduct research in terms of dimensions, defined by neurobiology and behavioral measures, which cut across traditional diagnostic categories. This framework incorporates genetics, neuroimaging and cognitive sciences for a new classification of mental disorders (Cuthbert, 2014).

In the last years, there is also growing evidence for impairment in some social cognition domains (this is a multi-dimensional concept) even during periods of remission (Samamé et al., 2012, 2015). In general, evidence supports a theory of mind deficit in euthymic bipolar patients, whereas it remains unclear whether substantial deficits in other social cognition dimensions could persist in euthymic patients with BD (Bora and Pantelis, 2016). Importantly, two points need to be kept in mind regarding social cognition: first, there is a large number of available tasks that evaluate social cognition domains with different level of complexity and quality; second, some findings point out that other neurocognitive deficits may influence social cognitive performance and this issue deserves further exploration (Samamé et al., 2012).

Neurocognitive impairment needs to be considered a therapeutic clinical target in order to improve both psychosocial functioning and quality of life of bipolar patients. Available evidence underscores that cognitive dysfunction is a critical mediator of adverse psychosocial outcomes in BD and a predictor of unfavorable employment outcomes (Tse et al., 2014; Baune and Malhi, 2015;). It is worthy to note that similarly to cognitive dysfunction, functional deficits persist even after symptomatic remission in a significant subset of patients with BD, thus adding additional burden to patients and also possibly increasing illness-related direct and indirect costs. In this sense, BD is ranked among the leading causes of burden of disease worldwide associated with a high level of disability-adjusted life years (Catalá-López, et al., 2013). Fortunately, in the last years some efforts have been done to develop interventions addressing neurocognitive function in BD. Nevertheless, this area of research is still on its early developmental stages.

The overarching aim of this review to draw a comprehensive picture of the complexity of cognitive dysfunction in BD, focusing on: 1) describing the magnitude of cognitive dysfunction in BD, 2) identifying potential moderators - factors that may influence cognition- that underlie cognitive impairment and 3) describing treatment approaches that primary address cognitive dysfunction in BD.

### **MAGNITUDE OF COGNITIVE IMPAIRMENT**

Evidence points out that the neurocognitive impairment profile observed in patients with BD is similar to that showed in patients with schizophrenia although in a lesser extent, therefore, differences between the two disorders seem to be predominantly quantitative rather than qualitative (Daban et al., 2006). Patients with schizophrenia also significantly underperform bipolar patients in social cognitive tasks, such as emotion recognition and theory of mind similarly to findings for other neurocognitive tasks (Bora and Pantelis, 2016). Nevertheless, an important matter is that studies comparing both psychiatric disorders have not taken into consideration the potential effect of the extant cognitive heterogeneity in both disorders.

Overall, approximately 40-60% of patients with BD exhibit neurocognitive impairment, with a large heterogeneity among them, as it will be discussed later. There has been disparity in the establishment of a threshold as well as criteria for the cognitive impairment in BD, with some studies being more conservative than others. Hence, percentages of cognitive impairment in the existing studies vary according to the neuropsychological tool or the cut-off used to consider impairment (i.e. 1 or 2 standard deviations below the mean). Beyond the percentage of neurocognitively impaired bipolar patients, converging data from a few recent studies suggest that there are several neurocognitive subtypes among bipolar patients. This neurocognitive heterogeneity may explain, at least in part, the extant variability on psychosocial functioning among patients. The use of cluster analysis approaches has enabled different authors to detect distinct neurocognitive profiles among both bipolar I and bipolar II

patients: one with a normal performance, one (or two groups) with selective modest impairments and, lastly, another showing a more globally severe cognitive impairment (i.e., encompassing several domains) (Burdick et al., 2014; Bora et al., 2016; Jensen et al., 2016; Solé et al., 2016). This qualitative and quantitative heterogeneity of neurocognitive impairment has also been observed in other psychiatric conditions such as schizophrenia (Lewandowski et al., 2014). In those studies assessing cross-diagnostic samples, patients with BD represent a larger proportion in the neuropsychologically normal group in comparison to schizophrenic patients which are more prone to be severely impaired (Lewandowski et al., 2014; Bora et al., 2016).

Differences in clinical or sociodemographic variables (e.g., schooling) might at least in part explain the variability of neurocognitive performances across studies, as will be reviewed later in this article. Across the different studies using clustering methods, patients belonging to the cluster with severe neurocognitive deficits had a significantly poorer educational attainment and/or a lower premorbid intelligence quotient (IQ) (Lewandowski et al., 2014; Bora et al., 2016; Solé et al., 2016). Burdick and colleagues also observed that their selectively impaired group had a higher number of total affective episodes supporting the negative impact of affective recurrences on neurocognitive functioning (Burdick et al., 2014). Moreover, these studies also showed that the more neurocognitive impairment, the poorer the functional outcome (Burdick et al., 2014; Jensen et al., 2016; Solé et al., 2016).

As some authors suggest, the neurocognitive variability might reflect an etiological heterogeneity in BD including potential different subtypes associated with different genetic susceptibility factors (Bora, 2016). Evidence shows that BD shares some susceptibility genes with schizophrenia whereas some other genetic susceptibility factors seem to be specific of each disorder (Lichtenstein et al., 2009; Craddock et al., 2010). Taking all these findings into consideration, it has been hypothesized the existence of two groups: a group of bipolar patients characterized by normal neurodevelopmental and cognitive functioning, whose



cognitive decline is probably influenced by the impact of repetitive affective episodes; and another much smaller group of patients presenting with a cognitive impairment comparable to that observed in schizophrenia, characterized by a low premorbid cognitive functioning before illness onset. This latter group of patients would share common genetic risk factors with schizophrenia and might be associated with neurodevelopmental abnormalities. At this point, further genetic and neurobiological research is needed to confirm this hypothesis.

Note that the neurocognitive variability should be considered when we interpret data coming from meta-analytic and systematic-review findings. As has been mentioned before, several clinical factors might influence this heterogeneity. Nevertheless, we cannot dismiss methodological issues as well as other intrinsic individual factors (e.g. motivation, self-esteem, etc.) as potential factors explaining, at least partially, this variability.

#### **MODERATORS OF NEUROCOGNITIVE FUNCTIONING**

The study of factors involved in neurocognitive impairment is one of the first steps to prevent or mitigate the neurocognitive dysfunction associated with BD. Nevertheless, it is not well established yet whether neurocognitive deficits are due to neurodevelopmental abnormalities, to the illness progression or if they reflect part of both processes (Goodwin et al., 2008). Altogether, neurocognitive dysfunction in BD seems to be multifactorial where a series of clinical factors has been suggested to exert some effect, direct or indirectly, on neurocognitive functioning (see Table 1.).

***Number of episodes and chronicity:*** In this regard, many of the cross-sectional studies point towards an association between number of affective episodes and neurocognitive impairment suggesting a progressive cognitive decline. When patients with multiple manic episodes were compared with patients with a first manic episode, those patients with a recurrence of manic episodes showed more deficits in executive functions and episodic memory and reduced psychomotor speed (López-Jaramillo et al., 2010; Hellvin et al., 2012). These findings also

support those studies on neuroanatomical differences in bipolar I patients. Nonetheless, we cannot fail to mention other several points. First, recurrence of depressive episodes has also been suggested to exert a detrimental effect on cognition in some study, though this association has been less replicated (Summers et al., 2006). Secondly, contrary to most cross-sectional studies, evidence from longitudinal studies is not in accordance with a neurocognitive progression in BD. As a whole, the follow-up studies have found a stable pattern of neurocognitive impairment over time (Budde and Schulze, 2014; Samamé et al., 2014). Nevertheless, the longer follow-up study carried so far revealed that most neurocognitive functions remained stable over time with the exception of a worsening in the executive functions, which was associated with illness duration and subdepressive symptoms (Torrent et al., 2012). Therefore, short time frames of assessment in most of the longitudinal studies could explain the lack of a cognitive function decline. Thirdly, data from a meta-analytic study indicate that neurocognitive impairment in first-episode bipolar patients was equivalent to that observed on patients with a chronic course of the illness (Bora and Pantelis, 2015). Explanations for this apparent contradiction may involve the fact that the patients who were able to stay for long-term follow-up might not be representative of the majority of bipolar patients; they may be more adherent and may have many of the “good prognosis” indicators that protect against cognitive decline.

Another interesting study showed that cognitive impairment was present in remitted bipolar patients from the first manic episode. However, those episode-free patients improved cognitively in the year following the first manic episode, so impairment seems reversible when compared to patients who experienced recurrences of affective episodes (Kozicky et al., 2014). Hence, these results would suggest the need to implement interventions in the early stages to avoid affective recurrences and to reverse neurocognitive deficits too.

Therefore, some inconsistent findings between the extant cross-sectional and longitudinal studies highlight the necessity to further research to elucidate the veritable evolution of cognitive dysfunction in bipolar illness and potential selection bias in longitudinal studies. During the last decade different staging model approaches have been proposed for BD (Berk et al., 2007; Kapczinski et al., 2008;). Both models assume an underlying pathophysiological process of neuroprogression associated with cognitive decline among other neurobiochemical changes, however, not every patient will proceed through all stages. Therefore, early identification of which patients will develop a neuroprogressive disorder as well as linking neurocognitive impairment with staging models is one of the challenges in the upcoming years (Rosa et al., 2014; Passos et al., 2016).

***Psychotic symptoms:*** Several studies reported an association between lifetime history of psychotic symptoms and neurocognitive impairment, mainly with a deleterious effect on executive functions and verbal memory (Glahn et al., 2007; Martinez-Aran et al., 2008). Besides, some authors suggested that neurocognitive dysfunction depends on history of psychosis rather than the diagnostic category *per se*, supporting a dimensional approach in the severe psychiatric illnesses classification systems (Simonsen et al., 2009). Nevertheless, we can also found some studies that did not found any relationship between cognitive impairment and psychotic symptomatology (Selva et al., 2007). These inconsistent findings are probably due to the use of small sample which include symptomatic patients. Overall, according to meta-analytic findings, the effect of the history of psychosis in BD is modest, not supporting a complete categorical distinction between patients with or without history of psychotic symptoms (Bora et al., 2010).

***Subclinical depressive symptoms:*** Another important factor that may exert an impact on psychosocial outcome and also on cognition is the subclinical depressive symptomatology, even at low levels (Bonnín et al., 2011; Torrent et al., 2012). A recent study using a path

analysis found that verbal memory had a significant indirect effect on functional outcome, partly mediating the relationship between depressive symptoms and functional status (Bonnín et al., 2014). Verbal memory has been linked to functional outcome in several studies and previous literature also links subclinical symptomatology with verbal memory (Martinez-Aran et al., 2007; Martino et al., 2009; Bonnín et al., 2010). Thus, depressive symptoms are not only negatively associated with functional outcome, but also affect verbal memory performance.

**Diagnostic subtype of bipolar disorder:** Due to the fact that the two main diagnostic subtypes of BD (bipolar I and II) present with a different clinical course and that manic and depressive episodes might exert a differential influence over cognitive function, a reduced number of studies have compared neurocognitive performance between both diagnostic subtypes. Despite some constraints affecting findings in this area of research, the vast majority of studies comparing euthymic samples detected a worse neurocognitive performance in patients with bipolar I subtype compared to bipolar II subtype, especially in verbal and visual memory domains and some tasks related with executive functions (e.g. verbal fluency) (Bora, 2011; Solé et al., 2011). Nevertheless, differences between the two subtypes seem to be small. This little but worse performance in bipolar I patients might be more consistently related with manic episodes which seem to exert a negative long-term effect on neuropsychological performance.

**Pharmacological treatment:** Medication constitutes an important confounding factor which is not easy to control for due to the fact that most patients with BD are usually treated with combinations of different drugs and few patients are medication-free. In addition, medication may exert conflicting effects on cognition: although some pharmacological treatments may have an indirectly protective role reducing affective and psychotic symptomatology, several drugs may not be free of neuropsychological negative side effects, especially in complex combinations. Besides, there is some controversy regarding neurotoxic and neuroprotective effects of several agents. Concerning lithium, the most widely studied mood stabilizer with

regards to its neurocognitive effects, seems to have a negative, although minor, effect on verbal memory and psychomotor speed (Wingo et al., 2009). Nevertheless, some preclinical studies point out the neuroprotective and neurotrophic effects of lithium. In fact, an interesting study showed that a small sample of excellent lithium responders exhibited normal cognitive functioning and plasma BDNF (brain derived neurotrophic factor) levels compared to the remaining lithium patients where the effect of lithium was not optimal (Rybakowski and Suwalska, 2010). With regards to antipsychotic and antiepileptic treatments, few studies have been specifically conducted with bipolar patients, with most data coming from schizophrenia or patients with epilepsy. Generally, antipsychotics have been associated with executive dysfunction and lower psychomotor speed (Frangou et al., 2005; Bora et al., 2009). After reviewing the available literature concerning BD treatment, Solé & Jimenez conclude that any atypical antipsychotic appears as better than others with regard to its cognitive profile (Solé and Jiménez, 2015). Concerning anticonvulsants in bipolar patients, they seem to exert similar cognitive effects than those described in volunteers and patients with epilepsy (Gualtieri and Johnson, 2006). Hence, lamotrigine and oxcarbazepine have more favorable cognitive profile, whereas carbamazepine, valproic acid and topiramate have the most cognitive toxicity. Benzodiazepines, another commonly prescribed drug type among bipolar patients, can interfere with memory. Lastly, most of the studies examining the effects of antidepressants on cognitive functioning come from patients suffering a major depressive disorder (Popovic et al., 2015), and a paucity of them were conducted with non-geriatric adult samples. As a whole, antidepressants seem to have beneficial effects in reducing cognitive impairment, especially with a positive effect on psychomotor speed and delayed recall, with no significant differences between different antidepressant classes (selective serotonin reuptake inhibitors –SSRIs-, serotonin and norepinephrine reuptake inhibitors –SNRIs-, dopamine modulators and norepinephrine inhibitors) (Keefe et al., 2014; Rosenblatt et al., 2016).

Despite potential side effects of pharmacological treatments, studies conducted with medication-free bipolar patients, or comparing patients with or without pharmacological treatment, indicate that cognitive impairment is caused by the illness impact and little effects are due to the medication *per se* (Goswami et al., 2009).

**Comorbidity:** BD is often accompanied by multiple medical comorbidities. Many of these conditions could constitute additional factors that influence neurocognitive performance in bipolar patients (e.g. substances use disorder, anxiety, attention deficit hyperactivity disorder –ADHD-, etc.) (Balanzá-Martínez et al., 2010).

Wu et al. found out that bipolar II patients with a comorbid anxiety disorder showed poorer cognitive performance than those without, especially in verbal and non-verbal memory, psychomotor speed and working memory (Wu et al., 2011). Volkert and colleagues also found that comorbid anxiety was significantly more prevalent among patients with cognitive impairment (Volkert et al., 2015). Furthermore, Levy and colleagues also found more symptoms of acute anxiety (a higher arousal) during neuropsychological tasks in those euthymic bipolar patients showing a poorer performance than in the healthy control group (Levy, 2013). This group of authors suggested that symptoms of acute anxiety triggered by cognitive challenge may compromise cognitive functioning in patients with BD.

Substance abuse is also particularly common. Bipolar patients with comorbid alcohol abuse exhibited more executive dysfunction and verbal and visual memory deficits than those without comorbidity (van Gorp et al., 1998; Levy et al., 2008). Even bipolar patients with previous history of alcohol consumption exhibit a poorer performance in inhibitory control in comparison with healthy controls (Sanchez-Moreno et al., 2009), however, no other differences were detected when compared to patients without history of consumption. Maybe, the potential negative effects of alcohol could be reversible after a long timeframe of abstinence. Nonetheless, comorbid patients had a less favorable period of recovery of

cognitive deficits over the course of mood remission after suffering an affective episode than those bipolar patients without comorbidity (Levy et al., 2013).

Alcohol abusers also showed higher impulsivity in all the components of the Barratt Impulsiveness Scale (i.e. non-planning, attentional and motor) when compared with non-comorbid bipolar patients (Nery et al., 2013). In this line, another study showed that performance in the Iowa Gambling Test (IGT) is a good predictor of risk for future drug use in bipolar patients with stimulant dependence (Nejtek et al., 2013).

Concerning ADHD, the first study including a comorbid BD+ADHD group showed that this group of patients performed significantly worse than both a pure ADHD and a healthy control group (Silva et al., 2014). Therefore, according to these results, authors suggested that some dysexecutive function, such as set-shifting deficits, in comorbid ADHD+BD patients may to a large extent be attributable to comorbid BD. Nevertheless this study was only focused on executive functions and was also lacking of a pure bipolar patients group. Recently, Torres and colleagues compared four groups (euthymic bipolar patients, ADHD patients, comorbid BD+ADHD patients and a healthy control group) for the first time (I. Torres, unpublished observations). This study detected that the comorbid group had a similar neurocognitive profile than pure bipolar patients, most likely reflecting the same neurobiological basis. The group of ADHD patients exhibited more selective cognitive impairment limited to memory and attention domains, although those ADHD patients without pharmacological treatment had a similar cognitive profile to bipolar patients. Therefore, these results also suggest that cognitive profile in adult comorbid ADHD+BD seems to be more related to BD than to ADHD.

Overweight and obesity are highly prevalent among bipolar patients and several studies report a negative impact on the course of the illness, with an increasing risk of developing diseases associated with obesity (e.g. diabetes mellitus, hypertension, etc.) (Liu et al., 2013). Overweight produces several pathophysiological changes (cardiovascular diseases, insulin

regulation, inflammation, etc.) affecting brain regions that may impact on neurocognitive function. Some studies showed that those euthymic bipolar patients with overweight/obesity have a poorer neurocognitive execution than those without, especially in verbal learning and memory domain (Yim et al., 2012; Depp et al., 2014; Lackner et al., 2015). Probably both conditions (BD and overweight) contribute to neurocognitive impairment in independent ways, however, the combination of both entails higher neurocognitive deficits.

Recognizing comorbidities is an important issue since some of them may complicate not only the course and treatment of patients but also may contribute to the magnitude of cognitive dysfunction adding other potential pathophysiological routes. But more important is the fact that treating some of these conditions may ameliorate cognitive impairment since some of them may be modifiable.

**Childhood adversity:** Adverse life events, as childhood abuse or neglect, are frequent in bipolar patients and have been related to worsening the clinical course of the illness (Daruy-Filho et al., 2011). Besides that, several studies indicate that traumatic events during childhood are associated with structural and functional brain alterations (Quidé et al., 2016). Savitz and colleagues found an association between sexual abuse, emotional abuse and neglect with poorer performance on measures of verbal/visual memory and executive functions (Savitz et al., 2008). Marshall and co-workers also found a significant main effect of trauma in inhibitory control accuracy not only in bipolar patients but also in healthy individuals (Marshall et al., 2016). In the same vein, Russo found that stable bipolar patients with a childhood history of emotional neglect performed worse in recognizing anger in tasks of facial emotion recognition than those without such a history (Russo et al., 2015). These authors speculated that the fact of suffering childhood maltreatment might lead to an increased vulnerability to develop more severe emotional dysregulation through abnormal neurodevelopment and consequently impaired emotional processing. Bückner and colleagues also detected an early influence of



childhood trauma on cognition early in the course of the illness, after the first manic episode, showing a more severe cognitive impairment in those patients with history of childhood trauma (Bücker et al., 2013). Furthermore, the same group of authors identified a significant reduction in total and regional volumes of the corpus callosum in a sample of bipolar patients recently recovered of a first manic episode with trauma compared with those without trauma (Bücker et al., 2014). Interestingly, another group of authors found that, although level of history of childhood trauma was higher in a group of schizophrenic patients, childhood trauma was more associated with poorer cognitive performance in the sample of first-episode affective psychoses (bipolar and psychotic depression) than in the non-affective psychoses (Aas et al., 2011). Lastly, a recent study examined the potential impact of childhood trauma specifically on distinct neurocognitive profiles using a data-driven analysis (E. Jimenez, personal communication). Higher levels of childhood trauma assessed with the Childhood Trauma Questionnaire (CTQ) were detected in those patients clustered in the low performance group. Moreover, the total CTQ score and the estimated IQ significantly contributed to differentiate among the different neurocognitive subgroups. Interestingly, this study found that physical negligence, which has been traditionally underreported, may also impact cognitive performance. Therefore, it should be necessary to assess and target mild forms of maltreatment too.

Despite the significant impact that may exert early life stress and maltreatment in bipolar patients, and particularly in cognitive functioning, there are few studies investigating its effects and it constitutes a crucial issue that awaits future research with longitudinal studies.

**Cognitive reserve:** Cognitive reserve reflects the capacity of the adult brain to endure neuropathology, minimizing clinical manifestations and allowing a successful accomplishment of cognitive tasks (Stern, 2009). It is a concept commonly used in neurology which has also been applied in neuropsychiatric disorders during the last years. Genetic and

neurodevelopmental factors determine cognitive reserve, however, exposure to specific environmental factors as education, lifestyle, mental and physical activities may also influence it. There are different proxy indicators of cognitive reserve, being the most commonly used years of educational attainment, leisure activities and premorbid IQ.

Very recently some studies have suggested that cognitive reserve may be a significant predictor of both cognitive and psychosocial outcome in euthymic bipolar patients which indicates that individual differences of brain characteristics and usage before illness onset may influence the future functional and neuropsychological outcome (Anaya et al., 2015; Forcada et al., 2015). Interestingly, a higher cognitive reserve has also been demonstrated to be associated with better neurocognitive, functional and clinical outcomes in first psychotic episodes and to be predictive of functioning at two-year follow-up of this group of patients (Amoretti et al., 2016). Altogether, these data suggested that early interventions to enrich cognitive reserve might result in minimizing the detrimental neuropsychological and functional impact caused by the illness (Vieta, 2015).

Cognitive reserve is still a new concept in the BD field, therefore, further research is guaranteed in the next years to have a better understanding of individual differences.

Other potential confounders that might be also influencing cognition have been less studied as rapid cycling (Wu et al., 2016), affective temperaments (Russo et al., 2014) or history of obstetric complications (Martino et al., 2008) and all of them deserve further research.

### **PREVENTION STRATEGIES**

**Pharmacotherapy:** The fact of considering cognitive dysfunction as a core feature of BD has led to a growing interest in developing both pharmacological and non-pharmacological prevention strategies to treat this type of deficits. Unfortunately, there is no well-established pharmacological treatment for neurocognitive impairment in BD since studies have yielded

mixed results with no convincing effects. From a pharmacologically therapeutic perspective, different drugs with potential beneficial effects for the treatment of neurocognitive impairment have been tested (e.g. some cholinesterase inhibitors, glutamate receptor antagonists as memantine, glucocorticoid receptor antagonists as mifepristone, dopaminergic agonists as pramipexole, intranasal insulin, some antioxidants, erythropoietin, etc.) (see Sanches et al., 2015 and Miskowiak et al., 2016). Despite the efforts done so far, no drug has been approved as a pro-cognitive enhancer for BD, although some of these drugs appear as promising candidates. Therefore, further research is required to find compounds which may become considered as pro-cognitive enhancers. Meanwhile, clinicians should bear in mind a rational use of drugs as well as the cognitive profile of each compound in order to minimize the cognitive side-effects.

***Non-pharmacological approaches:*** Prior research on *cognitive remediation*, even on other non-pharmacological interventions, in schizophrenia has provided some guides for BD. Nevertheless, as abovementioned, neurocognitive dysfunction in schizophrenia is of greater magnitude than observed in patients with BD, therefore, it is necessary to adjust or to develop new interventions specifically addressed to the characteristics of the latter group. Some authors have suggested some recommendations for developing specific cognitive remediation programs for bipolar illness (see Fuentes-Durá et al., 2012).

Very few studies have focused only on bipolar patients, and most of them were conducted with mixed affective disorder samples and were not rigorous controlled. The first study, focused specifically on bipolar patients, was a small open trial for subjects with residual depressive symptoms (Deckersbach et al., 2010). This study detected a reduction of depressive symptoms and an increase in psychosocial functioning in patients after receiving 14 individual sessions of cognitive remediation.

*Functional remediation* is an innovative intervention aimed at restoring psychosocial functioning specifically designed for bipolar patients. In 21 weekly sessions functional remediation provides several neurocognitive strategies and techniques for daily life to tackle the main neurocognitive deficits associated with BD (e.g. attention, memory, and executive functions). The first block of this intervention consists of a group of psychoeducative sessions about neurocognition and the impact of neurocognitive impairment in their daily lives, trying to increase, in this way, the insight about neurocognitive dysfunction. Family is involved in this process in order to increase the awareness of neurocognitive impairment associated with bipolar illness and to encourage the practice of all the learned strategies in each session and the reinforcement. The following three blocks are devoted to manage the three main affected cognitive areas related to the disorder, aforementioned. Then, the last block addresses some communication skills and techniques to the stress management in order to promote the autonomy of patients. The intervention includes both individual and group format tasks in an ecologic setting in order to establish a connection between the learned skills and strategies with daily life situations of patients (as work, autonomy, etc.).

The efficacy of functional remediation was proven in a randomized control trial comparing functional remediation to psychoeducation and to treatment as usual (Torrent et al., 2013). Patients receiving the functional remediation program improved the overall psychosocial outcome and, specifically, the interpersonal and occupational functioning. The intervention was also effective in bipolar II patients (Solé et al., 2015) and in secondary analysis this intervention proved to improve neurocognitive outcomes in cognitively impaired patients (Bonnin et al., 2015). Even more importantly, the functional improvement in the functional remediation group persisted at 1-year follow-up compared with the other two treatment groups (Bonnin et al., 2016). Autonomy was the functional domain that improved at 1-year follow-up and, interestingly, verbal memory also significantly improved from baseline to 1-year

follow-up. This cognitive improvement was not detected at the end of the intervention, at 6-month follow-up.

More recently, a small open pilot study assessing the feasibility of a briefer functional remediation program, which combined individual and group sessions, also showed a positive effect on overall psychosocial functioning in a sample of bipolar I patients (Zyto et al., 2016). This type of format seems to allow a more tailored approach to facilitate achieving personal goals.

In contrast, in another randomized control trial conducted by Demant and colleagues, cognitive remediation was not effective to remediate cognitive dysfunction in partial remitted bipolar patients with a 12 weeks group-based program (Demant et al., 2015). As the authors suggested, a more intensive and durable intervention may be necessary to improve cognition in bipolar patients.

Due to fast advance in information and communication technology (ICT) issues, one challenge in cognitive remediation is to implement computerized neurocognitive treatments in an effective manner. This kind of intervention delivery makes easier the accessibility to patients engaged in working life as well as to younger users who are familiar with. In fact, the aforementioned study by Demant and colleagues introduced a computer-assisted cognitive training as a part of the cognitive remediation program. Concerning this topic, Lewandowski and colleagues have published a study protocol for assessing the efficacy of an internet-based cognitive remediation program for patients with BD type I (Lewandowski et al., 2016). Unfortunately, results are not available yet.

Due to the fact that social cognition also seems to be involved in psychosocial functioning, an interest in developing strategies to enhance social cognition has emerged over the last years (Lahera et al., 2012). Nevertheless, any study has been exclusively focused on bipolar patients. Lahera and co-workers assessed the efficacy of the Social Cognition and Interaction Training

(SCIT), an intervention originally developed for patients with schizophrenia, in a sample mostly composed by bipolar patients but also some schizoaffective patients (Lahera et al., 2013). The intervention, addressed to improve emotion perception, attributional style and theory of mind, was found to be effective improving these social cognition domains but not social functioning. Maybe, interventions aimed to train social cognition would need to be adapted to the specific profile of social cognitive deficits observed in BD. In any case, an extension of social cognition training in cognitive/functional remediation interventions would be an interesting option of research for bipolar patients.

Therefore, interventions focused on the enhancement of cognition and psychosocial functioning in bipolar patients are still in its early stages and further research is needed in order to find the key components of cognitive and/or functional remediation.

Several positive effects have been associated with *physical exercise* such as increased production of brain neurotrophic factors and increased activity of specific neurotransmitters. The results obtained across other psychiatric or neurological conditions, as well as on aging, suggest that aerobic physical exercise may also have unequivocal beneficial effects on cognitive functioning in bipolar disorders, though no studies are available in this population (Kucyi et al., 2010; Malchow et al., 2013). Most of the studies about the effects of exercise in affective disorders, which are generally conducted with patients with major depressive disorder, are focused on mood, anxiety and quality of life. A reduction in these affective symptoms might lead to neurocognitive changes or target a common pathophysiology underlying both affective and neurocognitive mechanisms.

Lastly, some evidence grows concerning the role of *non-invasive brain stimulation techniques*, such as the transcranial magnetic stimulation (TMS), in the management of neurocognitive impairment in some neurological conditions. These interventions seem to modulate neuroplasticity processes. Deep transcranial magnetic stimulation (DTMS) is a novel alternative

to repetitive TMS which has been associated with small short-term improvement in sustained attention and larger improvement in spatial and visuospatial memory in unipolar depression (Minichino et al., 2012). Concerning BD, in a small open study Harel et al. detected a beneficial effect on working memory and psychomotor speed in a group of depressed bipolar patients after receiving 20 sessions of high-frequency DTMS when compared to healthy participants (Harel et al., 2011). Even so, DTMS in this group of patients also showed short-term antidepressant and anxiolytic effects, so it is difficult to conclude about the possible positive effect of DTMS on cognitive performance. Concerning the use of transcranial direct current stimulation (tDCS), on the one hand, repetitive sessions of tDCS in a group of euthymic bipolar patients provided preliminary results that concomitant prefrontal excitatory and cerebellar-inhibitory tDCS might have a positive effect on visuospatial memory (Minichino et al., 2015). Nevertheless, this study had significant limitations as a lack of a sham control group and a limited neuropsychological assessment. On the other hand, an intra-individual cross-over study failed in proving the efficacy of a single session of tDCS improving working memory and attention in euthymic bipolar patients (Martin et al., 2015). These last results would be in contrast to those reported in other studies conducted with healthy participants. Therefore, studies with brain stimulation techniques are still rare in BD and more studies will be required to assess the magnitude of effects compared to placebo and the durability of them. Even so, these extant studies raise several methodological considerations to keep in mind for further studies in order to achieve a potential cognitive enhancement.

As has been mentioned previously, due to the fact that cognitive reserve may contribute to functional and cognitive outcome, it has also been suggested that the implementation of intervention that allow the modification of CR in early stages of the illness might appear as a new target in bipolar patients in order to prevent a progressive impairment (Forcada et al., 2015).

While research in all these abovementioned areas proceed, it will also be important that clinicians prevent neurocognitive impairment through different steps. After achieving the remission of an acute episode it will be necessary to use an effective pharmacotherapy for relapse prevention, implement psychoeducation programs to avoid multiple episodes, and promote healthy habits. Another step towards mitigating cognitive impairment is via the treatment of those subclinical symptoms that may also impact cognitive function and functioning (see Table 2.).

### **FUTURE PROSPECTS**

Treating cognitive impairment in BD has become an important issue in the management of the patients, since it is a critical factor in psychosocial disability and lower quality of life for patients suffering this illness (Fountoulakis et al., 2016). The presence of these deficits in early stages of the disorder also indicates that cognitive dysfunction is a target for an early identification and intervention. Likewise, it is important to better elucidate the determinants of cognitive impairment which await further research in other potential factors influencing cognition. Nevertheless, not all the bipolar patients suffer cognitive dysfunction, therefore more research into cognitive heterogeneity is an important issue to have more valid and homogeneous neurocognitive phenotypes and a better understanding of those factors that may influence cognition and contribute to variability. The study of variability in non-chronic samples and in early stages (for instance, at first episodes or even at high-risk populations) will be needed in order to establish prevention strategies and to avoid a possible progression decline. Likewise, it is also necessary to study the longitudinal trajectories of the different neurocognitive subgroups. The characterization of neurocognitive profiles may also facilitate the emergence of genetic and neurobiological studies to delineate more valid subtypes of BD. Moreover, all these kind of studies may lead to a better definition of subgroups and would provide with helpful guidance for developing more effective pharmacological or



psychotherapeutic interventions contributing to enhance the management of the illness. A narrowly characterization of subtypes may also allow a better identification of key factors responsible for therapeutic response to treatments. Despite the previous research investigating new pharmacological and non-pharmacological treatments over the past decade, no robust evidence of therapeutic interventions targeting cognitive deficits is currently available, due to insufficient data, and further research is needed to be largely explored and draw firm conclusions. Meanwhile, functional remediation appears as a good option to ameliorate psychosocial outcome in bipolar patients. The combination of potential pro-cognitive drugs with cognitive/functional remediation is another via to be explored in bipolar patients since the combination of both may produce more robust efficacy with cognitive enhancers increasing the physiological mechanisms through which cognitive/functional remediation produce its therapeutics effects. In this line, there is a published protocol study about a randomized controlled trial to examine the utility of combining cognitive remediation and d-cycloserine (an NMDA receptor partial agonist) in individuals with BD (Breitborde et al., 2014). Unfortunately, results have not been published yet. Lurasidone, vortioxetine, omega-3-fatty acids, modafinil, vitamin-D, EPO, aspirine, and several other compounds are currently under investigation. Ultimately, novel neuromodulatory techniques deserve further research to be considered potential treatments to mitigate cognitive deficits in bipolar patients.

**Funding**

This work was supported by grants from the Spanish Ministry of Economy and Competitiveness (grant numbers PI11/00637, PI12/00912, PI15/00330) PN 2008–2011, Instituto de Salud Carlos III, Subdirección General de Evaluación y Fomento de la Investigación; Fondo Europeo de Desarrollo Regional. Unión Europea, “Una manera de hacer Europa”; CIBERSAM; and the Comissionat per a Universitats i Recerca del DIUE de la Generalitat de Catalunya (2014 SGR 398 to the Bipolar Disorders Group). Dr. Anabel Martinez-Aran’s project is supported, in part, by a 2013 NARSAD, Independent Investigator Grant from the Brain & Behavior Research Foundation. Dr. Carla Torrent is funded by the Spanish Ministry of Economy and Competitiveness, Instituto Carlos III, through a ‘Miguel Servet’ postdoctoral contract (CPI14/00175) and a FIS (PI 12/01498). Dr Torrent’s project is also supported in part by a 2014 NARSAD, Independent Investigator Grant from the Brain & Behavior Research Foundation (grant number 22039). Dr. I. Grande has received a Juan Rodés Contract (JR15/00012), Instituto de Salud Carlos III, Spanish Ministry of Economy and Competitiveness, Barcelona, Spain.

The other authors report no financial relationships with commercial interests.

**Acknowledgments**

The authors thank the support of the Spanish Ministry of Economy and Competitiveness, Instituto de Salud Carlos III, CIBERSAM, the Secretaria d’Universitats i Recerca del Departament d’Economia i Coneixement (2014\_SGR\_398), CERCAProgramme / Generalitat de Catalunya and the 2013 & 2014 NARSAD, Independent Investigator Grant from the Brain & Behavior Research Foundation.

**Conflict of interest**

Dr. Martinez-Aran has served as speaker or advisor for the following companies: Bristol-Myers Squibb, Otsuka, Lundbeck and Pfizer.

Dr. Vieta has received grants, CME-related honoraria, or consulting fees from AB-Biotics, Actavis, Alexza, Almirall, Allergan, AstraZeneca, Bristol-Myers Squibb, Cephalon, Dainippon Sumitomo Pharma, Eli Lilly, Ferrer, Forest Research Institute, Gedeon Richter, GlaxoSmith-Kline, Janssen, Janssen-Cilag, Jazz, Johnson & Johnson, Lundbeck, Merck, Novartis, Organon, Otsuka, Pfizer, Pierre-Fabre, Qualigen, Roche, Sanofi-Aventis, Schering-Plough, Servier, Shire, Solvay, Sunovion, Takeda, Telefónica, Teva, the Spanish Ministry of Science and Innovation (CIBERSAM), the Brain and Behaviour Foundation, the Seventh European Framework Programme (ENBREC), the Stanley Medical Research Institute, United Biosource Corporation, and Wyeth. The other authors report no financial relationships with commercial interests.

Dr. Carvalho received speaker honoraria from Abbott, GlaxoSmithKline, Lundbeck, Pfizer and Ache.

## References

- Aas M, Dazzan P, Fisher HL, Morgan C, Morgan K, Reichenberg A, Zanelli J, Fearon P, Jones PB, Murray RM, Pariante CM (2011) Childhood trauma and cognitive function in first-episode affective and non-affective psychosis. *Schizophr Res* 129: 12–9.
- Amoretti S, Bernardo M, Bonnin CM, Bioque M, Cabrera B, Mezquida G, Sole B, Vieta E, Torrent C (2016) The impact of cognitive reserve in the outcome of first-episode psychoses: 2-year follow-up study. *Eur Neuropsychopharmacol* 26:1638-1648.
- Anaya C, Torrent C, Caballero FF, Vieta E, Bonnin CM, Ayuso-Mateos JL (2016) Cognitive reserve in bipolar disorder: relation to cognition, psychosocial functioning and quality of life. *Acta Psychiatr Scand* 133:386-398.
- Arts B, Jabben N, Krabbendam L, van Os J (2008) Meta-analyses of cognitive functioning in euthymic bipolar patients and their first-degree relatives. *Psychol Med* 38: 771–785.
- Balanzá-Martínez V, Rubio C, Selva-Vera G, Martínez-Arán A, Sánchez-Moreno J, Salazar-Fraile J, Vieta E, Tabarés-Seisdedos R (2008) Neurocognitive endophenotypes (endophenocognities) from studies of relatives of bipolar disorder subjects: a systematic review. *Neurosci Biobehav Rev* 32:1426-1438.
- Balanzá-Martínez V, Selva G, Martínez-Arán A, Prickaerts J, Salazar J, González-Pinto A, Vieta E, Tabarés-Seisdedos R (2010) Neurocognition in bipolar disorders-A closer look at comorbidities and medications. *Eur J Pharmacol* 626: 87–96.
- Baune BT, Malhi GS (2015) A review on the impact of cognitive dysfunction on social, occupational, and general functional outcomes in bipolar disorder. *Bipolar Disord* 17: 41–55.
- Berk M, Conus P, Lucas N, Hallam K, Malhi GS, Dodd S, Yatham LN, Yung A, McGorry P (2007)

Setting the stage: from prodrome to treatment resistance in bipolar disorder. *Bipolar Disord* 9:, 671–678.

Bonnín CM, Martínez-Arán A, Torrent C, Pacchiarotti I, Rosa AR, Franco C, Murru A, Sanchez-Moreno J, Vieta E (2010) Clinical and neurocognitive predictors of functional outcome in bipolar euthymic patients: A long-term, follow-up study. *J Affect Disord* 121: 156–160.

Bonnín CM, Sánchez-Moreno J, Martínez-Arán A, Solé B, Reinares M, Rosa AR, Goikolea JM, Benabarre A, Ayuso-Mateos JL, Ferrer M, Vieta E, Torrent C (2011) Subthreshold symptoms in bipolar disorder : Impact on neurocognition , quality of life and disability. *J Affect Disord* 136:650-659.

Bonnín CM, González-Pinto A, Solé B, Reinares M, González-Ortega I, Alberich S, Crespo JM, Salamero M, Vieta E, Martínez-Arán A, Torrent C; CIBERSAM Functional Remediation Group (2014) Verbal memory as a mediator in the relationship between subthreshold depressive symptoms and functional outcome in bipolar disorder. *J Affect Disord* 160: 50–54.

Bonnin CM, Reinares M, Martinez-Aran A, Balanza-Martinez V, Sole B, Torrent C, Tabarés-Seisdedos R, García-Portilla MP, Ibáñez A, Amann BL, Arango C, Ayuso-Mateos JL, Crespo JM, González-Pinto A, Colom F, Vieta E; CIBERSAM Functional Remediation Group (2015) Effects of functional remediation on neurocognitively impaired bipolar patients: Enhancement of verbal memory. *Psychol Med* 46:291-301.

Bonnin CM, Torrent C, Arango C, Amann BL, Sole B, Gonzalez-Pinto A, Crespo JM, Tabarés-Seisdedos R, Reinares M, Ayuso-Mateos JL, García-Portilla MP, Ibañez Á, Salamero M, Vieta E, Martinez-Aran A; CIBERSAM Functional Remediation Group (2016) Functional remediation in bipolar disorder: 1-year follow-up of neurocognitive and functional outcome. *Br J Psychiatry* 208:87–93.

- Bora E (2016) Differences in cognitive impairment between schizophrenia and bipolar disorder : Considering the role of heterogeneity. *Psychiatry Clin Neurosci* 70:424-433.
- Bora E, Pantelis C (2015) Meta-analysis of cognitive impairment in first-episode bipolar disorder: Comparison with first-episode schizophrenia and healthy controls. *Schizophr Bull* 41: 1095–1104.
- Bora E, Pantelis C (2016) Social cognition in schizophrenia in comparison to bipolar disorder : A meta-analysis. *Schizophr Res* 175:72–78.
- Bora E, Yucel M, Pantelis C (2009) Cognitive endophenotypes of bipolar disorder: A meta-analysis of neuropsychological deficits in euthymic patients and their first-degree relatives. *J Affect Disord* 113: 1–20.
- Bora E, Yücel M, Pantelis C (2010) Neurocognitive markers of psychosis in bipolar disorder: a meta-analytic study. *J Affect Disord* 127:1–9.
- Bora E, Yücel M, Pantelis C, Berk M (2011) Meta-analytic review of neurocognition in bipolar II disorder. *Acta Psychiatr Scand* 123:165–174.
- Bora E, Hıdıroğlu C, Özerdem A, Kaçar ÖF, Sarısoy G, Civil Arslan F, Aydemir Ö, Cubukcuoglu Tas Z, Vahip S, Atalay A, Atasoy N, Ateşçi F, Tümkaya S (2016a) Executive dysfunction and cognitive subgroups in a large sample of euthymic patients with bipolar disorder. *Eur Neuropsychopharmacol* 26: 1338–1347.
- Bora E, Veznedaroğlu B, Vahip S (2016b) Theory of mind and executive functions in schizophrenia and bipolar disorder: A cross-diagnostic latent class analysis for identification of neuropsychological subtypes. *Schizophr Res* 176:500-505.
- Bortolato B, Miskowiak KW, Köhler CA, Vieta E, Carvalho AF (2015) Cognitive dysfunction in bipolar disorder and schizophrenia: a systematic review of meta-analyses. *Neuropsychiatr*

Dis Treat 11:3111–3125.

Bourne C, Aydemir O, Balanza-Martinez V, Bora E, Brissos S, Cavanagh JT, Clark L, Cubukcuoglu Z, Dias VV, Dittmann S, Ferrier IN, Fleck DE, Frangou S, Gallagher P, Jones L, Kieseppä T, Martínez-Aran A, Melle I, Moore PB, Mur M, et al. (2013) Neuropsychological testing of cognitive impairment in euthymic bipolar disorder: An individual patient data meta-analysis. *Acta Psychiatr Scand* 128:149–162.

Breitborde NJ, Dawson SC, Woolverton C, Dawley D, Bell EK, Norman K, Polsinelli A, Bernstein B, Mirsky P, Pletkova C, Grucci F 3rd, Montoya C, Nanadiego B, Sarabi E, DePalma M, Moreno F (2014) A randomized controlled trial of cognitive remediation and d-cycloserine for individuals with bipolar disorder. *BMC Psychol* 2: 41.

Bücker J, Kozicky J, Torres IJ, Kauer-Sant'anna M, Silveira LE, Bond DJ, Lam RW, Yatham LN (2013) The impact of childhood trauma on cognitive functioning in patients recently recovered from a first manic episode: data from the Systematic Treatment Optimization Program for Early Mania (STOP-EM). *J Affect Disord* 148:424–30.

Bücker J, Muralidharan K, Torres IJ, Su W, Kozicky J, Silveira LE, Bond DJ, Honer WG, Kauer-Sant'anna M, Lam RW, Yatham LN (2014) Childhood maltreatment and corpus callosum volume in recently diagnosed patients with bipolar I disorder: data from the Systematic Treatment Optimization Program for Early Mania (STOP-EM). *J Psychiatr Res* 48:65–72.

Budde M, Schulze TG (2014) Neurocognitive Correlates of the Course of Bipolar Disorder. *Harv Rev Psychiatry* 22: 342–347.

Burdick KE, Russo M, Frangou S, Mahon K, Braga RJ, Shanahan M, Malhotra AK (2014) Empirical evidence for discrete neurocognitive subgroups in bipolar disorder: clinical implications. *Psychol Med* 44:3083–3096.

- Catalá-López F, Gènova-Maleras R, Vieta E, Tabarés-Seisdedos R (2013) The increasing burden of mental and neurological disorders. *Eur Neuropsychopharmacol* 23: 1337–1339.
- Craddock N, Owen MJ (2010) The Kraepelinian dichotomy - going, going... but still not gone. *Br J Psychiatry* 196:92–95.
- Cuthbert BN (2014) The RDoC framework: Facilitating transition from ICD/DSM to dimensional approaches that integrate neuroscience and psychopathology. *World Psychiatry* 13: 28–35.
- Daban C, Martinez-Aran A, Torrent C, Tabarés-Seisdedos R, Balanzá-Martínez V, Salazar-Fraile J, Selva-Vera g, Vieta E (2006) Specificity of cognitive deficits in bipolar disorder versus schizophrenia. A systematic review. *Psychother Psychosom* 75: 72–84.
- Daruy-Filho L, Brietzke E, Lafer B, Grassi-Oliveira R (2011) Childhood maltreatment and clinical outcomes of bipolar disorder. *Acta Psychiatr Scand* 124:427–34.
- Deckersbach T, Nierenberg AA, Kessler R, Lund HG, Ametrano RM, Sachs G, Rauch SL, Dougherty D (2010) RESEARCH: Cognitive rehabilitation for bipolar disorder: An open trial for employed patients with residual depressive symptoms. *CNS Neurosci Ther* 16:298–307.
- Demant KM, Vinberg M, Kessing LV, Miskowiak KW (2015) Effects of Short-Term Cognitive Remediation on Cognitive Dysfunction in Partially or Fully Remitted Individuals with Bipolar Disorder: Results of a Randomised Controlled Trial. *PloS One*, 10: e0127955.
- Depp CA, Strassnig M, Mausbach BT, Bowie CR, Wolyniec P, Thornquist MH, Luke JR, McGrath JA, Pulver AE, Patterson TL, Harvey PD (2014) Association of obesity and treated hypertension and diabetes with cognitive ability in bipolar disorder and schizophrenia. *Bipolar Disord* 16: 422–431.



- Forcada I, Mur M, Mora E, Vieta E, Bartrés-Faz D, Portella MJ (2015) The influence of cognitive reserve on psychosocial and neuropsychological functioning in bipolar disorder. *Eur Neuropsychopharmacol* 25: 214–222.
- Fountoulakis KN, Vieta E, Young A, Yatham L, Grunze H, Blier P, Moeller HJ, Kasper S (2016) The International College of Neuropsychopharmacology (CINP) treatment guidelines for bipolar disorder in adults (CINP-BD-2017), part 4: Unmet needs in the treatment of Bipolar disorder and recommendations for future research. *Int J Neuropsychopharmacol* pii: pyw072.
- Frangou S, Donaldson S, Hadjulis M, Landau S, Goldstein LH (2005) The Maudsley Bipolar Disorder Project: executive dysfunction in bipolar disorder I and its clinical correlates. *Biol Psychiatry* 58: 859–864.
- Fuentes-Durá I, Balanzá-Martínez V, Ruiz-Ruiz JC, Martínez-Arán A, Girón M, Solé B, Sánchez-Moreno J, Gómez-Beneyto M, Vieta E, Tabarés-Seisdedos R (2012) Neurocognitive training in patients with bipolar disorders: Current status and perspectives. *Psychother Psychosom* 81: 250–252.
- Glahn DC, Bearden CE, Barguil M, Barrett J, Reichenberg A, Bowden CL, Soares JC, Velligan DI (2007) The neurocognitive signature of psychotic bipolar disorder. *Biol Psychiatry* 62 910–916.
- Goodwin, GM, Martinez-aran A, Glahn DC, Vieta E (2008) Cognitive impairment in bipolar disorder : Neurodevelopment or neurodegeneration ? An ECNP expert meeting report. *Eur Neuropsychopharmacol* 18: 787–793.
- Goswami U, Sharma A, Varma A, Gulrajani C, Ferrier IN, Young AH, Gallagher P, Thompson JM, Moore, PB (2009) The neurocognitive performance of drug-free and medicated euthymic bipolar patients do not differ. *Acta Psychiatr Scand* 120: 456–463.

- Gualtieri CT, Johnson LG (2006) Comparative neurocognitive effects of 5 psychotropic anticonvulsants and lithium. *MedGenMed* 8: 46.
- Harel EV, Zangen A, Roth Y, Reti I, Braw Y, Levkovitz Y (2011) H-coil repetitive transcranial magnetic stimulation for the treatment of bipolar depression: an add-on, safety and feasibility study. *World J Biol Psychiatry* 12: 119–26.
- Hellvin T, Sundet K, Simonsen C, Aminoff SR, Lagerberg TV, Andreassen OA, Melle I (2012) Neurocognitive functioning in patients recently diagnosed with bipolar disorder. *Bipolar Disord* 14:227–238.
- Jensen JH, Knorr U, Vinberg M, Kessing LV, Miskowiak KW (2016) Discrete neurocognitive subgroups in fully or partially remitted bipolar disorder: Associations with functional abilities. *J Affect Disord* 205:378–386.
- Kapczinski F, Vieta E, Cristina A, Frey BN, Gomes FA, Tramontina J, Kauer-Sant'anna M, Grassi-Oliveira R, Post RM (2008) Allostatic load in bipolar disorder: Implications for pathophysiology and treatment 32:675–692.
- Keefe RS, McClintock SM, Roth RM, Doraiswamy PM, Tiger S, Madhoo M (2014) Cognitive effects of pharmacotherapy for major depressive disorder: a systematic review. *J Clin Psychiatry* 75: 864–876.
- Kozicky JM, Torres IJ, Silveira LE, Bond DJ, Lam RW, Yatham LN (2014) Cognitive change in the year after a first manic episode: association between clinical outcome and cognitive performance early in the course of bipolar I disorder. *J Clin Psychiatry* 75: e587-93.
- Kucyi A, Alsuwaidan MT, Liauw SS, McIntyre RS (2010) Aerobic physical exercise as a possible treatment for neurocognitive dysfunction in bipolar disorder. *Postgrad Med* 122: 107–116.

- Kurtz MM, Gerraty RT (2009) A Meta-Analytic Investigation of Neurocognitive Deficits in Bipolar Illness : Profile and Effects of Clinical State. *Neuropsychology* 23: 551–562.
- Lackner N, Bengesser SA, Birner A, Painold A, Fellendorf FT, Platzer M, Reininghaus B, Weiss EM, Mangge H, McIntyre RS, Fuchs D, Kapfhammer HP, Wallner-Liebmann SJ, Reininghaus EZ (2015) Abdominal obesity is associated with impaired cognitive function in euthymic bipolar individuals. *World J Biol Psychiatry* 17:535-546.
- Lahera G, Ruiz-Murugarren S, Iglesias P, Ruiz-Bennasar C, Montes M, Ferna A (2012) Social Cognition and Global Functioning in Bipolar Disorder. *J Nerv Ment Dis* 200:135–141.
- Lahera G, Benito A, Montes JM, Fernandez-Liria A, Olbert CM, Penn DL (2013) Social cognition and interaction training (SCIT) for outpatients with bipolar disorder. *J Affect Disord* 146:132–136.
- Lee RSC, Hermens DF, Scott J, Redoblado-Hodge MA, Naismith SL, Lagopoulos J, Griffiths KR, Porter MA, Hickie IB (2014). A meta-analysis of neuropsychological functioning in first-episode bipolar disorders. *J Psychiatr Res* 57:1–11.
- Levy B (2013) Autonomic nervous system arousal and cognitive functioning in bipolar disorder. *Bipolar Disord* 15: 70–79.
- Levy B, Monzani BA, Stephansky MR, Weiss RD (2008) Neurocognitive impairment in patients with co-occurring bipolar disorder and alcohol dependence upon discharge from inpatient care. *Psychiatry Res* 161:28–35.
- Levy B, Medina AM, Weiss RD (2013) Cognitive and psychosocial functioning in bipolar disorder with and without psychosis during early remission from an acute mood episode: a comparative longitudinal study. *Compr Psychiatry* 54: 618–626.
- Lewandowski KE, Sperry SH, Cohen BM, Öngür D (2014) Cognitive variability in psychotic

disorders : a cross-diagnostic cluster analysis. *Psychol Med* 44:3239–3248.

Lewandowski KE, Sperry SH, Ongur D, Cohen BM, Norris LA, Keshavan MS (2016) Cognitive remediation versus active computer control in bipolar disorder with psychosis: study protocol for a randomized controlled trial. *Trials* 17:136.

Lichtenstein P, Yip BH, Björk C, Pawitan Y, Cannon TD, Sullivan PF, Hultman CM (2009) Common genetic determinants of schizophrenia and bipolar disorder in Swedish families: a population-based study. *Lancet* 373:234–239.

Liu CS, Carvalho AF, Mansur RB, McIntyre RS (2013) Obesity and Bipolar Disorder: Synergistic Neurotoxic Effects? *Adv Ther* 30: 987–1006.

López-Jaramillo C, Lopera-Vásquez J, Gallo A, Ospina-Duque J, Bell V, Torrent C, Martínez-Aran A, Vieta, E. (2010). Effects of recurrence on the cognitive performance of patients with bipolar I disorder: implications for relapse prevention and treatment adherence. *Bipolar Disord* 12:557–67.

Malchow B, Reich-Erkelenz D, Oertel-Knöchel V, Keller K, Hasan A, Schmitt A, Scheewe TW, Cahn W, Kahn RS, Falkai P (2013) The effects of physical exercise in schizophrenia and affective disorders. *Eur Arch PsychiatryClin Neurosci* 263:451–467.

Marshall DF, Passarotti AM, Ryan KA, Kamali M, Saunders EFH, PesterB, McInnis MG, Langenecker SA (2016) Deficient inhibitory control as an outcome of childhood trauma. *Psychiatry Res*:235, 7–12.

Martin DM, Chan H, Green MJ, Mitchell PB, Alonzo A, Loo C (2015) Transcranial direct current stimulation (tDCS) to remediate cognitive function in euthymic bipolar disorder patients. *Bipolar Disord* 17:849-858.

Martínez-Aran A, Vieta E (2015) Cognition as a target in schizophrenia, bipolar disorder and

depression. *Eur Neuropsychopharmacol* 25: 151–157.

Martinez-Aran A, Vieta E, Torrent C, Sanchez-Moreno J, Goikolea JM, Salamero M, Malhi GS, Gonzalez-Pinto A, Daban C, Alvarez-Grandi S, Fountoulakis K, Kaprinis G, Tabares-Seisdedos R, Ayuso-Mateos JL (2007) Functional outcome in bipolar disorder: the role of clinical and cognitive factors. *Bipolar Disord* 9: 103–113.

Martinez-Aran A, Torrent C, Tabares-Seisdedos R, Salamero M, Daban C, Balanza-Martinez V, Sanchez-Moreno J, Manuel Goikolea J, Benabarre A, Colom F, Vieta E (2008) Neurocognitive impairment in bipolar patients with and without history of psychosis. *J Clin Psychiatry* 69:233–239.

Martino DJ, Strejilevich SA, Scápola M, Igoa A, Marengo E, Ais ED, Perinot L (2008) Heterogeneity in cognitive functioning among patients with bipolar disorder. *J Affect Disord* 109:149–156.

Martino DJ, Marengo E, Igoa A, Scápola M, Ais ED, Perinot L, Strejilevich SA (2009) Neurocognitive and symptomatic predictors of functional outcome in bipolar disorders: a prospective 1 year follow-up study. *J Affect Disord* 116:37–42.

Martino DJ, Samamé C, Ibañez A, Strejilevich SA (2015) Neurocognitive functioning in the premorbid stage and in the first episode of bipolar disorder: a systematic review. *Psychiatry Res* 226:23–30.

Millan MJ, Goodwin GM, Meyer-Lindenberg A, Ove Ögren S (2015) Learning from the past and looking to the future: Emerging perspectives for improving the treatment of psychiatric disorders. *Eur Neuropsychopharmacol* 25: 599–656.

Minichino A, Bersani FS, Capra E, Pannese R, Bonanno C, Salviati M, Delle Chiaie R, Biondi M (2012) ECT, rTMS, and deepTMS in pharmacoresistant drug-free patients with unipolar

depression: a comparative review. *Neuropsychiatr Dis Treat* 8:55–64.

Minichino A, Bersani FS, Bernabei L, Spagnoli F, Vergnani L, Corrado A, Taddei I, Biondi M, Delle Chiaie R (2015) Prefronto-cerebellar transcranial direct current stimulation improves visuospatial memory, executive functions, and neurological soft signs in patients with euthymic bipolar disorder. *Neuropsychiatr Dis Treat* 11:2265–2270.

Miskowiak KW, Carvalho AF, Vieta E, Kessing LV (2016) Cognitive enhancement treatments for bipolar disorder: A systematic review and methodological recommendations. *Eur Neuropsychopharmacol* 26:1541-1561.

Moore RC, Harmell AL, Harvey PD, Bowie CR, Depp CA, Pulver AE, McGrath JA, Patterson TL, Cardenas V, Wolyniec P, Thornquist MH, Luke JR, Palmer BW, Jeste DV, Mausbach BT (2015) Improving the understanding of the link between cognition and functional capacity in schizophrenia and bipolar disorder. *Schizophrenia Research* 169:121–127.

Nejtek VA, Kaiser KA, Zhang B, Djokovic M (2013) Iowa Gambling Task scores predict future drug use in bipolar disorder outpatients with stimulant dependence. *Psychiatry Res* 210:871–879.

Nery FG, Hatch JP, Monkul ES, Matsuo K, Zunta-Soares GB, Bowden CL, Soares JC (2013) Trait impulsivity is increased in bipolar disorder patients with comorbid alcohol use disorders. *Psychopathology* 46: 145–152.

Passos IC, Mwangi B, Vieta E, Berk M, Kapczinski F (2016) Areas of controversy in neuroprogression in bipolar disorder. *Acta Psychiatr Scand* 134:91–103.

Popovic D, Vieta E, Fornaro M, Perugi G (2015) Cognitive tolerability following successful long term treatment of major depression and anxiety disorders with SSRI antidepressants. *J Affect Disord* 173:211–215.

- Quidé Y, O'Reilly N, Rowland JE, Carr VJ, Elzinga BM, Green MJ (2016) Effects of childhood trauma on working memory in affective and non-affective psychotic disorders. *Brain Imaging Behav*, 1–14.
- Robinson LJ, Thompson JM, Gallagher P, Goswami U, Young AH, Ferrier IN, Moore PB (2006) A meta-analysis of cognitive deficits in euthymic patients with bipolar disorder. *J Affect Disord* 93:105–115.
- Rosa AR, Magalhães PVS, Czepielewski L, Sulzbach MV, Goi PD, Vieta E, Gama CS, Kapczinski F (2014) Clinical staging in bipolar disorder: Focus on cognition and functioning. *J Clin Psychiatry* 75: 450–456.
- Rosenblat JD, Kakar R, McIntyre RS (2015) The Cognitive Effects of Antidepressants in Major Depressive Disorder: A Systematic Review and Meta-Analysis of Randomized Clinical Trials. *Int J Neuropsychopharmacol* 19: 1-13
- Russo M, Mahon K, Shanahan M, Ramjas E, Solon C, Braga RJ, Burdick KE (2014) Affective temperaments and neurocognitive functioning in bipolar disorder. *J Affect Disord* 169:51–56.
- Russo M, Mahon K, Shanahan M, Solon C, Ramjas E, Turpin J, Burdick KE (2015) The association between childhood trauma and facial emotion recognition in adults with bipolar disorder. *Psychiatry Res* 229: 771–776.
- Rybakowski JK, Suwalska A (2010) Excellent lithium responders have normal cognitive functions and plasma BDNF levels. *Int J Neuropsychopharmacol* 13:617–622.
- Samamé C, Martino DJ, Strejilevich SA (2012) Social cognition in euthymic bipolar disorder: Systematic review and meta-analytic approach. *Acta Psychiatr Scand* 125:266–280.
- Samamé C, Martino DJ, Strejilevich SA (2014) Longitudinal course of cognitive deficits in bipolar

disorder: a meta-analytic study. *J Affect Disord* 164:130–138.

Samamé C, Martino DJ, Strejilevich SA (2015) An individual task meta-analysis of social cognition in euthymic bipolar disorders. *J Affect Disord* 173:146–153.

Sanches M, Bauer IE, Galvez JF, Zunta-Soares GB, Soares JC (2015) The management of cognitive impairment in bipolar disorder: current status and perspectives. *Am J Ther* 22: 477–486.

Sanchez-Moreno J, Martinez-Aran A, Colom F, Scott J, Tabares-Seisdedos R, Sugranyes G, Torrent C, Daban C, Benabarre A, Goikolea JM, Franco C, González-Pinto A, Ayuso-Mateos JL, Vieta E (2009) Neurocognitive dysfunctions in euthymic bipolar patients with and without prior history of alcohol use. *J Clin Psychiatry* 70: 1120–1127.

Savitz JB, van der Merwe L, Stein DJ, Solms M, Ramesar RS (2008) Neuropsychological task performance in bipolar spectrum illness: genetics, alcohol abuse, medication and childhood trauma. *Bipolar Disord* 10:479–94.

Selva G, Salazar J, Balanzá-Martínez V, Martínez-Arán A, Rubio C, Daban C, Sánchez-Moreno J, Vieta E, Tabarés-Seisdedos R (2007). Bipolar I patients with and without a history of psychotic symptoms: do they differ in their cognitive functioning? *J Psychiatr Res* 41:265–272.

Silva KL, Rovaris DL, Guimaraes-da-Silva PO, Victor MM, Salgado CA, Vitola ES, Contini V, Bertuzzi G, Picon FA, Karam RG, Belmonte-de-Abreu P, Rohde LA, Grevet EH, Bau CH (2014) Could comorbid bipolar disorder account for a significant share of executive function deficits in adults with attention-deficit hyperactivity disorder? *Bipolar Disord* 16:270–276.

Simonsen C, Sundet K, Vaskinn A, Birkenaes B, Engh JA, Færden A (2009) Neurocognitive



Dysfunction in Bipolar and Schizophrenia Spectrum Disorders Depends on History of Psychosis Rather Than Diagnostic Group. *Schizophr Bull* 37:73-83.

Solé B, Jiménez E (2015) Tratamiento antipsicótico y aspectos neurocognitivos. *Psiquiatria Biologica* 22:36–38.

Solé B, Martínez-Arán A, Torrent C, Bonnin CM, Reinares M, Popovic D, Sancehz-Moreno J, Vieta, E. (2011). Are bipolar II patients cognitively impaired? A systematic review. *Psychol Med* 41: 1791–1803. Retrieved from

Solé B, Bonnin CM, Mayoral M, Amann BL, Torres I, González-Pinto A, Jimenez E, Crespo JM, Colom F, Tabarés-Seisdedos R, Reinares M, Ayuso-Mateos JL, Soria S, Garcia-Portilla MP, Ibáñez Á, Vieta E, Martinez-Aran A, Torrent C; CIBERSAM Functional Remediation Group (2015) Functional remediation for patients with bipolar II disorder: improvement of functioning and subsyndromal symptoms. *Eur Neuropsychopharmacol* 25:257–264.

Solé B, Jiménez E, Torrent C, Bonnin CM, Torres I, Reinares M, Reinares M, Priego Á, Salamero M, Colom F, Varo C, Vieta E, Martínez-Arán A (2016) Cognitive variability in bipolar II disorder: who is cognitively impaired and who is preserved. *Bipolar Disord* 18: 288–299.

Stern Y (2009) Cognitive reserve. *Neuropsychologia*, 47: 2015–28.

Summers M, Papadopoulou K, Bruno S, Cipolotti L, Ron MA (2006) Bipolar I and bipolar II disorder: cognition and emotion processing. *Psychol Med* 36:1799–1809.

Torrent C, Martinez-Arán A, Bonnin CM, Reinares M, Daban C, Solé B, Rosa AR, Tabarés-Seisdedos R, Popovic D, Salamero M, Vieta E (2012). Long-Term Outcome of Cognitive Impairment in Bipolar Disorder. *J Clin Psychiatry* 73: e899–e905.

Torrent C, Bonnin CM, Martínez-Arán A, Valle J, Amann BL, González-Pinto A, Crespo JM, Ibáñez Á, Garcia-Portilla MP, Tabarés-Seisdedos R, Arango C, Colom F, Solé B, Pacchiarotti

- I, Rosa AR, Ayuso-Mateos JL, Anaya C, Fernández P, Landín-Romero R, Alonso-Lana S, et al. (2013) Efficacy of functional remediation in bipolar disorder: a multicenter randomized controlled study. *Am J Psychiatry* 170:852–859.
- Torres JJ, Defreitas VG, Defreitas CM, Anna MK, Bond DJ, Honer WG, Lam RW, Yatham LN (2010) Neurocognitive Functioning in Patients With Bipolar I Disorder Recently Recovered From a First Manic Episode. *J Clin Psychiatry* 71:1234-1242.
- Tse S, Chan S, Ng KL, Yatham LN (2014) Meta-analysis of predictors of favorable employment outcomes among individuals with bipolar disorder. *Bipolar Disord* 16:217–229.
- van Gorp WG, Altshuler L, Theberge DC, Wilkins J, Dixon W (1998) Cognitive impairment in euthymic bipolar patients with and without prior alcohol dependence. A preliminary study. *Arch Gen Psychiatry* 55:41–6.
- Vieta E (2015) Staging and psychosocial early intervention in bipolar disorder. *Lancet Psychiatry* 2: 483–485.
- Vieta E (2016) DSM-5.1. *Acta Psychiatr Scand* 134:187–188.
- Volkert J, Kopf J, Kazmaier J, Glaser F, Zierhut KC, Schiele MA, Kittel-Schneider S, Reif A (2015) Evidence for cognitive subgroups in bipolar disorder and the influence of subclinical depression and sleep disturbances. *Eur Neuropsychopharmacol* 25: 192–202.
- Wingo AP, Wingo TS, Harvey PD, Baldessarini RJ (2009) Effects of lithium on cognitive performance: A meta-analysis. *J Clin Psychiatry* 70: 1588–1597.
- Wu HI, Chang YH, Lai CC, Wu JY, Chen SL, Chu CH, Lee IH, Yeh TL, Tzeng NS, Huang SY, Yang YK, Lu RB (2011) The effect of comorbid anxiety disorder on neuropsychological function in bipolar II disorder. *Prog NeuroPsychopharmacol Biol Psychiatry* 35: 1841–1845.
- Wu M, Passos IC, Bauer IE, Lavagnino L, Cao B, Zunta-soares GB, Kapczinski F, Soares JC (2016)

Individualized identification of euthymic bipolar disorder using the Cambridge Neuropsychological Test Automated Battery ( CANTAB ) and machine learning. J Affect Disord 192:219–225.

Yim CY, Soczynska JK, Kennedy SH, Woldeyohannes HO, Brietzke E, McIntyre RS (2012) The effect of overweight / obesity on cognitive function in euthymic individuals with bipolar disorder. Eur Psychiatry 27:223–228.

Zyto S, Jabben N, Schulte PF, Regeer BJ, Kupka RW (2016) A pilot study of a combined group and individual functional remediation program for patients with bipolar i disorder. J Affect Disord 194:9–15.

Table 1. Moderators of cognitive deficits in bipolar disorder (BD)
Educational attainment and premorbid Intelligence quotient (proxy variables of cognitive reserve)
Clinical symptomatology (remission vs. acute episode)
Subclinical depressive symptoms
Psychotic symptoms
Bipolar diagnostic subtype
Psychiatric or medical comorbidity
Illness duration (chronicity)
Number of episodes
Pharmacological treatment
Childhood adversity

<b>Table 2. Potential prevention strategies in cognitive dysfunction in bipolar disorder (BD)</b>
<b>Prevention of multiple episodes with an effective pharmacotherapy and implementation of psychoeducation programs</b>
<b>Avoid concomitant medications that interfere with cognitive function</b>
<b>Treat subclinical depressive symptoms</b>
<b>Control comorbidity (mental and psychiatric)</b>
<b>Implement Cognitive or Functional remediation</b>
<b>Promote healthy habits</b>
<b>Aerobic physical exercise</b>
<b>Prescribe adjunctive pro-cognitive treatment</b>
<b>Use of noninvasive brain stimulation techniques (TMS, DTMS, tDCS)</b>

Table 2. TMS= Transcranial magnetic stimulation; DTMS= Deep transcranial magnetic stimulation; tDCS= Transcranial direct current stimulation.

Figure legends

**Figure 1:** neuroprogression of bipolar disorder and strategies to prevent or to treat cognitive dysfunction.




Table 1. Moderators of cognitive deficits in bipolar disorder (BD)
Educational attainment and premorbid IQ (proxy variables of cognitive reserve)
Clinical symptomatology (remission vs. acute episode)
Subclinical depressive symptoms
Psychotic symptoms
Bipolar diagnostic subtype
Psychiatric or medical comorbidity
Illness duration (chronicity)
Number of episodes
Pharmacological treatment
Childhood adversity

Table 1. IQ= intelligence quotient.

Table 2. Potential prevention strategies in cognitive dysfunction in bipolar disorder (BD)
Prevention of multiple episodes with an effective pharmacotherapy and implementation of psychoeducation programs
Avoid concomitant medications that interfere with cognitive function
Treat subclinical depressive symptoms
Control comorbidity (mental and psychiatric)
Implement Cognitive or Functional remediation
Promote healthy habits
Aerobic physical exercise
Prescribe adjunctive pro-cognitive treatment
Use of noninvasive brain stimulation techniques (TMS, DTMS, tDCS)

Table 2. TMS= Transcranial magnetic stimulation; DTMS= Deep transcranial magnetic stimulation; tDCS= Transcranial direct current stimulation.








Figure 1: neuroprogression of bipolar disorder and strategies for cognitive functioning.

254x190mm (96 x 96 DPI)