

beam (sum over initial spin orientations divided by the number of such orientations) and when we do not detect the final polarizations (sum over final spin orientations),

$$\frac{d\sigma}{dk'_a dk'_b} = \frac{k'_a \mu_{aA} \mu_{ac}}{k_a 4\pi^2 \hbar^4} \frac{1}{(2J_A + 1)(2j_a + 1)} \times \sum_{\substack{m_a, m'_a \\ M_A, m'_b}} \left| \sum_{m_b} \langle j_b m_b j_c M_A - m_b | J_A M_A \rangle T_{m_a, m_b}^{m'_a, m'_b}(k'_a, k'_b) \right|^2. \quad (6.47)$$

The sum above can be simplified a bit. Let us consider a single particular value of m_b in the sum over m_b ,

$$\begin{aligned} \sum_{m_a, m'_a, m'_b} \left| T_{m_a, m_b}^{m'_a, m'_b}(k'_a, k'_b) \right|^2 \sum_{M_A} \left| \langle j_b m_b j_c M_A - m_b | J_A M_A \rangle \right|^2 = \\ \frac{2J_A + 1}{2j_b + 1} \sum_{m_a, m'_a, m'_b} \left| T_{m_a, m_b}^{m'_a, m'_b}(k'_a, k'_b) \right|^2 \\ \times \sum_{M_A} \left| \langle J_A - M_A j_c M_A - m_b | j_b m_b \rangle \right|^2, \end{aligned} \quad (6.48)$$

where we have used

$$\langle j_b m_b j_c M_A - m_b | J_A M_A \rangle = (-1)^{j_c - M_A + m_b} \sqrt{\frac{2J_A + 1}{2j_b + 1}} \langle J_A - M_A j_c M_A - m_b | j_b m_b \rangle. \quad (6.49)$$

As

$$\sum_{M_A} \left| \langle J_A - M_A j_c M_A - m_b | j_b m_b \rangle \right|^2 = 1, \quad (6.50)$$

we finally have

$$\frac{d\sigma}{dk'_a dk'_b} = \frac{k'_a \mu_{aA} \mu_{ac}}{k_a 4\pi^2 \hbar^4} \frac{1}{(2j_b + 1)(2j_a + 1)} \sum_{m_a, m'_a, m'_b} \left| \sum_{m_b} T_{m_a, m_b}^{m'_a, m'_b}(k'_a, k'_b) \right|^2. \quad (6.51)$$

Zero range approximation.

The zero range approximation consists in taking $v(r_{ab}) = D_0 \delta(r_{ab})$. Then, (see (2.147))

$$\begin{aligned} \mathbf{r}_{aA} &= \frac{c}{A} \mathbf{r}_{bc}, \\ \mathbf{r}_{ac} &= \mathbf{r}_{bc}. \end{aligned} \quad (6.52)$$

The angular dependence of the integral can be readily evaluated. From (6.20), noting that $\hat{\mathbf{r}}_{aA} = \hat{\mathbf{r}}_{ac} = \hat{\mathbf{r}}_{bc} \equiv \hat{\mathbf{r}}$,

$$\begin{aligned} [Y^{l_a}(\hat{\mathbf{r}}) Y^{l'_a}(\hat{\mathbf{r}})]_M^K [Y^{l_b}(\hat{\mathbf{r}}) Y^{l'_b}(\hat{\mathbf{r}})]_{-M}^K = \\ \frac{(-1)^{K-M}}{\sqrt{2K+1}} \left\{ [Y^{l_a}(\hat{\mathbf{r}}) Y^{l'_a}(\hat{\mathbf{r}})]^K [Y^{l_b}(\hat{\mathbf{r}}) Y^{l'_b}(\hat{\mathbf{r}})]^K \right\}_0^0. \end{aligned} \quad (6.53)$$

G.G. 2

We can as before evaluate this expression in the configuration shown in Fig. ~~6.6~~ ($\hat{\mathbf{r}} = \hat{\mathbf{z}}$), but now the multiplicative factor is 4π . The corresponding contribution to the integral is

$$\frac{(-1)^K}{4\pi(2K+1)} \langle l_a 0 l'_a 0 | K 0 \rangle \sqrt{(2l_a+1)(2l'_a+1)(2l_b+1)(2l'_b+1)}, \quad (6.F.54)$$

and

$$\begin{aligned} T_{m_a, m_b}^{m'_a, m'_b}(\mathbf{k}'_a, \mathbf{k}'_b) &= \frac{16\pi^2}{k_a k'_a k'_b} \frac{c}{A} D_0 T_\sigma \sum_{l_a, j_a} \sum_{l'_a, j'_a} \sum_{l_b, j_b} \sum_{l'_b, j'_b} e^{i(\sigma^{l_a} + \sigma^{l'_a} + \sigma^{l_b})} i^{l_a - l'_a - l_b} (-1)^{l_a + l'_a + l_b - j'_a - j'_b} \\ &\times \sqrt{(2l_a+1)(2l'_a+1)(2l_b+1)(2l'_b+1)} \langle l_a 0 l'_a 0 | K 0 \rangle \\ &\times \frac{2l_a+1}{2K+1} ((l'_a \frac{1}{2})_{j'_a} (l_a \frac{1}{2})_{j_a} | (l_a l'_a)_K (\frac{1}{2} \frac{1}{2})_0 \rangle_K ((l'_b \frac{1}{2})_{j'_b} (l_b \frac{1}{2})_{j_b} | (l_b l'_b)_K (\frac{1}{2} \frac{1}{2})_0 \rangle_K \\ &\times \langle l'_a m_a - m'_a - M \frac{1}{2} m'_a | j'_a m_a - M \rangle \langle l'_b m_b - m'_b + M \frac{1}{2} m'_b | j'_b M + m_b \rangle \\ &\times \langle l 0 \frac{1}{2} m_a | j m_a \rangle Y_{M+m_b+m'_b}^{l'_b}(\hat{\mathbf{k}}'_b) Y_{m_a+m'_a-M}^{l'_a}(\hat{\mathbf{k}}'_a) \mathcal{I}_{ZR}(l_a, l'_a, l'_b, j_a, j'_a, j'_b), \quad (6.F.55) \end{aligned}$$

where now the 1-dimensional integral to solve is

$$\mathcal{I}_{ZR}(l_a, l'_a, l'_b, j_a, j'_a, j'_b) = \int dr u_{l_b}(r) F_{l_a, j_a}(\frac{c}{A} r) F_{l'_a, j'_a}(r) F_{l'_b, j'_b}(r) / r. \quad (6.F.56)$$

6.F.3 One particle transfer

It may be interesting to state the expression for the one particle transfer reaction within the same context and using the same elements, in order to better compare these two type of experiments. In particle transfer, the final state of b is a bounded state of the $B(= a + b)$ nucleus, and we can carry on in a similar way as done previously just by substituting the distorted wave (continuum) wave function (6.F.34) with

$$\psi_{m'_b}^{l'_b, j'_b*}(\mathbf{r}_{ab}, \sigma_b) = u_{l'_b, j'_b}^*(r_{ab}) [Y_{l'_b}^{j'_b}(\hat{\mathbf{r}}_{ab}) \phi^{1/2}(\sigma_b)]_{m'_b}^{j'_b*}, \quad (6.F.57)$$

so the transition amplitude is now

$$\begin{aligned} T_{m_a, m_b}^{m'_a, m'_b}(\mathbf{k}'_a) &= \frac{8\pi^{3/2}}{k_a k'_a} \sum_{\sigma_a, \sigma_b} \sum_{l_a, j_a} \sum_{l'_a, m'_a, j'_a} e^{i(\sigma^{l_a} + \sigma^{l'_a})} i^{l_a - l'_a} (-1)^{l_a + l'_a - j'_a - j'_b} \\ &\times \sqrt{2l_a+1} \langle l'_a m'_a \frac{1}{2} m'_a | j'_a m'_a + m'_a \rangle \langle l_a 0 \frac{1}{2} m_a | j_a m_a \rangle \\ &\times Y_{-m'_a}^{l'_a}(\hat{\mathbf{k}}'_a) \int d\mathbf{r}_{aA} d\mathbf{r}_{bc} [Y_{l'_a}^{j'_a}(\hat{\mathbf{r}}_{bc}) \phi^{1/2}(\sigma_a)]_{-m'_a}^{j'_a} [Y_{l'_b}^{j'_b}(\hat{\mathbf{r}}_{ab}) \phi^{1/2}(\sigma_b)]_{-m'_b}^{j'_b} \\ &\times \frac{F_{l_a, j_a}(r_{aA}) F_{l'_a, j'_a}(r_{bc})}{r_{bc} r_{aA}} u_{l'_b, j'_b}^*(r_{ab}) u_{l_b, j_b}(r_{bc}) u(r_{ab}) u_\sigma(\sigma_a, \sigma_b) \\ &\times [Y_{l_a}^{j_a}(\hat{\mathbf{r}}_{aA}) \phi^{1/2}(\sigma_a)]_{m_a}^{j_a} [Y_{l_b}^{j_b}(\hat{\mathbf{r}}_{bc}) \phi^{1/2}(\sigma_b)]_{m_b}^{j_b}. \quad (6.F.58) \end{aligned}$$

no he encontrado
la referencia a la
Figura 6.F.4
en el texto




Figure 6.F.4: One particle transfer reaction $A(= c + b) + a \rightarrow B(= a + b) + c$.

6

Using (6.F.39), (6.F.40), (6.F.41), and setting $M = m_a - m'_a - m'_b$

$$\begin{aligned}
 T_{m_a, m'_b}^{m'_a, m'_b}(\mathbf{k}'_a) &= \frac{8\pi^{3/2}}{k_a k'_a} T_\sigma \sum_{l_a, j_a} \sum_{l'_a, j'_a} \sum_{K, M} e^{i(\sigma^{l_a} + \sigma^{l'_a})} i^{l_a - l'_a} (-1)^{l_a + l'_a - j'_a - j_b} \\
 &\quad \times ((l'_a \frac{1}{2})_{j'_a} (l_a \frac{1}{2})_{j_a} | (l_a l'_a)_K (\frac{1}{2} \frac{1}{2})_0 \rangle ((l'_b \frac{1}{2})_{j'_b} (l_b \frac{1}{2})_{j_b} | (l_b l'_b)_K (\frac{1}{2} \frac{1}{2})_0 \rangle_K \\
 &\quad \times \sqrt{2l_a + 1} \langle l'_a m_a - m'_a - M \ 1/2 \ m'_a j'_a m_a - M \rangle \langle l_a \ 0 \ 1/2 \ m_a j_a m_a \rangle \\
 &\quad \times Y_{m_a - m'_a - M}^{l'_a}(\hat{\mathbf{k}}'_a) \int d\mathbf{r}_{aA} d\mathbf{r}_{bc} \frac{F_{l_a, j_a}(r_{aA}) F_{l'_a, j'_a}(r_{bc})}{r_{bc} r_{aA}} u_{l'_b, j'_b}^*(r_{ab}) u_{l_b, j_b}(r_{bc}) v(r_{ab}) \\
 &\quad \times [Y^{l_a}(\hat{\mathbf{r}}_{aA}) Y^{l'_a}(\hat{\mathbf{r}}_{bc})]_M^K [Y^{l_b}(\hat{\mathbf{r}}_{bc}) Y^{l'_b}(\hat{\mathbf{r}}_{ab})]_{-M}^K. \quad (6.F.59)
 \end{aligned}$$

Aside from (6.F.59), we also need

$$\mathbf{r}_{bc} = \frac{a+B}{B} \mathbf{r}_{aA} + \frac{b}{A} \mathbf{r}_{bc}. \quad (6.F.60)$$

From (6.F.20), (8.1.2), (8.1.1), (8.1.3), (7.2.121), we get

$$\begin{aligned}
 T_{m_a, m'_b}^{m'_a, m'_b}(\mathbf{k}'_a) &= \frac{32\pi^3}{k_a k'_a} T_\sigma \sum_{l_a, j_a} \sum_{l'_a, j'_a} \sum_{K, M} e^{i(\sigma^{l_a} + \sigma^{l'_a})} i^{l_a - l'_a} (-1)^{l_a + l'_a - j'_a - j_b} \\
 &\quad \times ((l'_a \frac{1}{2})_{j'_a} (l_a \frac{1}{2})_{j_a} | (l_a l'_a)_K (\frac{1}{2} \frac{1}{2})_0 \rangle ((l'_b \frac{1}{2})_{j'_b} (l_b \frac{1}{2})_{j_b} | (l_b l'_b)_K (\frac{1}{2} \frac{1}{2})_0 \rangle_K \\
 &\quad \times \frac{2l_a + 1}{2K + 1} \langle l'_a m_a - m'_a - M \ 1/2 \ m'_a j'_a m_a - M \rangle \\
 &\quad \times \langle l_a \ 0 \ 1/2 \ m_a j_a m_a \rangle Y_{m_a - m'_a - M}^{l'_a}(\hat{\mathbf{k}}'_a) I(l_a, l'_a, j_a, j'_a, j'_b, K), \quad (6.F.61)
 \end{aligned}$$

with

$$\begin{aligned}
 I(l_a, l'_a, j_a, j'_a, K) &= \int d\mathbf{r}_{aA} d\mathbf{r}_{bc} d\theta r_{aA}^2 \frac{\sin \theta}{r_{bc}} \\
 &\quad \times F_{l_a, j_a}(r_{aA}) F_{l'_a, j'_a}(r_{bc}) u_{l'_b, j'_b}^*(r_{ab}) u_{l_b, j_b}(r_{bc}) v(r_{ab}) \\
 &\quad \times \sum_{M_K} \langle l_a \ 0 \ l'_a \ M_K | K \ M_K \rangle [Y^{l_b}(\cos \theta, 0) Y^{l'_b}(\cos \theta_{ab}, 0)]_{-M_K}^K Y_{M_K}^{l'_a}(\cos \theta_{bc}, 0), \quad (6.F.62)
 \end{aligned}$$

where (see (7.2.127), (6.F.60) and Fig. 7.42)

$$\cos \theta_{ab} = \frac{-r_{aA} - \frac{c}{A} r_{bc} \cos \theta}{\sqrt{\left(\frac{c}{A} r_{bc} \sin \theta\right)^2 + \left(r_{aA} + \frac{c}{A} r_{bc} \cos \theta\right)^2}}, \quad (6.F.63)$$

$$\cos \theta_{bc} = \frac{\frac{a+B}{B} r_{aA} + \frac{b}{A} r_{bc} \cos \theta}{\sqrt{\left(\frac{b}{A} r_{bc} \sin \theta\right)^2 + \left(\frac{a+B}{B} r_{aA} + \frac{b}{A} r_{bc} \cos \theta\right)^2}}, \quad (6.F.64)$$

and

$$r_{bc} = \sqrt{\left(\frac{b}{A} r_{bc} \sin \theta\right)^2 + \left(\frac{a+B}{B} r_{aA} + \frac{b}{A} r_{bc} \cos \theta\right)^2}. \quad (6.F.65)$$

Again, this is nothing new as many codes exist which deal with one particle transfer within the same DWBA formalism we have used here, but it may be useful to have our own code to better compare transfer and knock-out experiments. By the way, (6.F.61) can also be used when particle b populates a resonant state in the continuum of nucleus B .

✓ Appendix G: H Modified form factors

G.H.1 Two-particle transfer

(cf. p p. -11_a-
and -11_b- right
after p. 11)

G.H.2 One-particle transfer

G.H.3 One-particle knockout

G.H.4 Inelastic scattering

G.H.5 Elastic scattering

Likely it is better to have it
as an Appendix of Ch. 7 (2n-transfer)

Bibliography

- ✓ P. W. Anderson. *Basic notions of condensed matter physics*. Benjamin, Menlo Park, California, 1984.
- ✓ D. Bès and R. Broglia. In (A. Bohr and R. Broglia, editors), *International School of Physics "Enrico Fermi" Course LXIX, Elementary Modes of Excitation in Nuclei*, page 55, Amsterdam, 1977. North Holland.
- ✓✓ A. Bohr and B. R. Mottelson. *Nuclear Structure, Vol. II*. Benjamin, New York, 1975.
- pp. 55-134 ✓ D. Brink and R. A. Broglia. *Nuclear Superfluidity*. Cambridge University Press, Cambridge, 2005.
- ✓✓ R. Broglia and A. Winther. *Heavy Ion Reactions*. Westview Press, Cambridge, MA., 2004.
- ✓ R. Broglia, O. Hansen, and C. Riedel. Two-neutron transfer reactions and the pairing model. *Advances in Nuclear Physics*, 6:287, 1973. URL <http://arXiv.org/abs/nucl-th/9305001>
- R. Broglia, G. Coló, G. Onida, and H. Roman. *Solid State Physics of Finite Systems: metal clusters, fullerenes, atomic wires*. Springer Verlag, Berlin, Heidelberg, 2004.
- ✓ C. Mahaux et al. Dynamics of the shell model. *Physics Reports*, 120:1-274, 1985.
- ✓ R. Feynman. *Theory of fundamental processes*. Benjamin, Reading, Mass., 1975.
- I. Tanihata et al. Measurement of the two-halo neutron transfer reaction $^1\text{H}(^{11}\text{Li}, ^9\text{Li})^3\text{H}$ at 3A MeV. *Phys. Rev. Lett.*, 100:192502, 2008.
- G. Potel, F. Barranco, E. Vigezzi, and R. A. Broglia. Evidence for phonon mediated pairing interaction in the halo of the nucleus ^{11}Li . *Phys. Rev. Lett.*, 105:172502, 2010.
- ✓✓ G. Satchler. *Introduction to Nuclear Reactions*. Mc Millan, New York, 1980.
- italics
- ✓ Duguet, T. and Hagen, G. (2012) *Ab initio* approach to effective single-particle energies in doubly closed shell nuclei, *Phys. Rev. C* 85, 034330
- change as is now in the Review paper
- ✓ Jennings, B.K. (2011) Non observability of spectroscopic factors, arXiv: 1102.3721v2 [nucl-th]
- ✓ Dickhoff, W.H. and Barbieri, C. (2004) Self-consistent Green's function method for nuclei and nuclear matter, *Progress in Particle and Nuclear Physics* 52, 377-496

• Makhamedzhanov, A.M. and Kadyrov, A.S. (2010)
Unitary correlation in nuclear reaction theory:
separation of nuclear reactions and spectroscopic factors, Phys. Rev. C82, 051601.

• Furnstahl, R.J. and Schwenk, A. (2010)
How should one formulate, extract and interpret "non-observables" for nuclei?
J. of Phys. G 37, 064005 (6pp)

→ Schiffer, J.P., Hoffman, C.R., Kay, B.P., et al
(2012) Test of sum rules in nucleon transfer reactions, Phys. Rev. Lett. 108, 022501.

• Bang, J., Ershov, S.N., Gareev, F.A. -- (1980)
Discrete expansion of continuum wave-functions
Nucl. Phys. A 339, 89-124.

• Vaegen, J.S., Nilsson, B.S., Bang, J., -- (1979)
One-nucleon and two-nucleon overlaps
generated by a Sturmian method,
Nucl. Phys. A 319, 143-156.

• Brink, D.M. and Satchler, G.R. (1968) Angular Momentum, Clarendon Press, Oxford.

• Edmonds, A.R. (1960) Angular momentum in Quantum mechanics, Princeton University Press, Princeton, New Jersey.

• Bohr, A. and Mottelson, B.R. (1969) Nuclear Structure, Vol I, Benjamin, Amsterdam.

✓ Bortignon, P.F., Bracco, A. and Broglia, R.A. (1998) Giant Resonances, Harwood Academic Publishers, Amsterdam.

✓ Bertsch, G.F., Bortignon, P.F. and Broglia, R.A. (1983)
--- ^{title} --- Rev. Mod. Phys. 55 : 287.

✓ Migdal, A. (1958) Interaction between electrons and lattice vibrations in a normal metal ~~£~~, Soviet Physics JETP, 7, 996-1001.

Bardeen, J. and Pines, D. (1955) Electron-phonon interaction in metals, Phys. Rev. 99, 1140-1150

Fröhlich, H. (1952) Interaction of electrons ~~with~~ with lattice vibrations, Proc. Royal Soc. (London) A215, 291-297

Cobb W. R. and Guthe D.B., (1957)
Angular distribution of protons
from the $^{44}\text{Ca}(d,p)^{45}\text{Ca}$ reaction,
Phys. Rev. 107, 181~~44~~