

# Algebra: Chapter 0 Exercises

## Chapter 1, Section 4

David Melendez

February 9, 2017

**Problem 4.1.** Composition is defined for *two* morphisms. If more than two morphisms are given, e.g.:

$$A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{h} D \xrightarrow{i} E$$

then one may compose them in several ways, for example:

$$(ih)(gf), \quad (i(hg))f, \quad i((hg)f), \quad \text{etc.}$$

so that at every step one is only composing two morphisms. Prove that the result of any such nested composition is independent of the placement of the parentheses.

*Solution.* Let  $Z_m \in \text{Obj}(C)$  and  $f_m \in \text{Hom}(Z_{m+1}, Z_m)$  for every  $m \in \mathbb{N}$ . Let  $n$  be the number of morphisms we're composing. We will use induction on  $n$ .

Base case: Suppose  $n = 3$ . Then, since  $C$  is a category, we have  $f_1(f_2f_3) = (f_1f_2)f_3$ .

Induction: Suppose that all parenthesizations of  $f_1, \dots, f_{j-1}$  under composition are equivalent for all  $1 \leq j < n$ . Then, for some  $1 < k \leq n$ , let  $\alpha$  be some parenthesization of  $f_1, \dots, f_{k-1}$ , and let  $\beta$  be some parenthesization of  $f_k, \dots, f_n$ . Any parenthesization of  $f_1, \dots, f_n$  will then be of the form  $\alpha\beta$ . By associativity and our inductive hypothesis, we have  $\alpha = ((f_k \dots f_{n-1})f_n)$ . Thus, by associativity, we have

$$\begin{aligned} \alpha\beta &= (f_1 \dots f_{k-1}) ((f_k \dots f_{n-1})f_n) \\ &= ((f_1 \dots f_{k-1})(f_k \dots f_{n-1})) f_n \\ &= ((\dots ((f_1f_2)f_3) \dots) f_n \end{aligned}$$

as desired. ■