

ARTICLE TYPE

This is the sample article title[†]

Author One^{*1} | Author Two^{2,3} | Author Three³

¹Org Division, Org Name, State name,
Country name

²Org Division, Org Name, State name,
Country name

³Org Division, Org Name, State name,
Country name

Correspondence

*Corresponding author name, This is sample
corresponding address. Email:
authorone@gmail.com

Present Address

This is sample for present address text this is
sample for present address text

Summary

This is sample abstract text this is sample abstract text this is sample abstract text
this is sample abstract text this is sample abstract text this is sample abstract text
this is sample abstract text this is sample abstract text this is sample abstract text
this is sample abstract text this is sample abstract text this is sample abstract text
this is sample abstract text this is sample abstract text this is sample abstract text
this is sample abstract text this is sample abstract text this is sample abstract text
this is sample abstract text this is sample abstract text this is sample abstract text
this is sample abstract text this is sample abstract text this is sample abstract text
this is sample abstract text this is sample abstract text this is sample abstract text
this is sample abstract text this is sample abstract text this is sample abstract text
this is sample abstract text.

KEYWORDS:

keyword1, keyword2, keyword3, keyword4

1 | INTRODUCTION

2 | METHODOLOGY

3 | SIMULATION STUDY

4 | APPLICATION TO SPANISH DATA

- We now apply our changepoint detection algorithm to data from Spain.
- In Section 4.1, we introduce the dataset of temperatures and precipitation indexes used in our application.
- Section 4.3 shows the results of our initial screening step, while in Section 4.4, we discuss the results of our changepoint detection algorithm.
- Finally, Section 4.5 shows the clustering results for seasons before and after identified changepoints.

4.1 | Spanish Data

- We analyse temperature and precipitation observations from 40 meteorological stations distributed across mainland Spain.
- Their locations are shown in the left panel of Figure 1 .
- The analysis covers the period 1960-2020, inclusive.

[†]This is an example for title footnote.
⁰Abbreviations: ANA, anti-nuclear antibodies; APC, antigen-presenting cells; IRF, interferon regulatory factor

- Only stations satisfying the required record-length and data-availability criteria are retained. **State criteria for choosing stations/locations (description in code).**
- Daily maximum temperature and precipitation totals are obtained from the European Climate Assessment Dataset (ECAD) ([European Climate Assessment & Dataset \(ECA&D\) n.d.](#)).
- The ECAD provides quality-controlled daily meteorological observations from stations across Europe.
- We aggregate the daily observations to a weekly resolution, taking the maximum recorded temperature and total accumulated precipitation within each week.
- This aggregation reduces the influence of very short-range temporal variation and decreases the proportion of zero precipitation observations, although it does not necessarily remove temporal dependence completely.
- The relationship between high temperature and dry conditions may depend more strongly on precipitation accumulated over preceding weeks than on precipitation observed during the same week.
- We therefore represent precipitation conditions using the Standardised Precipitation Index (SPI) of [McKee, Doesken, and Kleist \(1993\)](#), rather than using weekly precipitation totals directly.
- At each station, we calculate SPI from precipitation accumulated over a rolling 13-week window.
- Thus, the value assigned to a given week summarises precipitation during that week and the preceding 12 weeks, providing a measure of precipitation conditions over an approximately three-month period.
- The accumulated precipitation is modelled using a Gamma distribution, whose parameters are estimated using unbiased probability-weighted moments.
- Its fitted cumulative probability is then transformed to a standard Gaussian scale.
- Under the usual SPI convention, negative values represent drier-than-normal conditions and positive values represent wetter-than-normal conditions.
- For the dependence analysis, we therefore define the drought index [$D = -\text{SPI}$,] so that larger values correspond to drier conditions.
- This sign convention means that high temperatures and dry conditions are expected to exhibit positive extremal dependence.
- The right-hand panels of Figure 1 illustrate the relationship between weekly maximum temperature and the drought index at Daroca, in Aragon, and Jerez de la Frontera, in Andalusia.
- These stations are highlighted in the map in the left-hand panel.
- At both stations, the distribution of weekly maximum temperature differs substantially between seasons.
- The seasonal separation in drought-index values is less pronounced at Daroca, whereas Jerez de la Frontera exhibits clearer seasonal variation in both temperature and drought conditions. **Mention explicit figures for means/95th quantiles??**
- These seasonal differences motivate fitting the dependence model and applying the changepoint procedure separately within each season.
- This allows both the marginal behaviour and the extremal dependence structure to vary seasonally.




FIGURE 1 Left: Elevation (m) profile of mainland Spain along with locations of 40 meteorological stations. Daroca and Jerez de la Frontera highlighted. Right: Weekly maximum temperature against 13-week drought index, ($D=-SPI$), at these two stations, with observations coloured by season.

4.2 | Setup

- The conditional extremes framework is applied after transforming each variable to a common marginal distribution.
- Consequently, systematic changes in the marginal distributions over time could otherwise be confounded with changes in the extremal dependence structure.
- Both variables exhibit seasonal variation, while weekly maximum temperature in particular also displays a long-term marginal trend.
- We therefore preprocess the marginal distributions before fitting the conditional extremes models.
- At each station, and for each variable, we consider observations from each calendar month separately and fit a linear trend in year.
- The estimated trend is removed from the observations for that month.
- Treating months separately allows the magnitude of the temporal trend to vary throughout the year while accounting for the pronounced marginal seasonality.
- This detrending removes an estimated linear change in the location of each monthly marginal distribution; it does not imply that all temporal dependence has been removed.
- In particular, some short-range dependence may remain between successive weekly observations, and consecutive values of the 13-week drought index are constructed from overlapping precipitation windows.
- Following detrending, the observations are grouped by meteorological season.
- For each station and variable, the resulting seasonal observations are transformed standard Laplace margins using the empirical rank transform.
- The conditional extremes and changepoint analyses are then performed separately for winter, spring, summer and autumn.
- We fit the conditional extremes model of [Heffernan and Tawn \(2004\)](#) using a common dependence threshold equal to the theoretical 80th percentile of the standard Laplace distribution.
- The same threshold is used for both variables, at every station, in both temporal blocks and at every candidate changepoint.
- This ensures that exceedances have a common interpretation throughout the analysis.

- The 80th-percentile threshold provided a satisfactory empirical compromise between restricting attention to the joint tail and retaining enough exceedances for stable model estimation.
- In particular, the resulting clustering solutions exhibited clear spatial contiguity. **Add a sensitivity analysis for the dependence threshold.**
- The Monte Carlo approximation used to construct the pairwise dissimilarities employs the same sample from the standard Laplace distribution in every temporal comparison.
- Holding this sample fixed avoids introducing additional Monte Carlo variation between candidate changepoints.
- The conditional fits from the two conditioning directions are combined into the aggregated dissimilarity matrix defined in Section ?? **. Test for others**
- All screening, testing and clustering calculations reported below use this matrix. **Confirm terminology and add sensitivity results for the alternative dissimilarity matrices, if required.**
- For the Spanish application, the permutation procedure is adapted to preserve the temporal and spatial structure within each season.
- Rather than permuting individual weekly observations, we permute complete season-years.
- For spring, summer and autumn, the season-year is the calendar year in which the observations occur.
- Winter spans two calendar years; December is therefore assigned to the same season-year as the following January and February.
- At a candidate season-year (t), the two temporal blocks consist of the (25) season-years ending at (t) and the (25) season-years beginning at ($t + 1$), respectively.
- Thus, (t) denotes a possible change between season-years (t) and ($t + 1$).
- In each pointwise permutation, the 50 season-years from the two blocks are pooled and randomly divided into two pseudo-blocks containing 25 season-years each.
- Every weekly observation belonging to a given season-year is moved as a single unit, and the same reassignment is applied simultaneously to both variables and all 40 stations.
- This preserves the dependence between observations within a season-year and the spatial correspondence between stations.
- The validity of this permutation procedure relies on treating season-years as exchangeable under the null hypothesis of no change in the extremal dependence structure.
- Although permuting complete season-years preserves within-year dependence, it does not preserve any residual dependence between successive season-years.
- The choice of block length involves a bias-variance trade-off.
- Shorter blocks provide better temporal localisation but contain fewer threshold exceedances, resulting in unstable or failed conditional extremes fits.
- Longer blocks provide more stable estimates but leave fewer feasible candidate changepoints and smooth changes over a wider temporal interval.
- We investigated block lengths of 15, 20, 25 and 30 season-years.
- With 15-year blocks, many candidate comparisons failed because at least one conditional fit contained too few exceedances.

- Twenty-year blocks were reliable for winter but produced substantial failure rates for several other seasons.
- All candidate comparisons fitted successfully with 25-year blocks.
- Increasing the length to 30 years also produced stable fits, but left only two to five candidate changepoints per season.
- We therefore use 25 season-years on either side of each candidate throughout the primary analysis.
- This was the shortest common block length that produced stable fits across all seasons while retaining a useful range of candidate changepoints. **Present the alternative block lengths as a sensitivity analysis or in the supplementary material(?).**

4.3 | Screening

- We first perform a computationally less intensive screening step, in which the observed discrepancy between the two estimated dissimilarity matrices is calculated at every feasible candidate season-year without generating a permutation distribution.
- Figure 2 presents the screening statistics for each season using the Frobenius, infinity and spectral norms.
- As discussed in Section 2, these norms provide complementary summaries of the difference between the dissimilarity matrices on either side of a candidate change.
- The three norms have different numerical scales, and their absolute values should therefore not be compared directly.
- Instead, we compare the timing and relative prominence of features in their respective screening curves.
- Red points in Figure 2 identify local peaks.
- For a given norm (T), an interior candidate season-year (t) is classified as a local peak when

$$T(t) > T(t - 1) \quad \text{and} \quad T(t) \geq T(t + 1).$$

The first and final candidate years are not classified as local peaks because they cannot be compared with candidates on both sides.

- For winter, all three norms increase during the late 1990s and exhibit a prominent shared peak in 1998.
- A smaller common peak is present around 1992.
- The agreement among the norms makes the late-1990s period a particularly notable candidate region.
- For spring, the clearest screening peak occurs around 1992, for all norms.
- Smaller local peaks occur around 1989 and 1997.
- For summer, the screening curves contain several prominent peaks.
- The largest norm values occur in 1988, with further substantial peaks in 1992 and 1997.
- For Autumn, the screening results are less concentrated around a single candidate year.
- The Frobenius curve has local peaks around 1990 and 1996, while the infinity and spectral curves identify broadly similar candidate regions during the late 1980s or early 1990s and the mid-1990s.
- The precise locations of the peaks are less consistent across norms than for winter, spring or summer.
- Across the four seasons, the largest Frobenius norm screening value occurs for summer around 1988. **Cite exact value**
- The clearest agreement among all three norms occurs around 1992 for spring and around 1998 for winter.
- Although the screening step highlights several candidate regions of particular interest, the total number of feasible candidate season-years is sufficiently small to apply the permutation test at every candidate.
- We therefore retain the complete candidate range rather than restricting the formal analysis to the local screening peaks.




FIGURE 2 Screening discrepancies between the estimated extremal-dependence dissimilarity matrices on either side of each candidate season-year, calculated using 25 season-years per temporal block. Results are shown for the test statistic D using the Frobenius, infinity and spectral norms. A candidate (t) represents a possible change between season-years (t) and ($t + 1$). Red points denote interior local peaks satisfying $(T(t) > T(t - 1))$ and $(T(t) \geq T(t + 1))$.

4.4 | Changepoint detection

- We next apply the permutation test at every feasible candidate season-year.
- For each candidate and norm, the null distribution is estimated using 1000 permutations of complete season-years between the two 25-year blocks.
- Figure 3 shows the resulting pointwise Monte Carlo p-values for the Frobenius, infinity and spectral norms.
- A candidate season-year (t) represents a possible change between season-years (t) and ($t + 1$), and the horizontal dashed line denotes the pointwise 5% significance level.
- The p-values in Figure 3 are pointwise: each tests a change at one specified candidate season-year.
- For spring, none of the three norms produces a pointwise p-value below 0.05.
- The smallest values occur around 1992, in agreement with the dominant feature identified during screening, but remain above the pointwise significance threshold.
- Also for Autumn, none of the three norms produces a pointwise p-value below 0.05, although the infinity norm came very close for 1986, the year with the highest test statistic in our screening step.
- For winter, the infinity norm provides pointwise evidence of a change over the period 1996–1999.
- The spectral norm supports a narrower region around 1997–1998, while the Frobenius norm is significant in 1998.

- Thus, all three norms identify 1998, and the overall pattern provides consistent evidence for a change in the winter extremal dependence structure during the late 1990s.
- For summer, the infinity norm produces pointwise-significant p-values over an extended period from approximately 1987 to 1993.
- The Frobenius and spectral norms support narrower subsets of this region, with particularly small p-values around 1988–1990 and 1992.
- The agreement among the norms indicates that the early-1990s signal is not confined to a single choice of discrepancy statistic.
- After approximately 1994, the summer p-values generally increase, particularly for the infinity norm.
- These results provide no pointwise evidence for a change at these later candidates.
- However, the relatively large p-values should not be interpreted as affirmative evidence that the dependence structure is unchanged; they indicate only that the observed discrepancies are not unusual under the corresponding permutation null distributions.
- The infinity norm identifies broader pointwise-significant candidate ranges than the Frobenius and spectral norms for both summer and winter.
- As discussed in Section ??, this may indicate that some changes are more pronounced in particular entries of the dissimilarity matrix than across its entire structure.
- Nevertheless, the principal summer and winter candidate regions are supported by more than one norm.
- In particular, all three norms provide pointwise evidence around 1990 for summer and around 1998 for winter.
- Taken together, the results suggest candidate changes in the summer extremal dependence structure during the late 1980s or early 1990s and in the winter structure during the late 1990s.
- No comparable evidence is found for spring or autumn.

4.5 | Clustering Results

- We finally examine how the detected changes affect the spatial clustering of the fitted extremal dependence structure.
- For each fitted conditional extremes model, the aggregated dissimilarity matrix is used to partition the 40 stations into three clusters using the clustering procedure described in Section ??.
- As the changepoint analysis provides no consistent evidence of a change in spring or autumn, we refit the conditional extremes model to the complete time series for these seasons.
- For summer, the pointwise tests identify a broader candidate change region during the late 1980s and early 1990s.
- We use 1990 as a representative division within this region and fit separate models to season-years 1960–1990 and 1991–2020.
- For winter, the three norms provide their clearest common evidence at season-year 1998.
- We therefore fit separate models to winter season-years 1960–1998 and 1999–2020.
- These divisions treat a candidate (t) as a change between season-years (t) and ($t + 1$). **Confirm or update the selected divisions after completing the multiplicity-adjusted changepoint analysis.**
- Figure 4 presents the resulting spatial clusterings.




FIGURE 3 Pointwise permutation-test p-values for candidate changes in the seasonal extremal dependence structure, calculated using 25 season-years per temporal block and 200 permutations at each candidate. Results are shown for the Frobenius, infinity and spectral norms. A candidate (t) represents a possible change between season-years (t) and ($t + 1$). The horizontal dashed line denotes the pointwise 5% significance level.

- Although station coordinates and elevation do not directly determine the clustering assignments, the fitted clusters exhibit substantial spatial coherence.
- Neighbouring stations are frequently assigned to the same cluster, suggesting that the estimated extremal dependence structure contains a strong geographical component.
- The spring clustering broadly separates groups of stations along the northern and Mediterranean coasts (in blue) from groups in the western (orange) and central interior (red).
- Nevertheless, the boundaries are not determined solely by physical proximity, with some geographically separated stations assigned to the same dependence cluster.
- The autumn clustering exhibits a stronger broadly coastal–interior structure.
- Several stations along the northern and eastern coasts are grouped together, similar to the spring clustering, while the other two clusters comprise locations in the west/south-west and elsewhere, respectively.
- For summer, the two fitted periods retain some common large-scale structure.
- However, a number of stations change their clustering relationships after 1990, most visibly in northwestern Spain, along the southern coast and across parts of the western interior.
- The summer changepoint therefore corresponds to a reorganization of particular regional relationships rather than a complete replacement of the earlier clustering structure.

- The winter clustering, in contrast, changes more extensively across the 1998 division.
- In particular, the post-1998 fit produces a different grouping of many stations along the northern coast, in the western and central interior, and along the Mediterranean coast.
- This broader spatial reorganisation, as opposed to the more localised changes seen in summer, is consistent with the late-1990s signal being identified by all three norms in the changepoint analysis.




FIGURE 4 Spatial clustering of the 40 Spanish stations obtained from the fitted conditional extremes dissimilarity matrices. Spring and autumn are fitted using their complete time series. Summer is fitted separately for season-years 1960–1990 and 1991–2020, while winter is fitted separately for season-years 1960–1998 and 1999–2020. Three clusters are used in each fit. Colours distinguish cluster membership within each panel. Background shading represents elevation.

5 | DISCUSSION

ACKNOWLEDGMENTS

This is acknowledgment text [Elbaum, Malishevsky, and Rothermel \(February 2002\)](#). Provide text here. This is acknowledgment text. Provide text here. This is acknowledgment text. Provide text here. This is acknowledgment text. Provide text here. This is acknowledgment text. Provide text here. This is acknowledgment text. Provide text here. This is acknowledgment text. Provide text here.

Author contributions

This is an author contribution text. This is an author contribution text. This is an author contribution text. This is an author contribution text. This is an author contribution text.

Financial disclosure

None reported.

Conflict of interest

The authors declare no potential conflict of interests.

SUPPORTING INFORMATION

The following supporting information is available as part of the online article:

How to cite this article: Williams K., B. Hoskins, R. Lee, G. Masato, and T. Woollings (2016), A regime analysis of Atlantic winter jet variability applied to evaluate HadGEM3-GC2, *Q.J.R. Meteorol. Soc.*, 2017;00:1–6.

APPENDIX

A SECTION TITLE OF FIRST APPENDIX

References

- Elbaum, S., Malishevsky, A. G., & Rothermel, G. (February 2002). Test case prioritization: a family of empirical studies. *IEEE Transactions on Software Engineering*, 28(2), 159–182. doi: 12345.2345
- European Climate Assessment & Dataset (ECA&D). (n.d.). *Predefined subsets of daily station data in ASCII format*. <https://www.ecad.eu/dailydata/predefinedseries.php>. Accessed 30 June 2026.
- Heffernan, J. E., & Tawn, J. A. (2004). A conditional approach for multivariate extreme values (with discussion). *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 66(3), 497–546.
- McKee, T. B., Doesken, N. J., & Kleist, J. (1993). The relationship of drought frequency and duration to time scales. In *Proceedings of the eighth conference on applied climatology* (pp. 179–184). Anaheim, California: American Meteorological Society.