YOUR STUDY RESULTS

PHARVARIS

RAPIDe-3

Title: Study of Oral Deucrictibant Soft Capsule for On-Demand Treatment of Angioedema Attacks in Adolescents and Adults with Hereditary Angioedema

(RAPIDe-3)

Drug Studied: deucrictibant

- In this study, researchers tested deucrictibant, a new drug currently being studied, to see if it helped people with hereditary angioedema (HAE).
- 134 participants took part in the study. 113 participants treated 201 HAE attacks. Of these, 88 participants treated paired HAE attacks, ie. 176 HAE attacks and completed the study. 46 participants left the study before they completed the study.
- Results showed that participants in this study who took deucrictibant immediate release (IR) had greater HAE attack symptoms relief compared with placebo treatment, and that this difference was unlikely due to chance.
- No participants had severe or serious side effects considered related to deucrictibant IR within 3 days of treatment. No participants stopped taking study drug because of side effects.

THANK YOU!

At Pharvaris, we think it is important for you to know the results of your study. We hope it helps you understand the important role you played in supporting medical research.

This is a summary of the main results of your study. If you have questions about these results, the study doctor or staff at your study site may be able to answer them.

WHY WAS THIS STUDY DONE?

HAE is a rare, genetic condition that causes unpredictable and painful swelling attacks in the limbs, face, intestines, and airways. HAE attacks can potentially be life-threatening. Current treatment involves on-demand medications to stop attacks, and preventative therapies.


Deucrictibant is an investigational drug, taken by mouth. Deucrictibant is thought to work by stopping opening of a receptor on blood vessels called the bradykinin B_2 receptor. The bradykinin B_2 receptor is like a control valve that can stop or allow fluid to move out of blood vessels into the surrounding tissues. By stopping the opening of this receptor, deucrictibant

might stop leakage of fluids from the blood vessel into surrounding tissues, which is what happens during angioedema attacks.

Researchers are studying whether an on-demand treatment taken by mouth could provide an additional option for people with HAE. Currently, deucrictibant is only available in clinical studies. It is not approved by any health authorities as a treatment for HAE. The deucrictibant used in this study was an "immediate release" pill, also called "IR". This means that deucrictibant is designed to be absorbed into the blood stream quickly after it is taken. The IR pill might work well for on-demand usage.


This study included:

- Deucrictibant IR (20 mg) or placebo taken by mouth to treat 2 HAE attacks where participants did not have breathing difficulties (1 attack treated with deucrictibant IR and 1 attack treated with placebo).



The main purpose of this study was to see what effect deucrictibant IR had as an on-demand treatment for HAE attacks, compared to the effect of a placebo.

To measure the response to treatment, participants were asked to complete questionnaires asking them about their attack symptoms. These questionnaires are called Patient Global Impression of Change (PGI-C) and Patient Impression of Severity (PGI-S). These questionnaires help to standardize the results participants reported by using a simple scale.



The main question researchers asked in this study was:

1 **How long did it take participants to feel at least "a little better" for 2 consecutive timepoints within 12 hours after treatment during an HAE attack? This was measured using the PGI-C scale.**

Researchers compared results for participants when they took deucrictibant IR to when they took placebo.

Other questions asked in this study included:

2 **What effect did taking deucrictibant IR have compared with taking placebo on:**
   a. **The time it took for the HAE attack to get a PGI-C rating of at least "better" for 2 consecutive timepoints within 12 hours after treatment?**
   b. **The time it took to get at least a 1-point reduction in PGI-S rating for 2 consecutive timepoints within 12 hours after treatment?**
   c. **The time it took for the HAE attack to get a PGI-S rating of "none" (defined as complete resolution)?**
   d. **The time it took for the HAE attack to stop getting worse (based on PGI-C)?**

   Researchers compared participant's questionnaire results.

3 **How safe is deucrictibant IR compared with placebo?**
   Researchers compared the number and type of side effects participants reported.

WHAT HAPPENED IN THE STUDY?


This was a "phase 3" study, where researchers compare the study drug to a standard treatment or placebo. This study used a placebo.


WHO WAS IN THE STUDY?

Males and females aged 12 to 75 years who had a diagnosis of HAE, with:

- History of at least 2 HAE attacks in the last 3 months before screening
- Experience with using standard-of-care drug to effectively manage on-demand treatment of HAE attacks

134 participants were enrolled in the study:






- 58 (43%) participants were men and 76 (57%) participants were women

- 10 (7%) participants were under 18 years old; 116 (87%) participants were aged 18 to 64 years and 8 (6%) participants were aged 65 years or older


This study was done at 76 study centers in the following countries:



WHAT TREATMENT DID PARTICIPANTS GET?

The basic design of this study is shown here:

- Adult participants were placed in 1 of 2 treatment sequences:



The placebo looked like deucrictibant IR but did not contain any medicine.

- This type of study is called a “cross over” study, because the treatment that participants took changed over during the study.
- Adolescent participants received a single dose of deucrictibant IR by mouth to test how much ended up in the blood stream. They were then assigned to the 2 groups in the cross over phase of the study, as above.

Participants were assigned to the treatment sequence by equal chance.

- Group 1: 66 people were assigned to get deucrictibant IR *then* placebo
- Group 2: 68 people were assigned to get placebo *then* deucrictibant IR

During the study:

- 114 participants received at least 1 dose of deucrictibant IR or placebo
- 113 participants treated a total of 201 HAE attacks
    - 100 participants treated 1 HAE attack with deucrictibant IR
    - 101 participants treated 1 HAE attack with placebo
    - 88 participants treated 2 HAE attacks, either 1 with deucrictibant IR then 1 with placebo, or 1 with placebo then 1 with deucrictibant IR. These were referred to as “paired attacks”

This study was “double-blind”, neither the participants nor the study doctors knew which treatment sequence they were assigned to. This was done to make sure the study results were not influenced in any way.

HOW LONG DID THE STUDY LAST?

Each participant was to be treated for 2 HAE attacks with either deucrictibant IR or placebo for the first attack, and then the opposite treatment for the second attack.

Of the 134 participants who entered the study, 46 left the study before they completed the study. This included:

- 35 participants who left because Pharvaris had collected enough information and stopped the study
- 6 participants who chose to stop taking part
- 2 participants who could not be contacted
- 2 participants who did not meet the requirements of the study
- 1 participant who was randomized but did not receive any study drug.

88 participants completed the study with 2 paired HAE attacks.

Depending on how often attacks occurred in each individual person, the duration of time on the study varied. The entire study lasted for 18 months and most participants were in the study for about 6 months. Some were in the study for up to a year. The study began 04 April 2024 and ended 17 October 2025.


WHAT WERE THE STUDY RESULTS?

Question 1

How long did it take participants to feel at least "a little better" for 2 consecutive timepoints within 12 hours after treatment during an HAE attack? This was measured using the PGI-C scale.

To study this, researchers calculated the median time to start of symptom relief for participants when taking deucrictibant IR or placebo. The median is the middle of a set of numbers when the set of numbers is sorted into a list of biggest to smallest or smallest to biggest. It was not possible for researchers to calculate this time for participants who took placebo. This is because fewer than half of the participants who took placebo reached this level of improvement within the study's observation period (12 hours), and the median can only be calculated when at least half of participants reach this target.

- The median time it took participants to feel at least "a little better" on the PGI-C scale after taking deucrictibant IR to treat their attack was 1.28 hours. The median time could not be calculated for participants who took placebo.



❖ Based on these results, researchers concluded that the difference between deucrictibant IR and placebo on time to start of symptom relief was unlikely due to chance.

Question 2a

What effect did taking deucrictibant IR have compared with taking placebo on the time it took for the HAE attack to get a PGI-C rating of at least "better" for 2 consecutive timepoints within 12 hours after treatment?


- In this study, the median time it took participants to feel at least “better” on the PGI-C scale for 2 consecutive timepoints after taking deucrictibant IR to treat their attack was 2.85 hours. As above, the median time could not be calculated for participants who took placebo.



Question 2b

What effect did taking deucrictibant IR have compared with taking placebo on the time it took to get at least a 1-point reduction in PGI-S rating for 2 consecutive timepoints within 12 hours after treatment?

- In this study, the median time it took participants to get symptom relief of at least 1 point on the PGI-S scale for 2 consecutive timepoints after taking deucrictibant IR was 2.41 hours. As above, the median time could not be calculated for participants who took placebo.



Question 2c

What effect did taking deucrictibant IR have compared with taking placebo on the time it took for the HAE attack to get a PGI-S rating of “none” (defined as “complete resolution”)?


- The median time it took participants to get completely better after taking deucrictibant IR was 11.95 hours. It was not possible for researchers to calculate this time for participants who took placebo.



Question 2d

What effect did deucrictibant IR have compared with taking placebo on the time it took for the HAE attack to stop getting worse (based on PGI-C)?

- The median time it took participants to stop getting worse after taking deucrictibant IR was 17.47 minutes. After taking placebo, it took participants a median of 228.67 minutes to stop getting worse.



Researchers concluded that all these differences were unlikely due to chance. Deucrictibant IR may improve the symptoms and length of HAE attacks compared with placebo.

This does not mean that everyone in this study had these results, and individual results could be better or worse than the overall group. Other studies may have different results. These are just some of the main results of this study. More results may be available at the websites listed at the end of this summary.

DID PARTICIPANTS HAVE SIDE EFFECTS?

None of the participants had severe or serious side effects considered related to deucrictibant IR within 3 days of treatment.

None of the participants stopped deucrictibant IR treatment due to side effects.

Two (2) participants experienced a single event of fatigue within 3 days of treatment, one of which was considered unrelated to deucrictibant IR treatment.

Information about other side effects may be available at the websites listed at the end of this summary.

WHERE CAN I LEARN MORE ABOUT THIS STUDY?

Researchers will use the information from this study and other similar studies to help determine if deucrictibant is safe and effective at improving symptoms of HAE attacks and stopping progression of HAE attacks. Additional studies with deucrictibant are ongoing.

For more information about this study, check these websites:

www.clinicaltrials.gov Use the study identifier **NCT06343779**

https://euclinicaltrials.eu/ Use the study identifier **2023-507268-37-00**

Full Title of This Study: A Phase 3, Randomized, Double-blind, Placebo-controlled, Crossover Study of Oral Deucrictibant Soft Capsule for On-demand Treatment of Attacks in Adolescents and Adults with Hereditary Angioedema

Pharvaris Study ID: **RAPIDe-3**

Pharvaris Study Number: **PHA022121-C306**

Date of This Report: 15 April 2026

Thank you again for being a part of this study!

We do research to try to find the best way to help patients, and you helped us to do that