1.	EU study number and full study title: 2024-516247-62. A Phase 3, Randomized, Double-blind, Placebo-controlled Study to Evaluate the Efficacy and Safety of Orally Administered Deucrictibant Extended-release Tablet for Prophylaxis Against Angioedema Attacks in Adolescents and Adults with Hereditary Angioedema
2.	Rationale: This is a study to test the safety and efficacy of a daily 40 mg dose of deucrictibant in preventing attacks of hereditary angioedema (HAE) in participants aged ≥12 years
3.	Objectives: Primary Objective: • To evaluate the efficacy of a daily 40 mg dose regimen of deucrictibant for preventative treatment of HAE attacks Secondary Objectives: • To evaluate the safety of deucrictibant of a daily 40 mg dose regimen for preventative treatment of HAE attacks • To evaluate what the patient's body does to deucrictibant and what deucrictibant does to the patient and the patient's well-being when deucrictibant is taken for 24 weeks
4.	Main study endpoints: The primary endpoint is a comparison of the number of HAE attacks confirmed by the study doctor in patients taking deucrictibant or placebo per 4 weeks of treatment
5.	Secondary study endpoints: Secondary efficacy endpoints: • The number of HAE attacks confirmed by the study doctor from time of the first dose to the last dose of deucrictibant • The number of attacks that are moderate or severe from time of the first dose to the last dose of deucrictibant • The number of attacks that require treatment with the patient's usual therapies from time of the first dose to the last dose of deucrictibant • Number of patients who have ≥50%, 70%, or 90% reduction in number of HAE attacks or have no HAE attacks from time of the first dose to the last dose of deucrictibant • Number of patients who did not have HAE attacks from time of the first dose to the last dose of deucrictibant • Number of patients without angioedema symptoms from time of the first dose to the last dose of deucrictibant Secondary safety endpoints: The safety endpoints include comparisons of the patient's clinical laboratory tests, pulse, temperature, breathing rate, blood pressure, and electrocardiogram (a test that

	measures the electrical activity of the heart) before and after taking deucrictibant or the placebo to see if taking the drugs adversely affects the patient's health
6.	Study design: This is a study to test the safety and efficacy of a 40 mg dose of deucrictibant taken once a day in preventing attacks of HAE in patients aged ≥12 years. The study is being conducted at approximately 80 sites in North America and Europe. Patients take either placebo or 40 mg deucrictibant once a day for 24 weeks and then completes an End of Study Visit 4 weeks after the last dose of either placebo or deucrictibant. Any patient who stops taking either placebo or deucrictibant before 24 weeks will complete the End of Treatment Visit and will be asked to continue coming into the clinic for visits for the remainder of the 24 weeks before completing the End of Study Visit. Any patient who withdraws consent or is discontinued from the study at any time completes an End of Treatment Visit and End of Study Visit 4 weeks after the last dose of either placebo or deucrictibant
7.	Study population: Approximately 81 male or female patients, including at least 8 adolescents, with HAE are participating in this study.
8.	Interventions: Patients take either placebo or 40 mg deucrictibant once per day for 24 weeks. The patient and the study doctor do not know whether the patient is taking placebo or deucrictibant. Patients are allowed to treat any HAE attack using standard of care on-demand therapy. An electronic diary is used to record any symptoms and details of HAE attacks and treatments. The electronic diary is also used as a dosing diary to record each administration of either placebo or deucrictibant and to complete electronic patient-reported outcome questionnaires.
9.	Ethical considerations relating to the clinical study including the expected benefit to the individual subject or group of patients represented by the study subjects as well as the nature and extent of burden and risks: Patients in this study may experience risks due to collection of blood samples, small increase in blood pressure, decreased heart rate, and worsening of the blood flow to parts of the body (ischemia). In previous clinical studies, deucrictibant was generally safe and well-tolerated and no serious side effects were reported. Patients taking deucrictibant may have a positive benefit from taking this drug.