

In-Class Quiz 1 (Gaussian Process)

1 Problem 1

1.1 a

Show that if a gaussian process is WSS then it is SSS.

1.2 b

consider the following gaussian process:

$$X_0 \sim \mathcal{N}(0, \sigma^2)$$
$$X_n = \frac{1}{2}X_{n-1} + Z_n, n \geq 1$$

where Z_1, Z_2, \dots are iid $\mathcal{N}(0, 1)$ and independent of X_0 . find σ such that X_n is SSS: [Hint: You can assume that there is some value of σ that this process is SSS]

1.3 Solution a

A random process $X(t)$ is said to be SSS if all it's finite order distributions are time invariant. i.e., the joint cdf (pdf, pmf) of $X(t_1), X(t_2), \dots, X(t_k)$ and $X(t_1 + \tau), X(t_2 + \tau), \dots, X(t_k + \tau)$ are the same for all k, all t_1, \dots, t_k and all time shift τ .

A random process $X(t)$ said to be wide-sense stationary WSS if it's mean and autocorrelation function are time invariant.

- $E[X(t)] = \mu$ independent of t.
- $R_x(t_1, t_2)$ is only a function of $t_1 - t_2$.

For Gaussian random processes, WSS \Rightarrow SSS, since the process is completely specified by its mean and autocorrelation functions.

1.4 Solution B

This process is gaussian so it is SSS if it is WSS.

$$E[X_n] = 0$$

Assume $n \geq m$:

$$\begin{aligned} R_x[n, m] &= E[X_n X_m] = E\left[\left(\frac{1}{2}X_{n-1} + Z_n\right)\left(\frac{1}{2}X_{m-1} + Z_m\right)\right] \\ &= E\left[\frac{1}{4}X_{n-1}X_{m-1} + \frac{1}{2}X_{n-1}Z_m + \frac{1}{2}X_{m-1}Z_n + Z_m Z_n\right] \end{aligned}$$

For the second term:

$$E[\frac{1}{2}X_{n-1}Z_m] = E[\frac{1}{2}(\frac{1}{2}X_{n-2} + Z_{n-1})Z_m]$$

At some point $n - i = m$ and we know that $E[Z_i^2] = 1$

$$E[\frac{1}{2}X_{n-1}Z_m] = 2^{-(n-m)}$$

So we have:

$$R_x[n, m] = \frac{1}{4}R_x[n-1, m-1] + 2^{-(n-m)}$$

If this process wants to be wss then $R_x[m, n] = R_x[m-1, n-1]$:

$$R_x[n, m] = \frac{4}{3}2^{-(n-m)}$$

Let $m = n$:

$$R_x[n, n] = E[X_n^2] = Var(X_n) = Var(X_0) = \sigma^2 = \frac{4}{3}$$