

Quiz 2 (Gaussian Processes) Solutions

1 -a)

$$E[X(t)] = E\left[\int_0^t W(\tau) d\tau\right] = \int_0^t E[W(\tau)] d\tau = \int_0^t 0 d\tau = 0$$

$$\begin{aligned} R_X(t, t + \tau) &= E[X(t)X(t + \tau)] = E\left[\int_0^t \int_0^{t+\tau} W(u)W(v) dv du\right] \\ &= \int_0^t \int_0^{t+\tau} E[W(u)W(v)] dv du \\ &= \int_0^t \int_0^{t+\tau} \alpha\delta(u - v) dv du \end{aligned}$$

Case 1: $\tau \geq 0$

$$R_X(t, t + \tau) = \int_0^t \alpha du = \alpha t$$

Case 2: $\tau < 0$

$$R_X(t, t + \tau) = \int_0^{t+\tau} \int_0^t \alpha\delta(u - v) du dv = \int_0^{t+\tau} \alpha dv = \alpha(t + \tau)$$

$$\Rightarrow R_X(t, t + \tau) = \alpha \min\{t, t + \tau\}$$

1-b)

Since $\text{Var}(X(t)) = \alpha \cdot t$ is dependent on t , it cannot be stationary.

2)

Let $W = [W(t_1), W(t_2), \dots, W(t_n)]^T$ denote a vector of samples of a Brownian motion process. To prove that $W(t)$ is a Gaussian process, we must show that W is a Gaussian random vector. To do so, let

$$X = [X_1, X_2, \dots, X_n]^T = [W(t_1), W(t_2) - W(t_1), \dots, W(t_n) - W(t_{n-1})]^T$$

denote the vector of increments. By the definition of Brownian motion, X_1, X_2, \dots, X_n is a sequence of independent Gaussian random variables. Thus, X is a Gaussian random vector. Finally,

$$W = \begin{bmatrix} X_1 \\ X_1 + X_2 \\ \dots \\ X_1 + X_2 + \dots + X_n \end{bmatrix} = \begin{bmatrix} 1 & \dots & \dots & 0 \\ 1 & 1 & 0 & 0 \\ \vdots & \ddots & \ddots & \vdots \\ 1 & \dots & 1 & 1 \end{bmatrix} \cdot X$$

Since X is a Gaussian random vector and $W = AX$, where A is a rank n matrix, so W is also a Gaussian random vector.