

COMP2610 / COMP6261 - Information Theory

Lecture 9 - Probabilistic Inequalities

Assignment Project Exam Help

Robert C. Williamson

<https://powcoder.com>

Research School of Computer Science



Australian  
National  
University

Add WeChat powcoder

20 August, 2018

Last time

# Assignment Project Exam Help

Mutual information chain rule

Jensen's inequality

<https://powcoder.com>

"Information cannot hurt"

Data processing inequality

Add WeChat powcoder

## Review: Data-Processing Inequality

### Theorem

Assignment Project Exam Help

- $X$  is the state of the world,  $Y$  is the data gathered and  $Z$  is the processed data

<https://powcoder.com>

- No “clever” manipulation of the data can improve the inferences that can be made from the data

Add WeChat powcoder

- No processing of  $Y$ , deterministic or random, can increase the information that  $Y$  contains about  $X$

This time

# Assignment Project Exam Help

- Markov's inequality

<https://powcoder.com>

- Chebyshev's inequality

Add WeChat powcoder

- Law of large numbers

## Outline

1 Properties of expectation and variance

2 Markov's inequality

3 Chebyshev's inequality

4 Law of large numbers

5 Wrapping Up

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

- 1 Properties of expectation and variance

# Assignment Project Exam Help

- 2 Markov's inequality

- 3 Chebyshev's inequality

<https://powcoder.com>

- 4 Law of large numbers

Add WeChat powcoder

- 5 Wrapping Up

## Expectation and Variance

Let  $X$  be a random variable over  $\mathcal{X}$ , with probability distribution  $p$

Expected value:

$$\mathbb{E}[X] = \sum_{x \in \mathcal{X}} x \cdot p(x).$$

Variance:

$$\begin{aligned}\mathbb{V}[X] &= \mathbb{E}[(X - \mathbb{E}[X])^2] \\ &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2.\end{aligned}$$

Standard deviation is  $\sqrt{\mathbb{V}[X]}$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

## Properties of expectation

A key property of expectations is **linearity**:

$$\mathbb{E} \left[ \sum_{i=1}^n X_i \right] = \sum_{i=1}^n \mathbb{E} [X_i]$$

$$\text{LHS} = \sum_{x_1 \in \mathcal{X}_1} \dots \sum_{x_n \in \mathcal{X}_n} \left( p(x_1, \dots, x_n) \cdot \sum_{i=1}^n x_i \right)$$

This holds even if the variables are dependent!

We have for any  $a \in \mathbb{R}$ ,

$$\mathbb{E}[aX] = a \cdot \mathbb{E}[X].$$



## Properties of variance

We have linearity of variance for independent random variables:

$$\mathbb{V}\left[\sum_{i=1}^n X_i\right] = \sum_{i=1}^n \mathbb{V}[X_i].$$

Does not hold if the variables are dependent

(prove this: expand the definition of variance and rely upon  $\mathbb{E}(X_i X_j) = \mathbb{E}(X_i)\mathbb{E}(X_j)$  when  $X_i \perp X_j$ )

We have for any  $a \in \mathbb{R}$ ,

$$\mathbb{V}[aX] = a^2 \cdot \mathbb{V}[X].$$

1 Properties of expectation and variance

# Assignment Project Exam Help

2 Markov's inequality

3 Chebyshev's inequality

<https://powcoder.com>

4 Law of large numbers

Add WeChat powcoder

5 Wrapping Up

# Markov's Inequality

## Motivation

# Assignment Project Exam Help

1000 school students sit an examination

The busy principal is only told that the average score is 40 (out of 100).

The principal wants to estimate the maximum possible number of students who scored more than 80.

- A question about the minimum number of students is trivial to answer. Why?

# Markov's Inequality

## Motivation

Call  $x$  the number of students who score  $> 80$

Call  $S$  is the total score of students who score  $\leq 80$

We know:

$$40 \cdot 1000 - S = \{\text{total score of students who score above } 80\} > 80x$$

Exam scores are nonnegative, so certainly  $S \geq 0$

Thus,  $80x < 40 \cdot 1000$ , or

$$x < 500.$$

Can we formalise this more generally?

## Markov's Inequality

# Assignment Project Exam Help

Theorem

Let  $X$  be a nonnegative random variable. Then, for any  $\lambda > 0$ ,

$$P(X \geq \lambda) \leq \frac{E[X]}{\lambda}.$$

<https://powcoder.com>

Bounds probability of observing a large outcome

Add WeChat powcoder

Vacuous if  $\lambda < E[X]$

# Markov's Inequality

## Alternate Statement

Assignment Project Exam Help

### Corollary

Let  $X$  be a nonnegative random variable. Then, for any  $\lambda > 0$ ,

$$P(X \geq \lambda \cdot \mathbb{E}[X]) \leq \frac{\mathbb{E}[X]}{\lambda}.$$

Observations of nonnegative random variable unlikely to be much larger than expected value

Add WeChat powcoder

Vacuous if  $\lambda < 1$

# Markov's Inequality

## Proof

Assignment Project Exam Help

$$\mathbb{E}[X] = \sum_{x \in \mathcal{X}} x \cdot p(x)$$

$$= \sum_{x < \lambda} x \cdot p(x) + \sum_{x \geq \lambda} x \cdot p(x)$$

$$\geq \sum_{x \geq \lambda} x \cdot p(x) \quad \text{nonneg. of random variable}$$

$$\geq \sum_{x \geq \lambda} \lambda \cdot p(x)$$

$$= \lambda \cdot p(X \geq \lambda).$$

Add WeChat powcoder

# Markov's Inequality


Illustration from

<https://justindomke.wordpress.com/2008/06/19/markovs-inequality/>

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder





# Markov's Inequality


Illustration from

<https://justindomke.wordpress.com/2008/06/19/markovs-inequality/>

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



# Markov's Inequality


Illustration from

<https://justindomke.wordpress.com/2008/06/19/markovs-inequality/>

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



# Markov's Inequality


Illustration from

<https://justindomke.wordpress.com/2008/06/19/markovs-inequality/>

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



1 Properties of expectation and variance

# Assignment Project Exam Help

2 Markov's inequality

3 Chebyshev's inequality

<https://powcoder.com>

4 Law of large numbers

Add WeChat powcoder



5 Wrapping Up

# Chebyshev's Inequality

## Motivation

Markov's inequality only uses the **mean** of the distribution

**Assignment Project Exam Help**  
What about the spread of the distribution (**variance**)?



<https://powcoder.com>

Add WeChat powcoder

# Chebyshev's Inequality

## Assignment Project Exam Help

Let  $X$  be a random variable with  $\mathbb{E}[X] < \infty$ . Then, for any  $\lambda > 0$ ,

$$p(|X - \mathbb{E}[X]| \geq \lambda) \leq \frac{\mathbb{V}[X]}{\lambda^2}.$$

Bounds the probability of observing an “unexpected” outcome

Does not require non negativity

Two-sided bound

# Chebyshev's Inequality

## Alternate Statement

# Assignment Project Exam Help

### Corollary

Let  $X$  be a random variable with  $\mathbb{E}[X] < \infty$ . Then, for any  $\lambda > 0$ ,

$$p(|X - \mathbb{E}[X]| \geq \lambda \cdot \sqrt{\mathbb{V}[X]}) \leq \frac{1}{\lambda^2}.$$

Observations are unlikely to occur several standard deviations away from the mean

# Chebyshev's Inequality

Proof

Define

$$Y = (X - \mathbb{E}[X])^2.$$

Then, by Markov's inequality, for any  $\nu > 0$ ,

$$p(Y \geq \nu) \leq \frac{\mathbb{E}[Y]}{\nu}.$$

But,

$$\mathbb{E}[Y] = \mathbb{V}[X].$$

Also,

$$Y \geq \nu \iff |X - \mathbb{E}[X]| \geq \sqrt{\nu}.$$

Thus, setting  $\lambda = \sqrt{\nu}$ ,

$$p(|X - \mathbb{E}[X]| \geq \lambda) \leq \frac{\mathbb{V}[X]}{\lambda^2}.$$



# Chebyshev's Inequality

## Illustration

For a binomial  $X$  with  $N$  trials and success probability  $\theta$ , we have e.g.

$$P(|X - N\theta| \geq \sqrt{2N\theta(1-\theta)}) \leq \frac{1}{2}.$$

# Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



# Chebyshev's Inequality

## Example

Suppose we have a coin with bias  $\theta$ , i.e.  $p(X = 1) = \theta$

Say we flip the coin  $n$  times, and observe  $x_1, \dots, x_n \in \{0, 1\}$

We use the maximum likelihood estimator of  $\theta$ :

$$\hat{\theta}_n = \frac{x_1 + \dots + x_n}{n}$$

Estimate how large  $n$  should be such that

$$p(|\hat{\theta}_n - \theta| \geq 0.05) \leq 0.01?$$

1% probability of a 5% error

(Aside: the need for two parameters here is generic: “Probably Approximately Correct”)

# Chebyshev's Inequality

## Example

Observe that

$$\mathbb{E}[\hat{\theta}_n] = \frac{\sum_{i=1}^n \mathbb{E}[x_i]}{n} = \theta$$

$$\mathbb{V}[\hat{\theta}_n] = \frac{\sum_{i=1}^n \mathbb{V}[x_i]}{n^2} = \frac{\theta(1-\theta)}{n}.$$

Thus, applying Chebyshev's inequality to  $\hat{\theta}_n$ ,

$$P(|\hat{\theta}_n - \theta| > 0.05) \leq \frac{\theta(1-\theta)}{(0.05)^2 n}.$$

We are guaranteed this is less than 0.01 if

$$n \geq \frac{\theta(1-\theta)}{(0.05)^2(0.01)}.$$

When  $\theta = 0.5$ ,  $n \geq 10,000$  (!)

1 Properties of expectation and variance

# Assignment Project Exam Help

2 Markov's inequality

3 Chebyshev's inequality

<https://powcoder.com>

4 Law of large numbers

Add WeChat powcoder

5 Wrapping Up

## Independent and Identically Distributed

Let  $X_1, \dots, X_n$  be random variables such that:

- Each  $X_i$  is independent of  $X_j$
- The distribution of  $X_i$  is the same as that of  $X_j$

Then, we say that  $X_1, \dots, X_n$  are independent and identically distributed (or iid)

Example: For  $n$  independent flips of an unbiased coin,  $X_1, \dots, X_n$  are iid from Bernoulli( $\frac{1}{2}$ )

# Law of Large Numbers

## Theorem

Let  $X_1, \dots, X_n$  be a sequence of iid random variables, with

$$\mathbb{E}[X_i] = \mu$$

and  $\mathbb{V}[X_i] < \infty$ . Define

$$\bar{X}_n = \frac{X_1 + \dots + X_n}{n}.$$

Then, for any  $\epsilon > 0$ ,

$$\lim_{n \rightarrow \infty} p(|\bar{X}_n - \mu| > \epsilon) = 0.$$

Given enough trials, the empirical “success frequency” will be close to the expected value

# Law of Large Numbers

## Proof

Since the  $X_i$ 's are identically distributed,

$$\mathbb{E}[X_n] = \mu.$$

Since the  $X_i$ 's are independent,

$$\mathbb{V}[X_n] = \mathbb{V}\left[\frac{X_1 + \dots + X_n}{n}\right]$$

$$= \frac{\mathbb{V}[X_1 + \dots + X_n]}{n^2}$$

$$= \frac{n\sigma^2}{n^2}$$

$$= \frac{\sigma^2}{n}.$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

# Law of Large Numbers

## Proof

Applying Chebyshev's inequality to  $\bar{X}_n$

# Assignment Project Exam Help

$$p(|\bar{X}_n - \mu| \geq \epsilon) \leq \frac{\mathbb{V}[\bar{X}_n]}{\epsilon^2}$$

$$= \frac{\sigma^2}{n\epsilon^2}.$$

# <https://powcoder.com>

For any fixed  $\epsilon > 0$ , as  $n \rightarrow \infty$ , the right hand side  $\rightarrow 0$ .

# Add WeChat powcoder

Thus,


$$p(|\bar{X}_n - \mu| < \epsilon) \rightarrow 1.$$



# Law of Large Numbers

## Illustration

$N = 1000$  trials with Bernoulli random variable with parameter  $\frac{1}{2}$



# Law of Large Numbers


## Illustration

$N = 50000$  trials with Bernoulli random variable with parameter  $\frac{1}{2}$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



1 Properties of expectation and variance

# Assignment Project Exam Help

2 Markov's inequality

3 Chebyshev's inequality

<https://powcoder.com>

4 Law of large numbers

Add WeChat powcoder

5 Wrapping Up

# Assignment Project Exam Help

- Markov's inequality

<https://powcoder.com>

- Chebyshev's inequality

Add WeChat powcoder

- Law of large numbers

Next time

# Assignment Project Exam Help

- Ensembles and sequences

<https://powcoder.com>

- Typical sets

Add WeChat powcoder

- Approximation Equipartition (AEP)