

COMP2610 / COMP6261 - Information Theory

Lecture 4: Bayesian Inference

Assignment Project Exam Help

Robert C. Williamson

<https://powcoder.com>

Research School of Computer Science



Australian
National
University

Add WeChat powcoder

July 31, 2018

Assignment Project Exam Help

- Examples of joint, marginal and conditional distributions

- When can we say that X , Y do not influence each other?

- What, if anything, does $p(X = x, Y = y)$ tell us about $p(Y = y|X = x)$?

Review Exercise

Suppose we have binary random variables X, Y such that

$$p(X = 1) = 0.6$$

$$p(Y = 1|X = 0) = 0.7$$

$$p(Y = 1|X = 1) = 0.8$$

Assignment Project Exam Help

Then,

<https://powcoder.com>

$$p(X = 1|Y = 1) =$$

Add WeChat powcoder

Review Exercise

Suppose we have binary random variables X, Y such that

$$p(X = 1) = 0.6$$

$$p(Y = 1|X = 0) = 0.7$$

$$p(Y = 1|X = 1) = 0.8$$

Assignment Project Exam Help

Then,

<https://powcoder.com>

$$p(X = 1|Y = 1) = \frac{p(Y = 1|X = 1)p(X = 1)}{p(Y = 1)} \quad \text{Bayes' rule}$$

Add WeChat powcoder

Review Exercise

Suppose we have binary random variables X, Y such that

$$p(X = 1) = 0.6$$

$$p(Y = 1|X = 0) = 0.7$$

$$p(Y = 1|X = 1) = 0.8$$

Assignment Project Exam Help

Then,

<https://powcoder.com>

$$p(X = 1|Y = 1) = \frac{p(Y = 1|X = 1)p(X = 1)}{p(Y = 1)}$$

Bayes' rule

Add WeChat powcoder

$$= \frac{p(Y = 1|X = 1)p(X = 1)}{p(Y = 1|X = 1)p(X = 1) + p(Y = 1|X = 0)p(X = 0)}$$

Review Exercise

Suppose we have binary random variables X, Y such that

$$p(X = 1) = 0.6$$

$$p(Y = 1|X = 0) = 0.7$$

$$p(Y = 1|X = 1) = 0.8$$

Assignment Project Exam Help

Then,

<https://powcoder.com>

$$\begin{aligned} p(X = 1|Y = 1) &= \frac{p(Y = 1|X = 1)p(X = 1)}{p(Y = 1)} && \text{Bayes' rule} \\ &= \frac{p(Y = 1|X = 1)p(X = 1)}{p(Y = 1|X = 1)p(X = 1) + p(Y = 1|X = 0)p(X = 0)} \\ &= \frac{(0.8)(0.6)}{(0.8)(0.6) + (0.7)(0.4)} \\ &\approx 0.63 \end{aligned}$$

Add WeChat powcoder

This time

Assignment Project Exam Help

- More examples on Bayes' theorem:
 - ▶ Eating hamburgers

▶ <https://powcoder.com>

- ▶ Detecting terrorists

▶ Add WeChat powcoder

- Are there notions of probability beyond frequency counting?

Outline

1 Bayes' Rule: Examples

- Eating Hamburgers
- Detecting Terrorists
- The Monty Hall Problem

<https://powcoder.com>

2 Moments for functions of two discrete Random Variables

3 The meaning of Probability

4 Wrapping Up

Add WeChat powcoder

1 Bayes' Rule: Examples

- Eating Hamburgers
- Detecting Terrorists
- The Monty Hall Problem

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

2 Moment Generating Functions and Multivariate Normal Variables

3 The meaning of Probability

4 Wrapping Up

Bayesian Inference:

Example 1 (Barber, BRML, 2011)

Assignment Project Exam Help

- 90% of people with McD syndrome are frequent hamburger eaters

<https://powcoder.com>

Add WeChat powcoder

Bayesian Inference:

Example 1 (Barber, BRML, 2011)

Assignment Project Exam Help

- 90% of people with McD syndrome are frequent hamburger eaters

- <https://powcoder.com>
- Probability of someone having McD syndrome: $1/10000$

Add WeChat powcoder

Bayesian Inference:

Example 1 (Barber, BRML, 2011)

Assignment Project Exam Help

- 90% of people with McD syndrome are frequent hamburger eaters

- <https://powcoder.com>
Probability of someone having McD syndrome: $1/10000$

- Add WeChat powcoder
Proportion of hamburger eaters is about 50%

Bayesian Inference:

Example 1 (Barber, BRML, 2011)

Assignment Project Exam Help

- 90% of people with McD syndrome are frequent hamburger eaters

- <https://powcoder.com>
Probability of someone having McD syndrome: $1/10000$

- Add WeChat powcoder
Proportion of hamburger eaters is about 50%

What is the probability that a hamburger eater will have McD syndrome?

Bayesian Inference:

Example 1: Formalization

Assignment Project Exam Help

Let $McD \in \{0, 1\}$ be the variable denoting having the McD syndrome and $H \in \{0, 1\}$ be the variable denoting a hamburger eater. Therefore:

<https://powcoder.com>

Add WeChat powcoder

Bayesian Inference:

Example 1: Formalization

Assignment Project Exam Help

Let $McD \in \{0, 1\}$ be the variable denoting having the McD syndrome and $H \in \{0, 1\}$ be the variable denoting a hamburger eater. Therefore:

$$p(H = 1 | \neg McD = 1) = 9/10$$

<https://powcoder.com>

Add WeChat powcoder

Bayesian Inference:

Example 1: Formalization

Assignment Project Exam Help

Let $McD \in \{0, 1\}$ be the variable denoting having the McD syndrome and $H \in \{0, 1\}$ be the variable denoting a hamburger eater. Therefore:

$$p(H = 1 | McD = 1) = 9/10 \quad p(McD = 1) = 10^{-4}$$

Add WeChat powcoder

Bayesian Inference:

Example 1: Formalization

Assignment Project Exam Help

Let $McD \in \{0, 1\}$ be the variable denoting having the McD syndrome and $H \in \{0, 1\}$ be the variable denoting a hamburger eater. Therefore:

$$\begin{aligned} p(H = 1 | McD = 1) &= 9/10 & p(McD = 1) &= 10^{-4} \\ p(H = 1) &= 1/2 \end{aligned}$$

Add WeChat powcoder

Bayesian Inference:

Example 1: Formalization

Assignment Project Exam Help

Let $McD \in \{0, 1\}$ be the variable denoting having the McD syndrome and $H \in \{0, 1\}$ be the variable denoting a hamburger eater. Therefore:

$$\begin{aligned} p(H = 1 | McD = 1) &= 9/10 & p(McD = 1) &= 10^{-4} \\ p(H = 1) &= 1/2 \end{aligned}$$

We need to compute $p(McD = 1 | H = 1)$, the probability of a hamburger eater having McD syndrome.

Bayesian Inference:

Example 1: Formalization

Assignment Project Exam Help

Let $McD \in \{0, 1\}$ be the variable denoting having the McD syndrome and $H \in \{0, 1\}$ be the variable denoting a hamburger eater. Therefore:

$$\begin{aligned} p(H = 1 | McD = 1) &= 9/10 & p(McD = 1) &= 10^{-4} \\ p(H = 1) &= 1/2 \end{aligned}$$

We need to compute $p(McD = 1 | H = 1)$, the probability of a hamburger eater having McD syndrome.

Any ballpark estimates of this probability?

Bayesian Inference:

Example 1: Solution

Assignment Project Exam Help

$$p(McD = 1 | H = 1) = \frac{p(H = 1 | McD = 1)p(McD = 1)}{p(H = 1)}$$
$$= 1.8 \times 10^{-4}$$

Add WeChat powcoder

Bayesian Inference:

Example 1: Solution

Assignment Project Exam Help

$$p(McD = 1 | H = 1) = \frac{p(H = 1 | McD = 1)p(McD = 1)}{p(H = 1)}$$
$$= 1.8 \times 10^{-4}$$

Repeat the above computation if the proportion of hamburger eaters is rather small. (say in France) 0.001.

Example 2: Detecting Terrorists:

From understandinguncertainty.org

Assignment Project Exam Help

- Scanner detects true terrorists with 95% accuracy

<https://powcoder.com>

Add WeChat powcoder

Example 2: Detecting Terrorists:

From understandinguncertainty.org

Assignment Project Exam Help

- Scanner detects true terrorists with 95% accuracy

- Scanner detects upstanding citizens with 95% accuracy

<https://powcoder.com>

Add WeChat powcoder

Example 2: Detecting Terrorists:

From understandinguncertainty.org

Assignment Project Exam Help

- Scanner detects true terrorists with 95% accuracy

- Scanner detects upstanding citizens with 95% accuracy

- There is 1 terrorist on your plane with 100 passengers aboard

<https://powcoder.com>
Add WeChat powcoder

Example 2: Detecting Terrorists:

From understandinguncertainty.org

Assignment Project Exam Help

- Scanner detects true terrorists with 95% accuracy

- Scanner detects upstanding citizens with 95% accuracy

<https://powcoder.com>

- There is 1 terrorist on your plane with 100 passengers aboard

Add WeChat powcoder

- The shift looking man sitting next to you tests positive (terrorist)

Example 2: Detecting Terrorists:

From understandinguncertainty.org

Assignment Project Exam Help

- Scanner detects true terrorists with 95% accuracy

- Scanner detects upstanding citizens with 95% accuracy

<https://powcoder.com>

- There is 1 terrorist on your plane with 100 passengers aboard


Add WeChat powcoder

- The shiftily looking man sitting next to you tests positive (terrorist)

What are the chances of this man being a terrorist?

Example 2: Detecting Terrorists:

Simple Solution Using "Natural Frequencies" (David Spiegelhalter)




Add WeChat powcoder

Figure reproduced from users.ics.forth.gr/~imanolis/addinguncertainty.org

Example 2: Detecting Terrorists:

Simple Solution Using “Natural Frequencies” (David Spiegelhalter)




Add WeChat powcoder

The chances of the man being a terrorist are $\approx \frac{1}{6}$

Example 2: Detecting Terrorists:

Simple Solution Using “Natural Frequencies” (David Spiegelhalter)

Assignment Project Exam Help



Add WeChat powcoder

Figure reproduced from <https://www.powcoder.com>

The chances of the man being a terrorist are $\approx \frac{1}{6}$

- Relation to disease example
- Consequences when catching criminals

Example 2: Detecting Terrorists:

Formalization with Actual Probabilities

Assignment Project Exam Help

Let $T \in \{0, 1\}$ denote the variable regarding whether the person is a terrorist and $S \in \{0, 1\}$ denote the outcome of the scanner.

<https://powcoder.com>

Add WeChat powcoder

Example 2: Detecting Terrorists:

Formalization with Actual Probabilities

Assignment Project Exam Help

Let $T \in \{0, 1\}$ denote the variable regarding whether the person is a terrorist and $S \in \{0, 1\}$ denote the outcome of the scanner.

$$p(S=1|T=1)=0.95, \quad p(S=0|T=1)=0.05$$

Add WeChat powcoder

Example 2: Detecting Terrorists:

Formalization with Actual Probabilities

Assignment Project Exam Help

Let $T \in \{0, 1\}$ denote the variable regarding whether the person is a terrorist and $S \in \{0, 1\}$ denote the outcome of the scanner.

$$\begin{aligned} p(S = 1 | T = 1) &= 0.95 & p(S = 0 | T = 1) &= 0.05 \\ p(S = 0 | T = 0) &= 0.95 & p(S = 1 | T = 0) &= 0.05 \end{aligned}$$

Add WeChat powcoder

Example 2: Detecting Terrorists:

Formalization with Actual Probabilities

Assignment Project Exam Help

Let $T \in \{0, 1\}$ denote the variable regarding whether the person is a terrorist and $S \in \{0, 1\}$ denote the outcome of the scanner.

$$\begin{aligned}p(S = 1 | T = 1) &= 0.95 & p(S = 0 | T = 1) &= 0.05 \\p(S = 0 | T = 0) &= 0.95 & p(S = 1 | T = 0) &= 0.05 \\p(T = 1) &= 0.01 & p(T = 0) &= 0.99\end{aligned}$$

Add WeChat powcoder

Example 2: Detecting Terrorists:

Formalization with Actual Probabilities

Assignment Project Exam Help

Let $T \in \{0, 1\}$ denote the variable regarding whether the person is a terrorist and $S \in \{0, 1\}$ denote the outcome of the scanner.

$$\begin{aligned}p(S = 1 | T = 1) &= 0.95 & p(S = 0 | T = 1) &= 0.05 \\p(S = 0 | T = 0) &= 0.95 & p(S = 1 | T = 0) &= 0.05 \\p(T = 1) &= 0.01 & p(T = 0) &= 0.99\end{aligned}$$

We want to compute $p(T = 1 | S = 1)$, the probability of the man being a terrorist given that he has tested positive.

Example 2: Detecting Terrorists:

Solution with Bayes' Rule

Assignment Project Exam Help

$$p(T = 1|S = 1) = \frac{p(S = 1|T = 1)p(T = 1)}{p(S = 1|T = 1)p(T = 1) + p(S = 1|T = 0)p(T = 0)}$$

Add WeChat powcoder

Example 2: Detecting Terrorists:

Solution with Bayes' Rule

Assignment Project Exam Help

$$p(T = 1|S = 1) = \frac{p(S = 1|T = 1)p(T = 1)}{p(S = 1|T = 1)p(T = 1) + p(S = 1|T = 0)p(T = 0)}$$
$$= \frac{(0.95)(0.01)}{(0.95)(0.01) + (0.05)(0.99)}$$

Add WeChat powcoder

Example 2: Detecting Terrorists:

Solution with Bayes' Rule

Assignment Project Exam Help

$$p(T = 1|S = 1) = \frac{p(S = 1|T = 1)p(T = 1)}{p(S = 1|T = 1)p(T = 1) + p(S = 1|T = 0)p(T = 0)}$$
$$= \frac{(0.95)(0.01)}{(0.95)(0.01) + (0.05)(0.99)}$$

≈ 0.16

Add WeChat powcoder

Example 2: Detecting Terrorists:

Solution with Bayes' Rule

Assignment Project Exam Help

$$p(T = 1|S = 1) = \frac{p(S = 1|T = 1)p(T = 1)}{p(S = 1|T = 1)p(T = 1) + p(S = 1|T = 0)p(T = 0)}$$
$$= \frac{(0.95)(0.01)}{(0.95)(0.01) + (0.05)(0.99)}$$

Add WeChat powcoder

The probability of the man being a terrorist is $\approx \frac{1}{6}$

Example 2: Detecting Terrorists:

Posterior Versus Prior Belief

While the man has a low probability of being a terrorist, our belief has

increased compared to our prior

$$\frac{p(T = 1|S = 1)}{p(T = 1)} = \frac{0.16}{0.01} = 16$$

i.e. our belief in him being a terrorist has gone up by a factor of 16

Since terrorists are so rare, a factor of 16 does not result in a very high (absolute) probability or belief

(Aside: They are indeed very rare. For an intriguing (and surprising) example of the implications of inability to take account of actual base rates (in the example above we made the numbers up), and the effect on people's subsequent decisions, see Gerd Gigerenzer, Dread Risk, September 11, and Fatal Traffic Accidents, *Psychological Science* 15(4), 286–287, (2004); Gerd Gigerenzer, Out of the Frying Pan into the Fire: Behavioural Reactions to Terrorist Attacks, *Risk Analysis* 26(2), 347–351 (2006). His calculation (which of course is based on some assumptions) is that in the year following 9/11, 6 times the number of people who were killed as passengers *additionally* died on roads (that is the increase in road deaths due to people choosing to drive instead of flying)! He calls the reaction to very low probability events with a bad outcome “dread risk”.)

Example 3: The Monty Hall Problem

Problem Statement

- Three boxes, one with a prize and the other two are empty

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Example 3: The Monty Hall Problem

Problem Statement

- Three boxes, one with a prize and the other two are empty
- Each box has equal probability of having the prize

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Example 3: The Monty Hall Problem

Problem Statement

- Three boxes, one with a prize and the other two are empty
- Each box has equal probability of having the prize
- Your goal is to pick up the box with the prize in it

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Example 3: The Monty Hall Problem

Problem Statement

- Three boxes, one with a prize and the other two are empty
- Each box has equal probability of having the prize
- Your goal is to pick up the box with the prize in it
- You select one of the boxes

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Example 3: The Monty Hall Problem

Problem Statement

- Three boxes, one with a prize and the other two are empty
- Each box has equal probability of having the prize
- Your goal is to pick up the box with the prize in it
- You select one of the boxes
- The host, who knows the location of the prize, opens the empty box out of the other two boxes

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Example 3: The Monty Hall Problem

Problem Statement

- Three boxes, one with a prize and the other two are empty
- Each box has equal probability of having the prize
- Your goal is to pick up the box with the prize in it
- You select one of the boxes
- The host, who knows the location of the prize, opens the empty box out of the other two boxes

Should you switch to the other box? Would that increase your chances of winning the prize?

Example 3: The Monty Hall Problem:

Formalization

Let $C \in \{r, g, b\}$ denote the box that contains the prize where r, g, b refer to the identity of each box.

WLOG assume the following:

<https://powcoder.com>

Add WeChat powcoder

Example 3: The Monty Hall Problem:

Formalization

Let $C \in \{r, g, b\}$ denote the box that contains the prize where r, g, b refer to the identity of each box.

WLOG assume the following:

- You have selected box r

<https://powcoder.com>

Add WeChat powcoder

Example 3: The Monty Hall Problem:

Formalization

Let $C \in \{r, g, b\}$ denote the box that contains the prize where r, g, b refer to the identity of each box.

WLOG assume the following:

- You have selected box r
- Denote the event: “the host opens box b ” with $H=b$

<https://powcoder.com>

Add WeChat powcoder

Example 3: The Monty Hall Problem:

Formalization

Let $C \in \{r, g, b\}$ denote the box that contains the prize where r, g, b refer to the identity of each box.

WLOG assume the following:

- You have selected box r
- Denote the event: "the host opens box b " with $H=b$

$$P(C=r) = \frac{1}{3} \quad P(C=g) = \frac{1}{3} \quad P(C=b) = \frac{1}{3}$$

Example 3: The Monty Hall Problem:

Formalization

Let $C \in \{r, g, b\}$ denote the box that contains the prize where r, g, b refer to the identity of each box.

WLOG assume the following:

- You have selected box r
- Denote the event: "the host opens box b " with $H=b$

$$P(C=r) = \frac{1}{3} \quad P(C=g) = \frac{1}{3} \quad P(C=b) = \frac{1}{3}$$

$$p(H=b|C=r) = \frac{1}{2} \quad p(H=b|C=g) = 1 \quad p(H=b|C=b) = 0$$

Example 3: The Monty Hall Problem:

Formalization

Let $C \in \{r, g, b\}$ denote the box that contains the prize where r, g, b refer to the identity of each box.

WLOG assume the following:

- You have selected box r
- Denote the event: "the host opens box b " with $H=b$

$$p(C=r) = \frac{1}{3} \quad p(C=g) = \frac{1}{3} \quad p(C=b) = \frac{1}{3}$$

$$p(H=b|C=r) = \frac{1}{2} \quad p(H=b|C=g) = 1 \quad p(H=b|C=b) = 0$$

We want to compute $p(C=r|H=b)$ and $p(C=g|H=b)$ to decide if we should switch from our initial choice.

Example 3: The Monty Hall Problem:

Solution

We have that:

$$P(H = b) = \sum_{c \in \{r, g, b\}} P(H = b | C = c) P(C = c)$$

<https://powcoder.com>

Add WeChat powcoder

Example 3: The Monty Hall Problem:

Solution

We have that:

$$P(H = b) = \sum_{c \in \{r, g, b\}} P(H = b | C = c) P(C = c) \\ = (1/2)(1/3) + (1)(1/3) + (0)(1/3)$$

<https://powcoder.com>

Add WeChat powcoder

Example 3: The Monty Hall Problem:

Solution

We have that:

$$P(H=b) = \sum_{c \in \{r, g, b\}} P(H=b|C=c)P(C=c)$$

$$= (1/2)(1/3) + (1)(1/3) + (0)(1/3)$$

$$= 1/2$$

<https://powcoder.com>

Add WeChat powcoder

Example 3: The Monty Hall Problem:

Solution

We have that:

$$\begin{aligned}P(H = b) &= \sum_{c \in \{r, g, b\}} P(H = b | C = c) P(C = c) \\&= (1/2)(1/3) + (1)(1/3) + (0)(1/3) \\&= 1/2\end{aligned}$$

Therefore:

Add WeChat powcoder

Example 3: The Monty Hall Problem:

Solution

We have that:

$$\begin{aligned} p(H = b) &= \sum_{c \in \{r, g, b\}} p(H = b | C = c) p(C = c) \\ &= (1/2)(1/3) + (1)(1/3) + (0)(1/3) \\ &= 1/2 \end{aligned}$$

Therefore:

$$p(C = r | H = b) = \frac{p(H = b | C = r) p(C = r)}{p(H = b)}$$

Example 3: The Monty Hall Problem:

Solution

We have that:

$$\begin{aligned} p(H = b) &= \sum_{c \in \{r, g, b\}} p(H = b | C = c) p(C = c) \\ &= (1/2)(1/3) + (1)(1/3) + (0)(1/3) \\ &= 1/2 \end{aligned}$$

Therefore:

$$p(C = r | H = b) = \frac{p(H = b | C = r) p(C = r)}{p(H = b)} = \frac{(1/2)(1/3)}{(1/2)} = 1/3$$

Example 3: The Monty Hall Problem:

Solution

We have that:

$$\begin{aligned} p(H = b) &= \sum_{c \in \{r, g, b\}} p(H = b | C = c) p(C = c) \\ &= (1/2)(1/3) + (1)(1/3) + (0)(1/3) \\ &= 1/2 \end{aligned}$$

Therefore:

$$p(C = r | H = b) = \frac{p(H = b | C = r) p(C = r)}{p(H = b)} = \frac{(1/2)(1/3)}{(1/2)} = 1/3$$

Similarly, $p(C = g | H = b) = 2/3$.

Example 3: The Monty Hall Problem:

Solution

We have that:

$$\begin{aligned} p(H = b) &= \sum_{c \in \{r, g, b\}} p(H = b | C = c) p(C = c) \\ &= (1/2)(1/3) + (1)(1/3) + (0)(1/3) \\ &= 1/2 \end{aligned}$$

<https://powcoder.com>

Therefore:

$$p(C = r | H = b) = \frac{p(H = b | C = r) p(C = r)}{p(H = b)} = \frac{(1/2)(1/3)}{(1/2)} = 1/3$$

Similarly, $p(C = g | H = b) = 2/3$.

You should switch from your initial choice to the other box in order to increase your chances of winning the prize!

Example 3: The Monty Hall Problem:

Illustration of the Solution




Illustration of the solution when you have initially selected box r.

Example 3: The Monty Hall Problem:

Another Perspective

Assignment Project Exam Help

Switching is bad if, and only if, we initially picked the prize box (because if not, the other remaining box must contain the prize)

<https://powcoder.com>

We picked the prize box with probability $1/3$. This is independent of the host's action

Hence, with probability $2/3$ switching will reveal the prize box

Example 3: The Monty Hall Problem:

Variants to Ponder

Assignment Project Exam Help

Would switching be rational if

- The host only revealed a box when he knew we picked the right one?

<https://powcoder.com>

- The host only revealed a box when he knew we picked the wrong one?

Add WeChat powcoder

- The host is himself unaware of the prize box, and reveals a box at random, which by chance does not have the prize?

1 Bayes' Rule: Examples

- Eating Hamburgers
- Detecting heartbeats
- The Monty Hall Problem

Assignment Project Exam Help

2 Moments for functions of two discrete Random Variables

<https://powcoder.com>

3 The meaning of Probability

Add WeChat powcoder

4 Wrapping Up

The Expected Value of a Function of Two Discrete Random Variables

(Assuming you have met **Expectation** $E[X]$ and **Variance** $\text{Var}(X)$ before.)

The expected value of a function $g(X, Y)$ of two discrete random variables is defined as

$$E[g(X, Y)] = \sum_x \sum_y g(x, y) p(X=x, Y=y). \quad (1)$$

In particular, the expected value of X is given by

$$E[X] = \sum_x \sum_y xp(X=x, Y=y). \quad (2)$$

It should be noted that if we have already calculated the marginal distribution of X , then it is simpler to calculate $E[X]$ using this.

Covariance and the Correlation Coefficient

The covariance between X and Y , $\text{Cov}(X, Y)$ is given by

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y) \quad (3)$$

Note that by definition $\text{Cov}(X, X) = E(X^2) - E(X)^2 = \text{Var}(X)$.
The coefficient of correlation between X and Y is given by

$$r(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \quad (4)$$

Always in $[-1, 1]$.

Example

Discrete random variables X and Y have the following joint distribution:

| | $Y = 0$ | $Y = 1$ | $Y = 2$ |
|---------|---------------|---------------|---------------|
| $X = 0$ | $\frac{1}{3}$ | $\frac{1}{3}$ | $\frac{1}{3}$ |
| $X = 1$ | $\frac{1}{3}$ | $\frac{1}{3}$ | $\frac{1}{3}$ |

Calculate

- 1 $p(X > Y)$
- 2 marginal distributions of X and Y
- 3 expected values and variances of X and Y
- 4 coefficient of correlation between X and Y

Are X and Y independent?

Example

Assignment Project Exam Help

To calculate the probability of such an event, note that we sum over all the cells which correspond to that event. Hence,

$$\begin{aligned} p(X \geq Y) &= p(X = 0, Y = -1) + p(X = 1, Y = -1) \\ &\quad + p(X = 1, Y = 0) = \frac{1}{3} \end{aligned}$$

Add WeChat powcoder

Example

Recall that

$$p(X = x) = \sum_y p(X = x, Y = y).$$

Assignment Project Exam Help

Hence,

$$p(X = 0) = \sum_{y=-1}^1 p(X = 0, Y = y) = 0 + \frac{1}{3} + 0 = \frac{1}{3}$$

$$p(X = 1) = \sum_{y=-1}^1 p(X = 1, Y = y) = \frac{1}{3} + 0 + \frac{1}{3} = \frac{2}{3}$$

Note that after obtaining $p(X = 0)$, we could calculate $p(X = 1)$ by using the fact that

$$p(X = 1) = 1 - p(X = 0), \quad (5)$$

since X only takes the values 0 and 1.

Example

Similarly, Assignment Project Exam Help

$$p(Y = -1) = \sum_{x=-1}^1 p(X = x, Y = -1) = 0 + \frac{1}{3} = \frac{1}{3}$$

<https://powcoder.com>

$$p(Y = 0) = \sum_{x=0}^1 p(X = x, Y = 0) = \frac{1}{3} + 0 = \frac{1}{3}$$

Add WeChat powcoder

$$p(Y = 1) = 1 - p(Y = -1) - p(Y = 0) = \frac{1}{3}$$

Example

We then calculate the expected values and variances of X and Y from these marginal distributions.

$$E(X) = \sum_{x=0}^1 x p(X=x) = 0 \times \frac{1}{3} + 1 \times \frac{2}{3} = \frac{2}{3}$$

$$E(Y) = \sum_{y=-1}^1 y p(Y=y) = -1 \times \frac{1}{3} + 0 \times \frac{1}{3} + 1 \times \frac{1}{3} = 0$$

Example

To calculate the variances of X and Y , $\text{Var}(X)$ and $\text{Var}(Y)$, we use the formula

$$\text{Var}(X) = E(X^2) - (E(X))^2$$

$$E(X^2) = \sum_{x=0}^1 x^2 p(X=x) = 0^2 \times \frac{1}{3} + 1^2 \times \frac{2}{3} = \frac{2}{3}$$

<https://powcoder.com>

$$E(Y^2) = \sum_{y=-1}^1 y^2 p(Y=y) = (-1)^2 \times \frac{1}{3} + 0^2 \times \frac{1}{3} + 1^2 \times \frac{1}{3} = \frac{2}{3}.$$

Add WeChat powcoder

Thus we get

$$\text{Var}(X) = \frac{2}{3} - \left(\frac{2}{3}\right)^2 = \frac{2}{9}$$

$$\text{Var}(Y) = \frac{2}{3} - (0)^2 = \frac{2}{3}$$

Example

To calculate the correlation coefficient, we first calculate the covariance between X and Y . We have

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y).$$

where

$$\begin{aligned} E(XY) &= \sum_{x=-1}^1 \sum_{y=-1}^1 xy p(X=x, Y=y) \\ &= 0(-1)0 + 0(0)\frac{1}{3} + 0(1)0 + 1(-1)\frac{1}{3} + 1(0)0 + 1(1)\frac{1}{3} = 0 \end{aligned}$$

Thus we get

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y) = 0 - \frac{2}{3} \times 0 = 0.$$

From the definition of the correlation coefficient,

$$\rho(X, Y) = 0.$$

Example - is X and Y independent

Assignment Project Exam Help

We have that

$$p(X=0, Y=-1) = 0 \neq p(X=0)p(Y=-1) = \left(\frac{1}{3}\right)^2$$

Add WeChat powcoder

1 Bayes' Rule: Examples

- Eating Hamburgers
- Detecting Spam Emails
- The Monty Hall Problem

Assignment Project Exam Help

2 Moment Generating Functions and Multivariate Normal Variables

<https://powcoder.com>

3 The meaning of Probability

Add WeChat powcoder

4 Wrapping Up

The meaning of Probability

Frequentist : Frequencies of random repeatable experiments

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

The meaning of Probability

Frequentist : Frequencies of random repeatable experiments

- E.g. Prob. of biased coin landing "Heads"

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

The meaning of Probability

Frequentist : Frequencies of random repeatable experiments

- E.g. Prob. of biased coin landing "Heads"

Bayesian : Degrees of Belief

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

The meaning of Probability

Frequentist : Frequencies of random repeatable experiments

- E.g. Prob. of biased coin landing "Heads"

Bayesian : Degrees of Belief

- E.g. Prob. of Tasmanian Devil disappearing by the end of this decade

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

The meaning of Probability

Frequentist : Frequencies of random repeatable experiments

- E.g. Prob. of biased coin landing "Heads"

Bayesian : Degrees of Belief

- E.g. Prob. of Tasmanian Devil disappearing by the end of this decade

Cox Axioms

Given $B(x)$, $B(\bar{x})$, $B(x, y)$, $B(x|y)$, $B(y)$:

- 1 Degrees of belief can be ordered
- 2 $B(\bar{x}) = f[B(x)]$
- 3 $B(x, y) = g[B(x|y), B(y)]$

The meaning of Probability

Frequentist : Frequencies of random repeatable experiments

- E.g. Prob. of biased coin landing "Heads"

Bayesian : Degrees of Belief

- E.g. Prob. of Tasmanian Devil disappearing by the end of this decade

Cox Axioms


Given $B(x)$, $B(\bar{x})$, $B(x, y)$, $B(x|y)$, $B(y)$:

- 1 Degrees of belief can be ordered
- 2 $B(x) = f[B(x)]$
- 3 $B(x, y) = g[B(x|y), B(y)]$

If a set of Beliefs satisfy these axioms they can be mapped onto probabilities satisfying the rules of probability.

Assignment Project Exam Help

Add WeChat powcoder



Frequentists versus Bayesians: Round II

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder




Image from <http://normaldeviate.wordpress.com/2012/11/09/anti-xkcd/>

In practice one needs to make use of both interpretations. Wise to be open to both. This is a huge topic which we can not get into further here. Note that Mackay was firmly in the Bayesian camp. . .

1 Bayes' Rule: Examples

- Eating Hamburgers
- Detecting Spam Emails
- The Monty Hall Problem

Assignment Project Exam Help

2 Moment Generating Functions and Multivariate Normal Variables

<https://powcoder.com>

3 The meaning of Probability

Add WeChat powcoder

4 Wrapping Up

Summary

- Examples of application of Bayes' rule
 - Formalization

- ▶ Solution by applying Bayes' theorem

- Intuition is usually helpful although it may sometimes deceive us

- Frequentist v Bayesian probabilities

- Cox axioms

Next time

Assignment Project Exam Help

- Working through some useful probability distributions

<https://powcoder.com>

- More on Bayesian inference

Add WeChat powcoder