

COMP2610 / COMP6261 - Information Theory

Lecture 10: Typicality and Asymptotic Equipartition Property

Assignment Project Exam Help

Robert C. Williamson

<https://powcoder.com>

Research School of Computer Science



Australian
National
University

Add WeChat powcoder

21 August, 2018

Last time

Assignment Project Exam Help

Markov's inequality

Chebyshev's inequality

Law of large numbers

<https://powcoder.com>
Add WeChat powcoder

Law of Large Numbers

Theorem

Let X_1, \dots, X_n be a sequence of iid random variables, with

Assignment Project Exam Help

$$\mathbb{E}[X_i] = \mu$$

and $\mathbb{V}[X_i] < \infty$. Define

$$\bar{X}_n = \frac{X_1 + \dots + X_n}{n}$$

Then, for any $\beta > 0$,

$$\lim_{n \rightarrow \infty} P(|\bar{X}_n - \mu| < \beta) = 1.$$

This is also called $\bar{X}_n \rightarrow \mu$ in probability.

Add WeChat powcoder

Definition: For random variables v_1, v_2, \dots , we say $v_n \rightarrow v$ in probability if for all $\beta > 0$ $\lim_{n \rightarrow \infty} P(|v_n - v| > \beta) = 0$.

β is fixed (not shrinking like $\frac{1}{n}$). Not max/min. Reduction in variability.

This time

Assignment Project Exam Help

- Ensembles and sequences

<https://powcoder.com>

- Typical sets

Add WeChat powcoder

- Asymptotic Equipartition Property (AEP)

- 1 Ensembles and sequences
 - Counting Types of Sequences

Assignment Project Exam Help

- 2 Typical sets

<https://powcoder.com>

- 3 Asymptotic Equipartition Property (AEP)

Add WeChat powcoder

- 4 Wrapping Up

Assignment Project Exam Help

Ensemble

An **ensemble** X is a triple $(x, \mathcal{A}_X, \mathcal{P}_X)$; x is a **random variable** taking **values** in $\mathcal{A}_X = \{a_1, a_2, \dots, a_l\}$ with **probabilities** $\mathcal{P}_X = \{p_1, p_2, \dots, p_l\}$.

We will call \mathcal{A}_X the **alphabet** of the ensemble

Add WeChat powcoder

Ensembles

Example: Bent Coin

Assignment Project Exam Help

Let X be an **ensemble** with outcomes h for *heads* with probability 0.9 and t for *tails* with probability 0.1.



<https://powcoder.com>

- The **outcome set** is $\mathcal{A}_X = \{h, t\}$
- The **probabilities** are

$\mathcal{P}_X = \{p_h = 0.9, p_t = 0.1\}$

Add WeChat powcoder

Extended Ensembles

We can also consider **blocks** of outcomes, which will be useful to describe sequences:

Assignment Project Exam Help

Example (Coin Flips):

hhhhthhththh \rightarrow hh hh th ht ht hh (6 \times 2 outcome blocks)
 \rightarrow hhh hth hth hth (4 \times 3 outcome blocks)
 \rightarrow hhhh thht hthh (3 \times 4 outcome blocks)

Add WeChat powcoder

Extended Ensembles

We can also consider **blocks** of outcomes, which will be useful to describe sequences:

Assignment Project Exam Help

Example (Coin Flips):

hhhhthhththh \rightarrow hh hh th ht ht hh (6 \times 2 outcome blocks)
 \rightarrow hhh hth hth hth (4 \times 3 outcome blocks)
 \rightarrow hhhh thht hthh (3 \times 4 outcome blocks)

<https://powcoder.com>

Add WeChat powcoder

Extended Ensemble

Let X be a single ensemble. The **extended ensemble** of blocks of size N is denoted X^N . Outcomes from X^N are denoted $\mathbf{x} = (x_1, x_2, \dots, x_N)$. The **probability** of \mathbf{x} is defined to be $P(\mathbf{x}) = P(x_1)P(x_2) \dots P(x_N)$.

Extended Ensembles

Example: Bent Coin



Let X be an ensemble with outcomes
 $\mathcal{A}_X = \{h, t\}$ with $p_h = 0.9$ and $p_t = 0.1$

Consider X^4 – i.e., 4 flips of the coin.

$\mathcal{A}_{X^4} = \{hhhh, hhht, hthh, \dots, tttt\}$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Extended Ensembles

Example: Bent Coin



Let X be an ensemble with outcomes
 $\mathcal{A}_X = \{h, t\}$ with $p_h = 0.9$ and $p_t = 0.1$

Consider X^4 – i.e., 4 flips of the coin.

$\mathcal{A}_{X^4} = \{hhhh, hhht, hthh, \dots, tttt\}$

$$P(hhhh) = (0.9)^4 \approx 0.6561$$

$$P(tttt) = (0.1)^4 = 0.0001$$

$$P(hthh) = 0.9 \cdot 0.1 \cdot 0.9 \cdot 0.9 = (0.9)^3(0.1) \approx 0.0729$$

$$P(htht) = 0.9 \cdot 0.1 \cdot 0.9 \cdot 0.1 = (0.9)^2(0.1)^2 \approx 0.0081.$$

Extended Ensembles

Example: Bent Coin

Entropy of extended ensembles

Assignment Project Exam Help

We can view X^4 as comprising 4 independent random variables, based on the ensemble X

Entropy is additive for independent random variables

Thus,

$$H(X^4) = 4H(X) = 4 \cdot (-0.9 \log_2 0.9 - 0.1 \log_2 0.1) = 1.88 \text{ bits.}$$

More generally,

$$H(X^N) = NH(X).$$

Counting Types of Sequences

Criteria for dividing 2^N sequences into **types**

Assignment Project Exam Help

In the best coin example,

$$\begin{aligned}(0.9)^2(0.1)^2 &= P(\text{hhtt}) \\ &= P(\text{htht}) \\ &= P(\text{htth}) \\ &= P(\text{thht}) \\ &= P(\text{ttht}) \\ &= P(\text{tthh}).\end{aligned}$$

<https://powcoder.com>

Add WeChat powcoder

The **order** of outcomes in the sequence is **irrelevant**

Counting Types of Sequences

Let X be an ensemble with alphabet $\mathcal{A}_X = \{a_1, \dots, a_I\}$.

Assignment Project Exam Help

Let $p(X = a_i) = p_i$.

For a sequence $\mathbf{x} = x_1, x_2, \dots, x_N$, how to compute $p(\mathbf{x})$?

let $n_i = \#$ of times symbol a_i appears in \mathbf{x} (symbol count)

Given the n_i 's, we can compute the probability of seeing \mathbf{x} :

Add WeChat powcoder

$$\begin{aligned} P(\mathbf{x}) &= P(x_1) \cdot P(x_2) \cdot \dots \cdot P(x_N) \\ &= P(a_1)^{n_1} \cdot P(a_2)^{n_2} \cdot \dots \cdot P(a_I)^{n_I} \\ &= p_1^{n_1} \cdot p_2^{n_2} \cdot \dots \cdot p_I^{n_I} \end{aligned}$$

Sufficient statistics: $\{n_1, n_2, \dots, n_I\}$. Use it as a criteria of partitioning.

Counting Types of Sequences

Sequence Types

Each unique choice of (n_1, n_2, \dots, n_I) gives a different **type** of sequence

- 4 heads, (3 heads, 1 tail), (2 heads, 2 tails), ...
- Sequences in each type are equiprobable

For a given **type** of sequence how many sequences are there with these symbol counts?

$$\# \text{ of sequences with } n_i \text{ copies of } a_i = \frac{N!}{n_1! n_2! \dots n_I!}$$

Add WeChat powcoder

$$\begin{aligned} & \binom{N}{n_1} \binom{N-n_1}{n_2} \binom{N-n_1-n_2}{n_3} \dots \\ &= \frac{N!}{n_1!(N-n_1)!} \cdot \frac{(N-n_1)!}{n_2!(N-n_1-n_2)!} \cdot \frac{(N-n_1-n_2)!}{n_3!(N-n_1-n_2-n_3)!} \dots \end{aligned}$$

Counting Types of Sequences

Example

Probability of types Assignment Project Exam Help

Let $\mathcal{A} = \{a, b, c\}$ with $P(a) = 0.2$, $P(b) = 0.3$, $P(c) = 0.5$.

<https://powcoder.com>

Add WeChat powcoder

Counting Types of Sequences

Example

Probability of types Assignment Project Exam Help

Let $\mathcal{A} = \{a, b, c\}$ with $P(a) = 0.2$, $P(b) = 0.3$, $P(c) = 0.5$.

Each sequence of type $(n_a, n_b, n_c) = (2, 1, 3)$ has length 6 and probability $(0.2)^2(0.3)^1(0.5)^3 = 0.0045$.

Add WeChat powcoder

Counting Types of Sequences

Example

Probability of types Assignment Project Exam Help

Let $\mathcal{A} = \{a, b, c\}$ with $P(a) = 0.2$, $P(b) = 0.3$, $P(c) = 0.5$.

Each sequence of type $(n_a, n_b, n_c) = (2, 1, 3)$ has length 6 and probability $(0.2)^2(0.3)^1(0.5)^3 = 0.0045$.

There are $\frac{6!}{2!1!3!} = 60$ such sequences.

Add WeChat powcoder

Counting Types of Sequences

Example

Assignment Project Exam Help

Let $\mathcal{A} = \{a, b, c\}$ with $P(a) = 0.2$, $P(b) = 0.3$, $P(c) = 0.5$.

Each sequence of type $(n_a, n_b, n_c) = (2, 1, 3)$ has length 6 and probability $(0.2)^2(0.3)^1(0.5)^3 = 0.0015$.

There are $\frac{6!}{2!1!3!} = 60$ such sequences.

The probability \mathbf{x} is of type $(2, 1, 3)$ is $(0.0015) \cdot 60 = 0.09$.

Study probabilities at the level of types (most likely, average/typical)

1 Ensembles and sequences

• Counting Types of Sequences

Assignment Project Exam Help

2 Typical sets

<https://powcoder.com>

3 Asymptotic Equipartition Property (AEP)

Add WeChat powcoder

4 Wrapping Up

Extended Ensembles

Example

With $p_h = 0.75$, what are the probabilities for X^N ?

$$N = 2$$

\mathbf{x}	$P(\mathbf{x})$
hh	0.5625
ht	0.1875
th	0.1875
tt	0.0625

<https://powcoder.com>

Add WeChat powcoder

Extended Ensembles

Example

With $p_h = 0.75$, what are the probabilities for X^N ?

$N = 2$

\mathbf{x}	$P(\mathbf{x})$
hh	0.5625
ht	0.1875
th	0.1875
tt	0.0625

$N = 3$

\mathbf{x}	$P(\mathbf{x})$
hhh	0.4219
hht	0.1406
hth	0.1406
htt	0.0469
thh	0.1406
tht	0.0469
tth	0.0469
ttt	0.0156

<https://powcoder.com>
Add WeChat powcoder

Extended Ensembles

Example

With $p_h = 0.75$, what are the probabilities for X^N ?

$N = 2$

\mathbf{x}	$P(\mathbf{x})$
hh	0.5625
ht	0.1875
th	0.1875
tt	0.0625

$N = 3$

\mathbf{x}	$P(\mathbf{x})$
hhh	0.4219
hht	0.1406
hth	0.1406
htt	0.0469
thh	0.1406
tht	0.0469
tth	0.0469
ttt	0.0156

$N = 4$

\mathbf{x}	$P(\mathbf{x})$
hhhh	0.3164
hhht	0.1055
hhth	0.1055
hthh	0.1055
htth	0.0352
htth	0.0352
thhh	0.1055
thht	0.0352
thth	0.0352
thth	0.0352
thtt	0.0117
thtt	0.0117
ttth	0.0117
ttth	0.0117
ttth	0.0117
ttth	0.0117
tttt	0.0039

Assignment Project Exam Help

As N increases, there is an increasing spread of probabilities

The most likely single sequence will always be the all h's

However, for $N = 4$, the most likely sequence **type** is 3 h's and 1 t

Not surprising because $3 = N \cdot p_h$, pretty much average case.

<https://powcoder.com>

Add WeChat powcoder

Symbol Frequency in Long Sequences

To judge if a sequence is typical/average, a natural question to ask is:

How often does each symbol appear in a sequence \mathbf{x} from \mathcal{X}^N ?

Assignment Project Exam Help

Intuitively, in a sequence of length N , let a_i appear for n_i times.

Then **in expectation**

$n_i \approx N \cdot P(a_i)$
<https://powcoder.com>

Note $p_i = P(a_i)$, and

$$P(\mathbf{x}) = P(a_1)^{n_1} P(a_2)^{n_2} \dots P(a_l)^{n_l} \approx p_1^{Np_1} p_2^{Np_2} \dots p_l^{Np_l}$$

Add WeChat powcoder

Symbol Frequency in Long Sequences

To judge if a sequence is typical/average, a natural question to ask is:

How often does each symbol appear in a sequence \mathbf{x} from \mathcal{X}^N ?

Assignment Project Exam Help

Intuitively, in a sequence of length N , let a_i appear for n_i times.

Then **in expectation**

$n_i \approx N \cdot P(a_i)$
<https://powcoder.com>

Note $p_i = P(a_i)$, and

$P(\mathbf{x}) = P(a_1)^{n_1} P(a_2)^{n_2} \dots P(a_I)^{n_I} \approx p_1^{Np_1} p_2^{Np_2} \dots p_I^{Np_I}$
Add WeChat powcoder

So the *information content* $-\log_2 P(\mathbf{x})$ of that sequence is approximately

$$-p_1 N \log_2 p_1 - \dots - p_I N \log_2 p_I = -N \sum_{i=1}^I p_i \log_2 p_i = NH(X)$$

Typical Sets

We want to consider elements \mathbf{x} that have $-\log_2 P(\mathbf{x})$ "close" to $NH(X)$

Typical Set

For "closeness" $\beta > 0$ the typical set $T_{N\beta}$ for X^N is

$$T_{N\beta} \stackrel{\text{def}}{=} \{\mathbf{x} : |-\log_2 P(\mathbf{x}) - NH(X)| < N\beta\}$$

$$= \left\{ \mathbf{x} : \left| -\frac{1}{N} \log_2 P(\mathbf{x}) - H(X) \right| < \beta \right\}$$

Union of types

Typical Sets

We want to consider elements \mathbf{x} that have $-\log_2 P(\mathbf{x})$ “close” to $NH(X)$

Typical Set

For “closeness” $\beta > 0$, the typical set $T_{N\beta}$ for X^N is

$$T_{N\beta} \stackrel{\text{def}}{=} \{\mathbf{x} : |-\log_2 P(\mathbf{x}) - NH(X)| < N\beta\}$$

$$= \left\{ \mathbf{x} : \left| -\frac{1}{N} \log_2 P(\mathbf{x}) - H(X) \right| < \beta \right\}$$

Union of types



What when $\beta = 0$ (and replace $<$ by \leq)?

Criterion based on information content. Other criterion (KL divergence)?

Typical Sets

Properties

Assignment Project Exam Help

Typical sequences are nearly equiprobable. Every $\mathbf{x} \in T_{N\beta}$ has

$$2^{-N(H(X)+\beta)} \leq P(\mathbf{x}) \leq 2^{-N(H(X)-\beta)}.$$

Variation is small when β is small

Number of sequences in the typical set: For any N, β ,

Add WeChat powcoder

$$|T_{N\beta}| \leq 2^{N(H(X)+\beta)}.$$

Typical Sets

Proof of Cardinality Bound

For every $\mathbf{x} \in T_{N\beta}$,

$$p(\mathbf{x}) \geq 2^{-N(H(X)+\beta)}.$$

Thus,

$$\begin{aligned} 1 &= \sum_{\mathbf{x} \in T_{N\beta}} p(\mathbf{x}) \\ &\geq \sum_{\mathbf{x} \in T_{N\beta}} 2^{-N(H(X)+\beta)} \\ &= 2^{-N(H(X)+\beta)} \cdot |T_{N\beta}|. \end{aligned}$$

Thus

$$|T_{N\beta}| \leq 2^{N(H(X)+\beta)}$$

Typical Sets

Most Likely Sequence

The most likely sequence may not belong to the typical set

Assignment Project Exam Help

e.g. with $p_h = 0.75$, we have

$$-\frac{1}{4} \log_2 P(\text{hhhh}) = 0.4150$$

<https://powcoder.com>

whereas $H(X) = 0.8113$

Add WeChat powcoder

The most likely single sequence \rightarrow hhhh


The most likely single sequence type $\rightarrow \{\text{hhht}, \text{hthh}, \dots\}$

Typical Sets

Most Likely Sequence

Probability of most likely sequence decays like $(p_h)^N$ ($p_h = 0.75$)

Sequences with $N \cdot p_h$ heads contain much more total probability mass



Blue curve corresponds to typical set with $\beta = 0$. What if $\beta > 0$?

1 Ensembles and sequences

• Counting Types of Sequences

Assignment Project Exam Help

2 Typical sets

<https://powcoder.com>

3 Asymptotic Equipartition Property (AEP)

Add WeChat powcoder

4 Wrapping Up

Asymptotic Equipartition Property

Eventually
Informally

Equally Divided



Asymptotic Equipartition Property (Informal)

As $N \rightarrow \infty$, $\log_2 P(x_1, \dots, x_N)$ is close to $-NH(X)$ with high probability.

For large block sizes “almost all sequences are typical” (i.e., in $T_{N\beta}$)

<https://powcoder.com>

$$n(r)P(\mathbf{x}) = \binom{N}{r} p_1^r (1 - p_1)^{N-r}$$



Probability sequence \mathbf{x} has r heads for $N = 100$ (left) and $N = 1000$ (right). Here $P(X = \text{head}) = 0.1$.

Asymptotic Equipartition Property

Formally

Asymptotic Equipartition Property

If x_1, x_2, \dots are i.i.d. with distribution P then, in probability

$$-\frac{1}{N} \log_2 P(x_1, \dots, x_N) \rightarrow H(X).$$

In precise language:

$$(\forall \beta > 0) \lim_{N \rightarrow \infty} p \left(\left| -\frac{1}{N} \log_2 P(x_1, \dots, x_N) - H(X) \right| < \beta \right) = 1.$$

Exactly the probability of $\mathbf{x} \in \Gamma_{N\beta}$.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Asymptotic Equipartition Property

Formally

Asymptotic Equipartition Property

If x_1, x_2, \dots are i.i.d. with distribution P then, in probability,

$$-\frac{1}{N} \log_2 P(x_1, \dots, x_N) \rightarrow H(X).$$

In precise language:

$$(\forall \beta > 0) \lim_{N \rightarrow \infty} P \left(\left| -\frac{1}{N} \log_2 P(x_1, \dots, x_N) - H(X) \right| < \beta \right) = 1.$$

Exactly the probability of $\mathbf{x} \in \Gamma_{N\beta}$.

Recall definition: for random variables v_1, v_2, \dots , we say $v_N \rightarrow v$ in **probability** if for all $\beta > 0$ $\lim_{N \rightarrow \infty} P(|v_N - v| > \beta) = 0$

Here v_N corresponds to $-\frac{1}{N} \log_2 P(x_1, \dots, x_N)$.

Asymptotic Equipartition Property

Comments

Why is it surprising/significant? Assignment Project Exam Help

For an ensemble with binary outcomes, and low entropy,

$$|T_N| \leq e^{NH(X) + \beta} \ll 2^N$$

<https://powcoder.com>

i.e. the typical set is a **small fraction** of all possible sequences

AEP says that for N sufficiently large, we are virtually guaranteed to draw a sequence from this small set

Add WeChat powcoder

Significance in information theory

Asymptotic Equipartition Property

Proof

Since x_1, \dots, x_N are independent,

$$\begin{aligned} -\frac{1}{N} \log p(x_1, \dots, x_N) &= -\frac{1}{N} \log \prod_{n=1}^N p(x_n) \\ &= -\frac{1}{N} \sum_{n=1}^N \log p(x_n) \end{aligned}$$

Let $Y = -\log p(X)$ and $y_n = -\log p(x_n)$. Then, $y_n \sim Y$, and

$$\mathbb{E}[Y] = H(X).$$

But then by the law of large numbers,

$$(\forall \beta > 0) \lim_{N \rightarrow \infty} p \left(\left| \frac{1}{N} \sum_{n=1}^N y_n - H(X) \right| > \beta \right) = 0.$$

1 Ensembles and sequences

• Counting Types of Sequences

Assignment Project Exam Help

2 Typical sets

<https://powcoder.com>

3 Asymptotic Equipartition Property (AEP)

Add WeChat powcoder

4 Wrapping Up

Assignment Project Exam Help

- Ensembles and sequences

- Typical sets

- Asymptotic Equipartition Property (AEP)

<https://powcoder.com>

Add WeChat powcoder

Next: Source Coding.