Week 2

Assignment Project Exam Help

Properties of Numbers II Udaya Parampalli

https://powcoder.com

University of Melbourne



Part -2 Symmetric key Cryptography

https://powcoder.com

Properties of Numbers

Add WeChat powcoder

Workshop 2: Workshops start from this week

Quizz 2



- 2.1 More on Inverse Modulo n
- 2.2 https://powcoder.com
- 2.3 How can you use Euler's Phi to compute inverses?

2.1 https://powcoder.com

Modular Arithmetic

Let a and b be integers and let n be a positive integer. We say "a" is congruent to "b", modulo n and write

Assignment Project Exam Help

if a and b differ by a multiple of n; i.e.; if n is a factor of |b-a|. Every integer is congruent mod n to exactly one of the integers in the sentence of t

We can define the following operations:

Add WeChat, powcoder

$$x \otimes_n y = (xy) \mod n$$

When the context is clear we use the above special addition and multiplication symbols interchangeably with their counterpart regular symbols.

Modular Multiplicative Inverse

Assingament Project Exam Help

Let $x \in Z_n$, if there is an integer y such that

https://powcoder.com
then we say y is the multiplicative inverse of x. It is denoted by

then we say y is the multiplicative inverse of x. It is denoted by $y = x^{-1}$ usually.

Example General in 20W General 2 is inverse of 3 modulo 5.

Determining multiplicative inverse

Assignment Project Exam Help

For any integers a and b, there exist integers x and y such that

https://poweoder.com

2.2 https://powcoder.com

Euler Phi function

Definition

Two numbers a and b are relatively prime if gcd(a, b) is 1.

ssignment Project Exam Help

Euler phi function(or Euler totient function): For $n \ge 1$, let $\phi(n)$ denote the number of integers less than n but are relatively prime to n. **Proof** 100 **Proof** 100

Definition

Reduced set of residues, Remain as set of residues, Remain as set of residues modulo nowhich are relatively prime to n.

Example:
$$\phi(6) = 2$$
: Observe, $gcd(1,6) = 1, gcd(2,6) = 2, gcd(3,6) = 3, gcd(4,6) = 2, gcd(5,6) = 1$. Then $R(6) = \{1,5\}$. Hence $\phi(6) = 2$.

Some Relations

Fact

 $\phi(p) = p - 1$, for any prime p.

Assignment Project Exam Help

 $\underset{\text{for any prime}}{\text{https}} \overset{\phi(p^a)}{\underset{\text{and any prime}}{\text{powcoder.com}}} \overset{p^a-p^{a-1}}{\underset{\text{integer}}{\text{powcoder.com}}} \overset{p^{a-1}(p-1)}{\underset{\text{integer}}{\text{der.com}}}$

Consider numbers from 0 to p^a-1 , then only numbers which have some divisivity p^a are those numbers which are multiple of p. There are exactly p^a are those numbers which are multiple of p. There are exactly p^a such numbers including the number 0. All other numbers are relatively prime to p^a . Hence, $\phi(p^a)=p^a-p^{a-1}=p^{a-1}(p-1)$ as needed. Example: $\phi(8)=4$, the numbers which are multiple of 2 are $\{2,4,6,8\}$ and hence the relatively prime numbers are all odd numbers up to 7, i.e $R(8)=\{1,3,5,7\}$.

Some Relations, cont.

Fact

 $\phi(pq) = (p-1)(q-1)$, for any pair of primes p and q.

Stignine at icker no perce but stix no infeur to visualize. Again consider numbers from 1 to pq. Like before, we can exclude all those numbers which are multiple of p and q to form R(pq). Then dan we say the following? POWCOGET.COM

|R(pq)| = pq - ((pq)/q) - ((pq)/p) = (pq - p - q)

In the above counting, we have excluded multiple of pq twice, once while claim the wutters of the design will be the multiples of q. So we need to make the following change

$$\phi(pq) = |R(pq)| = pq - p - q + 1 = (p-1)(q-1).$$

Example: $\phi(15) = 8$, the relatively prime numbers are 1, 2, 4, 7, 8, 11, 13, 14.

Euler Phi function is multiplicative

Fact

Assignment $\Pr_{\phi(ab)}^{\text{If a and b are relatively prime numbers (gcd(a, b) = 1), then, }} \Pr_{\phi(ab)} \Pr_{\phi(b)} \text{Exam Help}$

This is not directly obvious with whatever we have studied so far.
But the this ear fact you we could be supported in the studied so far.

But the this ear fact you we could be supported in the support of the studied so far.

Using the above fact, we can derive a general result about eulers ϕ function. We know that any number has a unique factorization:

Add WeChat powcoder
$$n = \prod_{i=1}^{n} p_i^{a_i} = p_1^{a_1} p_2^{a_2} \cdots p_{\tau}^{a_{\tau}}$$
,

where τ is a positive number, p_i are primes and $a_i \ge 1$ and Π is the symbol for product. Find $\phi(n)$ for this case. Example: What is $\phi(200) = \phi(2^3 5^2)$?.

$$\phi(n) = \phi(\Pi_{i=1}^{\tau} p_i^{a_i}) = \phi(p_1^{a_1} p_2^{a_2} \cdots p_{\tau}^{a_{\tau}}),$$
 From the point of t

$$\phi(n) = \prod_{i=1}^{\tau} p_i^{a_i-1}(p_i-1)).$$

Example dia is we chan power ener

2.3 Inteps: ys/ publishicoder.com

A system of the first of the state of the st

We will prove the following result later, but let us state it now. let \mathbf{Z}_n^{\star} be set of numbers from 1 to n-1 but are relatively prime.

Theorem

If $a \in \mathbf{Z}_n^{\star}$, then $a^{\phi(n)} = 1 \pmod{n}$.

Now, how de you we the above the orp for Wn Gutil Cler's e of a mod n?

https://powcoder.com

```
Function(a, n)

inva := a^{\phi(n)-1} \pmod{n}.

Return with the function;

We Chat powcoder
```

Part -2 Symmetric key Cryptography

https://powcoder.com

Properties of Numbers

Add WeChat powcoder

Workshop 2: Workshops start from this week

Quizz 2

