

Assignment Project Exam Help

Karush-Kuhn-Tucker conditions

<https://powcoder.com>

Geoff Gordon & Ryan Tibshirani
Optimization 10-725 / 36-725

Add WeChat powcoder

Remember duality

Given a minimization problem

$$\begin{aligned} \min_{x \in \mathbb{R}^n} \quad & f(x) \\ \text{subject to} \quad & h_i(x) \leq 0, \quad i = 1, \dots, m \\ & \ell_j(x) = 0, \quad j = 1, \dots, r \end{aligned}$$

we defined the Lagrangian

$$L(x, u, v) = f(x) + \sum_{i=1}^m u_i h_i(x) + \sum_{j=1}^r v_j \ell_j(x)$$

Add WeChat powcoder
and **Lagrange dual function:**

$$g(u, v) = \min_{x \in \mathbb{R}^n} L(x, u, v)$$

The subsequent **dual problem** is:

$$\max_{u \in \mathbb{R}^m, v \in \mathbb{R}^r} g(u, v)$$

subject to $u \geq 0$

Assignment Project Exam Help

Important properties:

- Dual problem is always convex, i.e., g is always concave (even if primal problem is not convex)
- The primal and dual optimal values, f^* and g^* , always satisfy weak duality: $f^* \geq g^*$
- Slater's condition: for convex primal, if there is an x such that

$$h_1(x) < 0, \dots, h_m(x) < 0 \quad \text{and} \quad \ell_1(x) = 0, \dots, \ell_r(x) = 0$$

then **strong duality** holds: $f^* = g^*$. (Can be further refined to strict inequalities over nonaffine h_i , $i = 1, \dots, m$)

Duality gap

Given primal feasible x and dual feasible u, v , the quantity

Assignment Project Exam Help
is called the **duality gap** between x and u, v . Note that

$$f(x) - f^* \leq f(x) - g(u, v)$$

https://powcoder.com

so if the duality gap is zero, then x is primal optimal (and similarly, u, v are dual optimal)

Add WeChat powcoder
From an algorithmic viewpoint, provides a stopping criterion: if $f(x) - g(u, v) \leq \epsilon$, then we are guaranteed that $f(x) - f^* \leq \epsilon$

Very useful, especially in conjunction with iterative methods ...
more dual uses in coming lectures

Dual norms

Let $\|x\|$ be a **norm**, e.g.,

- ℓ_p norm: $\|x\|_p = (\sum_{i=1}^n |x_i|^p)^{1/p}$, for $p \geq 1$
- Nuclear norm: $\|X\|_{\text{nuc}} = \sum_{i=1}^r \sigma_i(X)$

Assignment Project Exam Help

We define its **dual norm** $\|x\|_*$ as

$$\|x\|_* = \max_{\|z\| \leq 1} z^T x$$

<https://powcoder.com>

Gives us the inequality $|z^T x| \leq \|z\| \|x\|_*$, like Cauchy-Schwartz.

Back to our examples

- ℓ_p norm dual: $(\|x\|_p)_* = \|x\|_q$, where $1/p + 1/q = 1$
- Nuclear norm dual: $(\|X\|_{\text{nuc}})_* = \|X\|_{\text{spec}} = \sigma_{\max}(X)$

Dual norm of dual norm: it turns out that $\|x\|_{**} = \|x\|$...
connections to duality (including this one) in coming lectures

Assignment Project Exam Help

Today.

- KKT conditions
- Examples
- Constrained and Lagrange forms
- Uniqueness with 1-norm penalties

<https://powcoder.com>

Add WeChat powcoder

Karush-Kuhn-Tucker conditions

Given general problem

Assignment Project Exam Help

$$\begin{aligned} \min_{x \in \mathbb{R}^n} & f(x) \\ \text{subject to} & h_i(x) \leq 0, \quad i = 1, \dots, m \\ & \ell_j(x) = 0, \quad j = 1, \dots, r \end{aligned}$$

<https://powcoder.com>

The **Karush-Kuhn-Tucker conditions** or **KKT conditions** are:

- $0 \in \partial f(x) + \sum_{i=1}^m u_i \partial h_i(x) + \sum_{j=1}^r v_j \partial \ell_j(x)$ (stationarity)
- $u_i \cdot h_i(x) = 0$ for all i (complementary slackness)
- $h_i(x) \leq 0, \ell_j(x) = 0$ for all i, j (primal feasibility)
- $u_i \geq 0$ for all i (dual feasibility)

Add WeChat powcoder

Necessity

Let x^* and u^*, v^* be primal and dual solutions with zero duality gap (strong duality holds, e.g. under Slater's condition). Then

$$f(x^*) = g(u^*, v^*)$$

$$= \min_{x \in \mathbb{R}^n} f(x) + \sum_{i=1}^m u_i^* h_i(x) + \sum_{j=1}^r v_j^* \ell_j(x)$$

$$\begin{aligned} &\leq f(x^*) + \sum_{i=1}^m u_i^* h_i(x^*) + \sum_{j=1}^r v_j^* \ell_j(x^*) \\ &\leq f(x^*) \end{aligned}$$

In other words, all these inequalities are actually equalities

Two things to learn from this:

- The point x^* minimizes $L(x, u^*, v^*)$ over $x \in \mathbb{R}^n$. Hence the subdifferential of $L(x, u^*, v^*)$ must contain 0 at $x = x^*$ —this is exactly the **stationarity** condition.
- We must have $\sum_{i=1}^m u_i^* h_i(x^*) = 0$, and since each term here is ≤ 0 , this implies $u_i^* h_i(x^*) = 0$ for every i —this is exactly **complementary slackness**.

<https://powcoder.com>

Primal and dual feasibility obviously hold. Hence, we've shown:

If x^* and u^*, v^* are primal and dual solutions, with zero duality gap, then x^*, u^*, v^* satisfy the KKT conditions.

(Note that this statement assumes nothing a priori about convexity of our problem, i.e. of f, h_i, ℓ_j)

Sufficiency

If there exists x^*, u^*, v^* that satisfy the KKT conditions, then

$$\begin{aligned} g(u^*, v^*) &= f(x^*) + \sum_{i=1}^m u_i^* h_i(x^*) + \sum_{j=1}^r v_j^* \ell_j(x^*) \\ &= f(x^*) \end{aligned}$$

where the first equality holds from stationarity, and the second holds from complementary slackness

Therefore duality gap is zero (and x^* and u^*, v^* are primal and dual feasible) so x^* and u^*, v^* are primal and dual optimal. I.e., we've shown:

If x^* and u^*, v^* satisfy the KKT conditions, then x^* and u^*, v^* are primal and dual solutions

Putting it together

In summary, KKT conditions:

- always sufficient
- necessary under strong duality

Assignment Project Exam Help

Putting it together:

For a problem with strong duality (e.g. assume Slater's condition: convex problem and there exists x strictly satisfying non-affine inequality constraints),

x^* and u^*, v^* are primal and dual solutions
 $\Leftrightarrow x^*$ and u^*, v^* satisfy the KKT conditions

(Warning, concerning the stationarity condition: for a differentiable function f , we cannot use $\partial f(x) = \{\nabla f(x)\}$ unless f is convex)

What's in a name?

Older folks will know these as the KT (Kuhn-Tucker) conditions:

- First appeared in publication by Kuhn and Tucker in 1951
- Later people found out that Karush had the conditions in his unpublished master's thesis of 1939

Many people (including instructor!) use the term KKT conditions for unconstrained problems i.e., to refer to stationarity condition

Note that we could have alternatively derived the KKT conditions from studying optimality entirely via subgradients

$$0 \in \partial f(x^*) + \sum_{i=1}^m \mathcal{N}_{\{h_i \leq 0\}}(x^*) + \sum_{j=1}^r \mathcal{N}_{\{\ell_j = 0\}}(x^*)$$

where recall $\mathcal{N}_C(x)$ is the normal cone of C at x

Quadratic with equality constraints

Consider for $Q \succeq 0$,

$$\begin{aligned} \min_{x \in \mathbb{R}^n} \quad & \frac{1}{2} x^T Q x + c^T x \\ \text{subject to} \quad & Ax = 0 \end{aligned}$$

E.g., as in Newton step for $\min_{x \in \mathbb{R}^n} f(x)$ subject to $Ax = b$

Convex problem, no inequality constraints, so by KKT conditions:
 x is a solution if and only if

$$\begin{bmatrix} Q & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} x \\ u \end{bmatrix} = \begin{bmatrix} -c \\ 0 \end{bmatrix}$$

for some u . Linear system combines stationarity, primal feasibility
(complementary slackness and dual feasibility are vacuous)

Water-filling

Example from B & V page 245: consider problem

$$\min_{x \in \mathbb{R}^n} - \sum_{i=1}^n \log(\alpha_i + x_i)$$

subject to $x \geq 0, 1^T x = 1$

Information theory: think of $\log(\alpha_i + x_i)$ as communication rate of i th channel. KKT conditions:

$$\begin{aligned} -1/(\alpha_i + x_i) - u_i + v &= 0, \quad i = 1, \dots, n \\ x_i(x_i + \alpha_i) &= 0, \quad i = 1, \dots, n, \quad x \geq 0, \quad 1^T x = 1, \quad v \geq 0 \end{aligned}$$

Eliminate u :

$$\begin{aligned} 1/(\alpha_i + x_i) &\leq v, \quad i = 1, \dots, n \\ x_i(v - 1/(\alpha_i + x_i)) &= 0, \quad i = 1, \dots, n, \quad x \geq 0, \quad 1^T x = 1 \end{aligned}$$

Can argue directly stationarity and complementary slackness imply

$$x_i = \begin{cases} 1/v - \alpha & \text{if } v \leq 1/\alpha \\ 0 & \text{if } v > 1/\alpha \end{cases} = \max\{0, 1/v - \alpha\}, \quad i = 1, \dots, n$$

Assignment Project Exam Help

Still need x to be feasible, i.e., $1^T x = 1$, and this gives

$$\sum_{i=1}^n \max\{0, 1/v - \alpha_i\} = 1$$


<https://powcoder.com>

Univariate equation, piecewise linear in $1/v$ and not hard to solve

Add WeChat powcoder

This reduced problem is called **water-filling**

(From B & V page 246)



Lasso

Let's return the lasso problem: given response $y \in \mathbb{R}^n$, predictors $A \in \mathbb{R}^{n \times p}$ (columns A_1, \dots, A_p), solve

Assignment Project Exam Help

$$\min_{x \in \mathbb{R}^p} \frac{1}{2} \|y - Ax\|^2 + \lambda \|x\|_1$$

KKT conditions:

$$A^T(y - Ax) = \lambda s$$

where $s \in \partial \|x\|_1$, i.e.,

$$s_i \in \begin{cases} \{1\} & \text{if } x_i > 0 \\ \{-1\} & \text{if } x_i < 0 \\ [-1, 1] & \text{if } x_i = 0 \end{cases}$$

Now we read off important fact: if $|A_i^T(y - Ax)| < \lambda$, then $x_i = 0$
... we'll return to this problem shortly


Group lasso

Suppose predictors $A = [A_{(1)} \ A_{(2)} \ \dots \ A_{(G)}]$, split up into groups, with each $A_{(i)} \in \mathbb{R}^{n \times p(i)}$. If we want to select entire groups rather than individual predictors, then we solve the **group lasso** problem:

$$\min_{x=(x_{(1)}, \dots, x_{(G)}) \in \mathbb{R}^p} \frac{1}{2} \|y - Ax\|^2 + \lambda \sum_{i=1}^G \sqrt{p(i)} \|x_{(i)}\|_2$$

<https://powcoder.com>

Add WeChat powcoder



(From Yuan and Lin (2006), "Model selection and estimation in regression with grouped variables")

KKT conditions:

$$A_{(i)}^T(y - Ax) = \lambda \sqrt{p(i)} s_{(i)}, \quad i = 1, \dots, G$$

where each $s_{(i)} \in \partial \|x_{(i)}\|_2$, i.e.,

$$s_{(i)} \in \begin{cases} \{x_{(i)} / \|x_{(i)}\|_2\} & \text{if } x_{(i)} \neq 0 \\ \{z \in \mathbb{R}^{p(i)} : \|z\|_2 \leq 1\} & \text{if } x_{(i)} = 0 \end{cases}, \quad i = 1, \dots, G$$

Hence if $\|A_{(i)}^T(y - Ax)\|_2 < \lambda \sqrt{p(i)}$, then $x_{(i)} = 0$. On the other hand, if $x_{(i)} \neq 0$, then

$$x_{(i)} = \left(A_{(i)}^T A_{(i)} + \frac{\lambda \sqrt{p(i)}}{\|x_{(i)}\|_2} I \right)^{-1} A_{(i)}^T r_{-(i)},$$

$$\text{where } r_{-(i)} = y - \sum_{j \neq i} A_{(j)} x_{(j)}$$

Constrained and Lagrange forms

Often in statistics and machine learning we'll switch back and forth between **constrained** form, where $t \in \mathbb{R}$ is a tuning parameter,

Assignment Project Exam Help

$$\min_{x \in \mathbb{R}^n} f(x) \text{ subject to } t(x) \leq t \quad (C)$$

and **Lagrange** form, where $\lambda \geq 0$ is a tuning parameter,

<https://powcoder.com>

$$\min_{x \in \mathbb{R}^n} f(x) + \lambda \cdot h(x) \quad (L)$$

and claim these are equivalent. Is this true (assuming convex f, h)?

Add WeChat powcoder

(C) to (L): if problem (C) is strictly feasible then strong duality holds, and there exists some $\lambda \geq 0$ (dual solution) such that any solution x^* in (C) minimizes

$$f(x) + \lambda \cdot (f(x) - t)$$

so x^* is also a solution in (L)

(L) to (C): if x^* is a solution in (L), then the KKT conditions for (C) are satisfied by taking $t = h(x^*)$, so x^* is a solution in (C)

Conclusion:

$$\bigcup_{\lambda \geq 0} \{\text{solutions in (L)}\} = \bigcup_t \{\text{solutions in (C)}\}$$

$$\bigcup_{\lambda \geq 0} \{\text{solutions in (L)}\} \supseteq \bigcup_{\substack{t \text{ such that (C) \\ is strictly feasible}}} \{\text{solutions in (C)}\}$$

Strictly speaking this is not a perfect equivalence (albeit minor nonequivalence). Note: when the only value of t that leads to a feasible but not strictly feasible constraint set is $t = 0$, i.e.,

$$\{x : g(x) \leq t\} \neq \emptyset, \{x : g(x) = t\} = \emptyset \Rightarrow t = 0$$

(e.g., this is true if g is a norm) then we do get perfect equivalence

Uniqueness in 1-norm penalized problems

Using the KKT conditions and simple probability arguments, we can produce the following (perhaps surprising) result:

Theorem: Let f be differentiable and strictly convex. $A \in \mathbb{R}^{n \times p}$, $\lambda > 0$. Consider

$$\min_{x \in \mathbb{R}^p} f(Ax) + \lambda \|x\|_1$$

If the entries of A are drawn from a continuous probability distribution (on \mathbb{R}^{np}), then with probability 1 the solution $x^* \in \mathbb{R}^p$ is unique and has at most $\min\{n, p\}$ nonzero components

Remark: here f must be strictly convex, but no restrictions on the dimensions of A (we could have $p \gg n$)

Proof: the KKT conditions are

$$-A^T \nabla f(Ax) = \lambda s, \quad s_i \in \begin{cases} \{\text{sign}(x_i)\} & \text{if } x_i \neq 0 \\ [-1, 1] & \text{if } x_i = 0 \end{cases}, \quad i = 1, \dots, n$$

Note that Ax, s are unique. Define $S = \{j : |A_j^T \nabla f(Ax)| = \lambda\}$, also unique, and note that any solution satisfies $x_i = 0$ for all $i \notin S$

First assume that $\text{rank}(A_S) < |S|$ (here $A \in \mathbb{R}^{n \times |S|}$, submatrix of A corresponding to columns in S). Then for some $i \in S$,

$$A_i = \sum_{j \in S \setminus \{i\}} c_j A_j$$

for constants $c_j \in \mathbb{R}$, hence

$$s_i A_i = \sum_{j \in S \setminus \{i\}} (s_i s_j c_j) \cdot (s_j A_j)$$


Taking an inner product with $-\nabla f(Ax)$,

$$\lambda = \sum_{j \in S \setminus \{i\}} (s_i s_j c_j) \lambda, \quad \text{i.e.,} \quad \sum_{j \in S \setminus \{i\}} s_i s_j c_j = 1$$

In other words, we've proved that $\text{rank}(A_S) < |S|$ implies $s_i A_i$ is in the affine span of $s_j A_j$, $j \in S \setminus \{i\}$ (subspace of dimension $< n$)

We say that the matrix A has columns in general position if any affine subspace E of dimension $k < n$ does not contain more than $k + 1$ elements; of $\{\pm A_1, \dots, \pm A_p\}$ (excluding antipodal pairs)

It is straightforward to show that, if the entries of A have a density over \mathbb{R}^{nA} , then A is in general position with probability 1



Therefore, if entries of A are drawn from continuous probability distribution, any solution must satisfy $\text{rank}(A_S) = |S|$

Assignment Project Exam Help

Recalling the KKT conditions, this means the number of nonzero components in any solution is $\leq |S| \leq \min\{n, p\}$

Furthermore, we can reduce our optimization problem (by partially solving) to

$$\min_{x_S \in \mathbb{R}^{|S|}} f(A_S x_S) + \lambda \|x_S\|_1$$

Finally, strict convexity implies uniqueness of the solution in this problem, and hence in our original problem \square

Add WeChat powcoder

Back to duality

One of the most important uses of duality is that, under strong duality, we can **characterize primal solutions** from dual solutions

Recall that under strong duality, the KKT conditions are necessary for optimality. Given dual solutions u^*, v^* , any primal solution x^* satisfies the stationarity condition

$$0 \in \partial f(x^*) + \sum_{i=1}^m u_i^* \partial h_i(x^*) + \sum_{j=1}^r v_j^* \partial \ell_j(x^*)$$

In other words, x^* achieves the minimum in $\min_{x \in \mathbb{R}^n} L(x; u^*, v^*)$

- Generally, this reveals a characterization of primal solutions
- In particular, if this is satisfied uniquely (i.e., above problem has a unique minimizer), then the corresponding point must be the primal solution

Assignment Project Exam Help

- S. Boyd and L. Vandenberghe (2004), *Convex Optimization*, Cambridge University Press, Chapter 5
- R. T. Rockafellar (1970), *Convex Analysis*, Princeton University Press, Chapters 28–30

<https://powcoder.com>

Add WeChat powcoder