

---

# Assignment 0: Exercise 1


```
clear all; close all
```

a) Rosenbrock function

```
f = @(x,y) 100.*(y - x.^2).^2 + (1 - x).^2;
```

b) Discretisation of the unit square. It can be scaled with factor  $\alpha$  using simple multiplication  $\alpha X, \alpha Y$   
Plot the function of two variables

```
n = 300;  
x = linspace(-1,1,n+1);  
h = 2/n;  
y = x;  
[X,Y] = meshgrid(x,y);  
  
alpha = 2;  
%figure, surf(alpha*X, alpha*Y, f(alpha*X,alpha*Y), 'EdgeColor',  
%figure, surf(alpha*X, alpha*Y, f(alpha*X,alpha*Y), 'EdgeColor',  
%figure, contour(alpha*X, alpha*Y, log(f(alpha*X,alpha*Y)))  
figure, contourf(alpha*X, alpha*Y, log(f(alpha*X,alpha*Y)))
```



c) Gradient

$$df/dx = -400(y - x^2)x - 2(1 - x)$$

$$df/dy = 200(y - x^2)$$

c) Hessian

$$d^2f/dx^2 = -400(y - 3x^2) + 2$$

$$d^2f/dxdy = -400x$$

$$d^2f/dydx = -400x$$

$$d^2f/dy^2 = 200$$

d) Find the minimiser  $x^*$  of the function  $f$ . Show that  $x^*$  is unique and that  $\nabla^2 f(x^*)$  is positive definite.

We observe that  $f(x, y) \geq 0$  because is the sum of two non-negative terms. Also,

$$f(x, y) = 0 \Leftrightarrow y = x^2 \text{ and } x = 1$$

This is only possible if  $x = 1$  and  $y = 1$ . Hence, we have  $x^* = (1, 1)^T$ .

The Hessian at  $x^*$  is

$$\nabla^2 f(x^*) = \begin{pmatrix} 802 & -400 \\ -400 & 200 \end{pmatrix}$$

Hence, to verify that it is positive definite, for all  $(x, y) \neq 0$  we must have:

$$(x \ y) \begin{pmatrix} 802 & -400 \\ -400 & 200 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} > 0$$

This is equivalent to

$$802x^2 - 800xy + 200y^2 > 0$$

Finally, we have

$$2x^2 + 200(2x - y)^2 > 0,$$

that obviously holds for all  $(x, y) \neq (0, 0)$ .

---

e) Finite difference implementation of the gradient

```
% Construct central finite difference to be applied columnwise
D = 1/(2*h)*spdiags([-ones(n+1,1) ones(n+1,1)], [-1 1], n+1,n+1);
D(1,1:2) = 1/h*[-1 1];
D(n+1,n:n+1) = 1/h*[-1 1];

% Gradient via finite differences
% d/dy
Dyf = D*f(alpha*X, alpha*Y);
% d/dx, central finite difference applied rowwise
Dxf = (D*(f(alpha*X, alpha*Y).')).';

% Hessian via finite difference
% d^2/dxdy
Dxyf = D*Dxf;
% d^2/dx^2
Dxxf = (D*(Dxf.')).';
% d^2/dy^2
Dyyf = D*Dyf;
% d^2/dydx
Dyxf = (D*(Dyf.')).';
```

Assignment Project Exam Help

*Published with MATLAB® R2016a*

<https://powcoder.com>

Add WeChat powcoder