

EM & Factor Analysis

- Introduce general EM Algorithm
- Gaussian mixture as EM
- Factor Analysis

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

RECAP

WE EXAMINED GMMs (light sources)

GIVEN $x^{(1)} \dots x^{(n)} \in \mathbb{R}^d$ AND $K > 0$

DO: find $P(z^{(i)} = j)$ for $i=1 \dots n, j=1 \dots K$

WE CALLED $z^{(i)}$ A LATENT VARIABLE (NOT OBSERVED DIRECTLY)

TODAY: Derive Gmm Algo in more general way

Keep: E-STEP, M-STEP

Guess $z^{(i)}$ \leftarrow \rightarrow fit parameters of model

Assignment Project Exam Help


<https://powcoder.com>

Add WeChat powcoder

LATENT VARIABLE model

$$\begin{aligned}
 \ell(\theta) &= \sum_{i=1}^n \log P(x^{(i)} | \theta) \quad \begin{matrix} \text{DATA} \\ \text{PARAMETERS} \end{matrix} \\
 &= \sum_{i=1}^n \sum_z \log P(x^{(i)} | z, \theta) P(z | \theta) \quad \begin{matrix} \text{latent variable} \end{matrix}
 \end{aligned}$$

Picture of our algorithm (cf w/ Gmm)



our estimate at

Add WeChat powcoder

HOPE: $L_t(\theta)$ EASIER to optimize than $\ell(\theta)$

(E-STEP) find $L_t(\theta)$ given $\theta^{(t)}$ Why does this Abstract Gmm?

(M-STEP) $\theta^{(t+1)} = \text{Argmax}_{\theta} L_t(\theta)$

Next: How do we find $L_t(\theta)$ given $\theta^{(t)}$

IDEA WE GO TEAM-BY-TEAM $\log P(x^{(i)}; \theta)$ Single team

$$\log P(x^{(i)}; \theta) = \log \sum_z P(x^{(i)}, z^{(i)} = z; \theta)$$

let $Q^{(i)}(z)$ s.t. $\sum_z Q^{(i)}(z) = 1, Q^{(i)}(z) \geq 0$ (*)

Symbol pushing!
Just def of \mathbb{E}

$$= \log \sum_z \frac{Q^{(i)}(z)}{Q^{(i)}(z)} P(x^{(i)}, z^{(i)})$$

$$= \log \mathbb{E} \left[\frac{P(x^{(i)}, z^{(i)})}{Q^{(i)}(z)} \right]$$

Assignment Project Exam Help

RECALL JENSEN, WE CAN SAY $\log(\mathbb{E}[X]) \geq \mathbb{E}[\log(X)]$

(SIDEBAR, next page)
<https://powcoder.com>

Add WeChat powcoder

$$\geq \mathbb{E} \left[\log \frac{P(x^{(i)}, z^{(i)})}{Q^{(i)}(z)} \right]$$

$$= \sum_z Q^{(i)}(z) \log \frac{P(x^{(i)}, z^{(i)})}{Q^{(i)}(z)}$$

for ANY $Q^{(i)}$ ABOVE THIS HOLDS, AND FOR EACH TEAM

SO PICK ANY $Q^{(i)} \rightarrow$ GIVES AN $\mathcal{L}_i(\theta)$!

CALL THIS $\text{ELBO}(x, Q; \theta) = \sum_z Q(z) \log \frac{P(x, z; \theta)}{Q(z)}$

SHOWS $P(x^{(i)}; \theta) \geq \text{ELBO}(x, Q^{(i)}; \theta)$

Is Property 1

Property 2

WE PICK A SPECIFIC $Q^{(i)}$ depending on $x^{(i)} \neq \theta^{(i)}$

SO THAT $l(\theta^{(i)}) = L_t(\theta^{(i)})$

Goal: Pick $Q^{(i)}$ s.t. \rightarrow

$$\log \sum_z P(x^{(i)}, z^{(i)}) = \sum_z Q^{(i)}(z) \log \frac{P(x^{(i)}, z^{(i)}; \theta^{(i)})}{Q^{(i)}(z)}$$

$$\text{SET } Q^{(i)}(z) = P(z | x^{(i)}; \theta^{(i)})$$

\Rightarrow

$$\frac{P(x^{(i)}, z^{(i)})}{Q^{(i)}(z)} = \frac{P(z^{(i)} | x^{(i)}; \theta^{(i)}) P(x^{(i)}; \theta^{(i)})}{P(z^{(i)} | x^{(i)}; \theta^{(i)})}$$

\rightarrow DOES NOT DEPEND ON z

Assignment Project Exam Help

Hence $\sum_z Q^{(i)}(z) \cdot \log L = \log L$ [RHS/ELBO]

(LHS) $\log \sum_z P(x^{(i)}, z^{(i)}) = \log \sum_z Q^{(i)}(z) \frac{P(x^{(i)}, z^{(i)})}{Q^{(i)}(z)}$

$$= \log L$$

RESTATE EM

(E-STEP) for $i=1 \dots n$, SET $Q_i(z) = P(z | x^{(i)}; \theta^{(i)})$

(M-STEP) $\theta^{(i+1)} = \underset{\theta}{\text{ARGMAX}} L_t(\theta)$


$$= \underset{\theta}{\text{Argmax}} \sum_{i=1}^n \text{ELBO}(x^{(i)}, \theta^{(i)}, \theta)$$

SENSEU REMINDER (SIDE BAR)

$$\log(\mathbb{E}[x]) \geq \mathbb{E}[\log(x)]$$

\log is concave function

TO help you remember inequality, thought experiment $x \rightarrow \begin{matrix} a & \text{w/ prob } \lambda \\ b & \text{w/ prob } 1-\lambda \end{matrix}$



SINCE z is like $\mathbb{E}[x] = \lambda a + (1-\lambda)b$

Picture says $f(\mathbb{E}[z]) \geq \mathbb{E}[f(z)]$ for $f = \log$.

for any $z = \lambda a + (1-\lambda)b$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder