

156 Let n and r be natural variables in the refinement

$P \Leftarrow \text{if } n=1 \text{ then } r:=0 \text{ else } n:=\text{div } n \ 2. \ P. \ r:=r+1 \text{ fi}$

Suppose the operations div and $+$ each take time 1 and all else is free (even the call is free). Insert appropriate time increments, and find an appropriate P to express the execution time in terms of

(a) the initial values of the memory variables. Prove the refinement for your choice of P .

§ With time increments added, I must prove

$P \Leftarrow \text{if } n=1 \text{ then } r:=0 \text{ else } t:=t+1. \ n:=\text{div } n \ 2. \ P. \ t:=t+1. \ r:=r+1 \text{ fi}$

How should we choose P ? Execution of P proceeds as follows. If n is initially 0, then n is divided by 2, making it again 0, and we are in an infinite loop. If n is initially positive, then it is repeatedly divided by 2 (rounding down) until it becomes 1, then r is assigned 0, then r is incremented as many times as n was divided by 2. The number of times n is divided by 2 until it becomes 1 is the logarithm (base 2) of n . This may not be obvious, so I can easily code this procedure in any implemented programming language I like, and run it for a variety of initial values for n and r and for initial time 0, and see that the final value of t is $2 \times \text{floor}(\log n)$. So P can be

$(n=0 \Rightarrow t'=\infty) \wedge (n>0 \Rightarrow t' = t + 2 \times \text{floor}(\log n))$

But floor is an awkward function to work with, so I'll get rid of it by replacing the exact time with an upper bound. My choice of P is

$(n=0 \Rightarrow t'=\infty) \wedge (n>0 \Rightarrow t' \leq t + 2 \times \log n)$

I prove it in parts (each conjunct separately), and I prove each part by cases.

First part, first case:

$$\begin{aligned}
 & (n=0 \Rightarrow t'=\infty) \Leftarrow n=1 \wedge (r:=0) \quad \text{portation} \\
 = & n=0 \wedge n=1 \wedge (r:=0) \Rightarrow t'=\infty \\
 = & \perp \wedge (r:=0) \Rightarrow t'=\infty \\
 = & \perp \Rightarrow t'=\infty \\
 = & \top
 \end{aligned}$$

First part, last case:

$$\begin{aligned}
 & (n=0 \Rightarrow t'=\infty) \Leftarrow n \neq 1 \wedge (t:=t+1. \ n:=\text{div } n \ 2. \ n=0 \Rightarrow t'=\infty. \ t:=t+1. \ r:=r+1) \\
 & \quad \text{portation and expand final assignment} \\
 = & n=0 \wedge n \neq 1 \wedge (t:=t+1. \ n:=\text{div } n \ 2. \ n=0 \Rightarrow t'=\infty. \ t:=t+1. \ r'=r+1 \wedge n'=n \wedge t'=t) \\
 \Rightarrow & t'=\infty \quad \text{simplify, and substitution law in two parts} \\
 = & n=0 \wedge (\text{div } n \ 2 = 0 \Rightarrow t'=\infty. \ r'=r+1 \wedge n'=n \wedge t'=t+1) \\
 \Rightarrow & t'=\infty \quad \text{eliminate dependent composition} \\
 = & n=0 \wedge (\exists r'', n'', t''. (\text{div } n \ 2 = 0 \Rightarrow t''=\infty) \wedge r'=r''+1 \wedge n'=n'' \wedge t'=t''+1) \\
 \Rightarrow & t'=\infty \quad \text{context: } n=0 \\
 = & n=0 \wedge (\exists r'', n'', t''. t''=\infty \wedge r'=r''+1 \wedge n'=n'' \wedge t'=t''+1) \Rightarrow t'=\infty \quad \text{one-point} \\
 = & n=0 \wedge t'=\infty+1 \Rightarrow t'=\infty \quad \text{absorption and specialization} \\
 = & \top
 \end{aligned}$$

Last part, first case:

$$\begin{aligned}
 & (n>0 \Rightarrow t' \leq t + 2 \times \log n) \Leftarrow n=1 \wedge (r:=0) \quad \text{portation and expand assignment} \\
 = & n=1 \wedge r'=0 \wedge n'=n \wedge t'=t \Rightarrow t' \leq t + 2 \times \log n \quad \text{context, and } \log 1 = 0 \\
 = & \top
 \end{aligned}$$

Last part, last case:

$$\begin{aligned}
& (n > 0 \Rightarrow t' \leq t + 2 \times \log n) \\
\Leftarrow & \quad n \neq 1 \wedge (t := t+1. \ n := \text{div } n \ 2. \ n > 0 \Rightarrow t' \leq t + 2 \times \log n. \ t := t+1. \ r := r+1) \\
& \text{portation and expand final assignment} \\
= & \quad t' \leq t + 2 \times \log n \\
\Leftarrow & \quad n > 1 \wedge (t := t+1. \ n := \text{div } n \ 2. \ n > 0 \Rightarrow t' \leq t + 2 \times \log n. \ t := t+1. \ r' = r+1 \wedge n' = n \wedge t' = t) \\
& \text{substitution law in two parts} \\
= & \quad t' \leq t + 2 \times \log n \\
\Leftarrow & \quad n > 1 \wedge (\text{div } n \ 2 > 0 \Rightarrow t' \leq t+1 + 2 \times \log(\text{div } n \ 2). \ r' = r+1 \wedge n' = n \wedge t' = t+1) \\
& \text{eliminate dependent composition} \\
= & \quad t' \leq t + 2 \times \log n \\
\Leftarrow & \quad n > 1 \wedge (\exists r'', n'', t''. \ (\text{div } n \ 2 > 0 \Rightarrow t'' \leq t+1 + 2 \times \log(\text{div } n \ 2)) \\
& \quad \wedge r' = r''+1 \wedge n' = n'' \wedge t' = t''+1) \quad \text{one-point for } n'' \text{ and } t'' \\
= & \quad t' \leq t + 2 \times \log n \\
\Leftarrow & \quad n > 1 \wedge (\exists r''. \ (\text{div } n \ 2 > 0 \Rightarrow t' \leq t+2 + 2 \times \log(\text{div } n \ 2)) \wedge r' = r''+1) \quad \text{distributive} \\
= & \quad t' \leq t + 2 \times \log n \\
\Leftarrow & \quad n > 1 \wedge (\exists r'': \text{nat} \cdot r' = r''+1) \wedge (\text{div } n \ 2 > 0 \Rightarrow t' \leq t+2 + 2 \times \log(\text{div } n \ 2)) \\
& \text{in preparation for one-point, rewrite } r' = r''+1 \text{ and make } r'': \text{nat} \text{ an explicit conjunct} \\
= & \quad t' \leq t + 2 \times \log n \\
\Leftarrow & \quad n > 1 \wedge (\exists r'': \text{nat} \cdot r'' = r'-1 \wedge (r'': \text{nat})) \wedge (\text{div } n \ 2 > 0 \Rightarrow t' \leq t+2 + 2 \times \log(\text{div } n \ 2)) \\
& \text{now use one-point} \\
= & \quad t' \leq t + 2 \times \log n \\
\Leftarrow & \quad n > 1 \wedge (r'-1: \text{nat}) \wedge (\text{div } n \ 2 > 0 \Rightarrow t' \leq t+2 + 2 \times \log(\text{div } n \ 2)) \\
= & \quad t' \leq t + 2 \times \log n \Leftarrow n > 1 \wedge (r'-1: \text{nat}) \wedge (\text{div } n \ 2 > 0 \Rightarrow t' \leq t + 2 + 2 \times \log(\text{div } n \ 2)) \\
& \text{simplify } \text{div } n \ 2 > 0 \\
= & \quad t' \leq t + 2 \times \log n \Leftarrow n > 1 \wedge r' \geq 1 \wedge (n > 1 \Rightarrow t' \leq t + 2 + 2 \times \log(\text{div } n \ 2)) \\
& \text{increase } \text{div } n \ 2 \text{ to } n/2 \\
& \text{this will increase } t + 2 + 2 \times \log(\text{div } n \ 2) \\
& \text{this will weaken } t' \leq t + 2 + 2 \times \log(\text{div } n \ 2) \\
& \text{this will weaken } n > 1 \Rightarrow t' \leq t + 2 + 2 \times \log(\text{div } n \ 2) \\
& \text{this will weaken } n > 1 \wedge r' \geq 1 \wedge (n > 1 \Rightarrow t' \leq t + 2 + 2 \times \log(\text{div } n \ 2)) \\
& \text{this will strengthen } t' \leq t + 2 \times \log n \Leftarrow n > 1 \wedge r' \geq 1 \wedge (n > 1 \Rightarrow t' \leq t + 2 + 2 \times \log(\text{div } n \ 2)) \\
& \text{so we need to put } \Leftarrow \text{ in the left margin} \\
\Leftarrow & \quad t' \leq t + 2 \times \log n \Leftarrow n > 1 \wedge r' \geq 1 \wedge (n > 1 \Rightarrow t' \leq t + 2 + 2 \times \log(n/2)) \quad \text{drop } r' \geq 1 \\
& \text{this will weaken } n > 1 \wedge r' \geq 1 \wedge (n > 1 \Rightarrow t' \leq t + 2 + 2 \times \log(n/2)) \\
& \text{this will strengthen } t' \leq t + 2 \times \log n \Leftarrow n > 1 \wedge r' \geq 1 \wedge (n > 1 \Rightarrow t' \leq t + 2 + 2 \times \log(n/2)) \\
& \text{so again we need to put } \Leftarrow \text{ in the left margin} \\
\Leftarrow & \quad t' \leq t + 2 \times \log n \Leftarrow n > 1 \wedge (n > 1 \Rightarrow t' \leq t + 2 + 2 \times \log(n/2)) \quad \text{discharge and simplify} \\
= & \quad t' \leq t + 2 \times \log n \Leftarrow n > 1 \wedge t' \leq t + 2 \times \log n \quad \text{specialization} \\
= & \quad \top
\end{aligned}$$

(b) the final values of the memory variables. Prove the refinement for your choice of P .

§ I prove

$$\begin{aligned}
t' = t + 2 \times r' & \Leftarrow \text{if } n=1 \text{ then } r:=0 \\
& \text{else } t:=t+1. \ n:=\text{div } n \ 2. \ t' = t+2 \times r'. \ t:=t+1. \ r:=r+1 \text{ fi}
\end{aligned}$$

by cases. First case:

$$\begin{aligned}
t' = t + 2 \times r' & \Leftarrow n=1 \wedge (r:=0) & \text{expand assignment} \\
= & \quad t' = t + 2 \times r' \Leftarrow n=1 \wedge r'=0 \wedge n'=n \wedge t'=t \\
= & \quad \top
\end{aligned}$$

Last case:

$$\begin{aligned} & t' = t+2 \times r' \Leftarrow n \neq 1 \wedge (t := t+1. \ n := \text{div } n \ 2. \ t' = t+2 \times r'. \ t := t+1. \ r := r+1) \\ = & t' = t+2 \times r' \Leftarrow n \neq 1 \wedge (t := t+1. \ n := \text{div } n \ 2. \ t' = t+2 \times r'. \ t := t+1. \ r' = r+1 \wedge n' = n \wedge t' = t) && \text{expand final assignment} \\ = & t' = t+2 \times r' \Leftarrow n \neq 1 \wedge (t' = t+1+2 \times r'. \ r' = r+1 \wedge n' = n \wedge t' = t+1) && \text{substitution law in two parts} \\ = & t' = t+2 \times r' \Leftarrow n \neq 1 \wedge t' = t+2+2 \times (r'-1) && \text{one-point} \\ = & \top \end{aligned}$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder