

Assignment Project Exam Help

ECON 61001: *Hypothesis Testing*

<https://powcoder.com>

Alastair R. Hall

The University of Manchester

Add WeChat powcoder

- asset returns: $R - R_f = \beta_0(R_m - R_f) + \text{error}$

Assignment Project Exam Help

- $\beta_0 < 0 \Rightarrow$ stock is inversely related to market index;
- $\beta_0 = 1 \Rightarrow$ stock moves in line with market index.
- returns to education (dropping exp to simplify and reparameterizing).

$$\ln(w) = \beta_{0,1} + \beta_{0,2} * \text{edu} + \beta_{0,3} * D + \beta_{0,4} * (D * \text{edu}) + \text{error}$$

Add WeChat powcoder

- $D = 1$ is female and zero else
- $\beta_{0,3} = 0, \beta_{0,4} = 0 \Rightarrow$ no difference between men and women.

Assignment Project Exam Help

- aggregate production function:

$$\ln Q = \beta_{0,1} + \beta_{0,2} * \ln(L) + \beta_{0,3} * \ln(K) + \text{error}$$

<https://powcoder.com>

- $\beta_{0,2} + \beta_{0,3} \begin{cases} < \\ = \\ > \end{cases} 1 \Rightarrow \begin{cases} \text{diminishing} \\ \text{constant} \\ \text{increasing} \end{cases} \text{ returns to scale}$

Add WeChat powcoder

Assignment Project Exam Help

- A statistical hypothesis is a conjecture about the distribution of one or more random variables.
- The classical theory of hypothesis testing provides a framework for deciding whether a particular hypothesis is correct.
- Within this framework, there are only two possible decisions: the hypothesis is true or it is not. A decision procedure for such a problem is called a test.

<https://powcoder.com>

Add WeChat powcoder

Assume: our hypothesis involves θ , the parameter vector indexing distribution of V , and Θ denotes the parameter space with

$$\Theta \subset \mathbb{R}^p.$$

Divide Θ into two mutually exclusive and exhaustive parts:

$$\Theta_0 = \{\theta : \text{such that the hypothesis is true}\},$$

$$\Theta_1 = \{\theta : \text{such that the hypothesis is false}\}$$

Using this partition, we can state the object as being to test the null hypothesis

$$H_0 : \theta \in \Theta_0$$

against the alternative hypothesis,

$$H_1 : \theta \in \Theta_1.$$

To facilitate the choice between H_0 and H_1 , collect sample of T observations.

Base inference on some function of this sample, known as a test statistic; denoted by S_T .

In a test procedure, divide sample space of S_T into two mutually exclusive and exhaustive regions, R_0 and R_1 , such that

$S_T \in R_0 \Rightarrow H_0$ is accepted or rather not rejected

$S_T \in R_1 \Rightarrow H_0$ is rejected in favour of H_1

where

- R_0 is known as the acceptance region.
- R_1 is known as the rejection region or the critical region.

Choice of R_0 and R_1 based on possible outcomes of test.

Decision may be correct, but can also make error.

- a **Type I error**, in which H_0 is rejected when it is true;
- a **Type II error**, in which H_0 is not rejected when its false.

<https://powcoder.com>

Ideally we would make both errors as small as possible but there is a tension between them.

So limit $P(\text{Type I error})$ to be no larger than: 0.1, 0.05 or 0.01.

To do this, we need to know the distribution of the test statistic under H_0 .

Size and significance level of a test

Let $P_\theta(\cdot)$ denote the probability of the event in parentheses if the parameter vector takes the value θ .

Define $\alpha(\theta) = P_\theta(R_1)$ that is, $\alpha(\theta)$ describes the probability of a type I error for values of θ that satisfy H_0 .

The quantity $\sup_{\theta \in \Theta_0} \alpha(\theta)$ is known as the size of the test, and equals the maximal probability of a type I error.

To implement test, specify α such that $\alpha(\theta) \leq \alpha$ for all $\theta \in \Theta_0$.

- α is an upper bound on the probability of a type one error.
- $100\alpha\%$ is known as the **significance level** of the test.
- In general, the size and α coincide (but they need not.)

Suppose that:

- $v_t \sim N(\theta, \sigma^2)$, $t = 1, 2, \dots, T$; assume σ^2 known.

- wish to test $H_0 : \theta = 0$ versus $H_1 : \theta \neq 0$.

- can show $\bar{v}_T \sim N(0, \sigma^2/T)$, where $\bar{v}_T = T^{-1} \sum_{t=1}^T v_t$.

→ test statistic is:

$$\tau_T = \frac{\bar{v}_T}{\sqrt{\sigma^2/T}}$$

Under H_0 , $\tau_T \sim N(0, 1)$.

Reject H_0 if τ_T is sufficiently far away from zero.

So the decision rule takes form: reject H_0 if $|\tau_T| > c$ for some constant c .

Choose c to control $P(\text{Type I error})$.

- $\Theta_0 = \{0\}$ and so $\sup_{\theta \in \Theta_0} \alpha(\theta) = P(|\tau_T| > c \mid \theta = 0)$.

- $\Rightarrow c = t_{1-\alpha/2}$


\Rightarrow (for example): reject $H_0 : \theta = 0$ in favour of $H_1 : \theta \neq 0$ at the 5% significance level if $|\tau_T| > 1.96$.

Example 2.8 in Lecture Notes: Acceptance and rejection regions

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



Assignment Project Exam Help

The observed significance level or p-value is the significance level for which the test statistic lies on the boundary of the acceptance and rejection regions.

<https://powcoder.com>

In our example: p-value = δ such that $|\tau_T| = z_{1-\delta/2}$.

Interpretation of p-value: we reject H_0 at all significance levels $100\alpha\%$ for which $\alpha > p\text{-value}$.

Add WeChat powcoder

Further discussion of the topics in this podcast can be found in Section 2.3.1 of the Lecture Notes.

Assignment Project Exam Help

Same source discusses application of framework to hypotheses of the form:

<https://powcoder.com>

- $H_0 : \theta \leq 0$ versus $H_1 : \theta > 0$
- $H_0 : \theta \geq 0$ versus $H_1 : \theta < 0$.

Add WeChat powcoder