## Assignment Project Exam Help

https://powcoder.com

The University of Manchester

Add WeChat powcoder

#### Outline of this lecture

• Time series regression models with non-spherical errors

## Assignment Project Exam Help

https://powcoder.com

- Instrumental Variables estimation Add WeChat powcoder
  - Models with endogenous regressors
  - ullet OLS as MoM o IV
  - Examples of instruments

#### OLS

In Lecture 5 we show that conventional OLS inference framework goes through under the following assumptions.

# Assignment Project Exam Help • Assumption TS1: $y_t = x_t'\beta_0 + u_t$ , t = 1, 2, ... T

- Assumption TS2;  $(y_t, h'_t)$  is a weakly stationary, weakly entropy. Sime specific WCOCET.COM
- Assumption TS3:  $E[x_t x_t'] = Q$ , a finite, positive definite
- . Ashdd WeChatopowcoder
- Assumption TS5:  $Var[u_t | x_t] = \sigma_0^2$ .
- Assumption TS6: For all  $t \neq s$ ,  $E[u_t u_s | x_t, x_s] = 0$ .

#### OLS - with conditionally heteroscedastic errors

In video on "OLS-based inference in time series regression models with conditionally heteroscedastic errors" we consider OLS based School Consumption TS1:  $y_t = x_t'\beta_0 + u_t$ , t = 1, 2, ..., T

- Assumption TS2-SS:  $(y_t, h'_t)$  is a strongly stationary, weakly entropy in Sime Spowcoder.com
- Assumption TS3:  $E[x_t x_t'] = Q$ , a finite, positive definite matrix.
- . Add WeChatopowcoder
- Assumption TS5-H:  $Var[y_t | \mathcal{I}_t] = \sigma_t^2$ .
- Assumption TS6: For all  $t \neq s$ ,  $E[u_t u_s \mid x_t, x_s] = 0$

### Dynamic completeness

### A sasignment of the person of

• kespinging 75/p b. W. Code there to information set at time t.

- if Assumption The local the the model is said to be der

#### Serial correlation in errors

The framework adopted affects how we view the consequences of serial correlation in the errors and so how we proceed.

### A significant substituting the point A significant A sign

- QLS is still consistent as Assumption TS4 holds.
- . https://poiwcodercecomols

- OLS is likely inconsistent as TS4 may not hold.
- Need to reconsider specification.

#### **OLS-based** inference

If Assumptions TS1, TS2-SS, TS3, TS4, & TS5-H hold then:  $\hat{\beta}_{\mathcal{T}}$  is consistent and Project Exam Help

$$T^{1/2}(\hat{\beta}_T - \beta_0) \stackrel{d}{\rightarrow} N(0, V_{sc})$$

https://powcoder.com

- $V_{sc} = Q^{-1}\Omega Q^{-1}$ ,
- . Add We Chat, powcoder

Estimation of long run variance  $\Omega$ ?

#### Covariance matrix estimation

Use heteroscedasticity and autocorrelation covariance (HAC) matrix estimator:

### Assignment Project Exam Help

 $\hat{\Omega}_{HAC} = \hat{\Gamma}_0 + \sum_{i} \omega(i, T)(\hat{\Gamma}_i + \hat{\Gamma}'_i)$ where type  $\frac{1}{2} \frac{1}{2} \frac$ 

- $\omega(\,\cdot\,,\,\cdot\,)$  chosen so that  $\hat{\Omega}_{HAC}$  is both consistent and psd.

#### Covariance matrix estimation

Popular choice of kernel:

Assignment Projectin Exam Help
$$\omega(i,T) = 1 - a_i, \text{ for } a_i < 1$$

$$\text{https://powcoder.com}$$

• This is an example of a funcated kernel estimator: der Add Wechat powcoder

$$\hat{\Omega}_{HAC} = \hat{\Gamma}_0 + \sum_{i=1}^{b_T} \omega(i, T) \left\{ \hat{\Gamma}_i + \hat{\Gamma}_i' \right\}$$

### Assignment Project Exam Help

• Set  $\hat{V}_{sc} = \hat{Q}^{-1} \hat{\Omega}_{HAC} \hat{Q}^{-1}$ . https://powcoder.com

So can perform integrate pased on QLS provided substitute  $\hat{V}_{sc}$  for  $\hat{\sigma}_T^2 \hat{Q}^{T^1}$  installistic. Viscussed in Adum BOSEV Circular I

### OLS if Assumption TS7 fails

If justify standard OLS-based inference framework via Assumptions TS1-TS3, TS5 & TS7 then serial correlation arises through failure  $\mathbf{Assigntment}$   $\mathbf{Project}$   $\mathbf{Exam}$   $\mathbf{Help}$   $E[y_t \mid \mathcal{I}_t] \neq x_t'\beta_0$ 

This https://poweroder.com

- model is linear but have omitted variables
- . Add We Chat powcoder

So need to re-consider specification.

Recall need model for  $\Sigma$ .

### Assignment Project Exam Help

## $\overset{\text{where}}{\bullet} \underset{\varepsilon_t}{\text{Add}} \underset{\sim}{\text{WeChat powcoder}}$

- Regularity conditions hold (esp  $E[x_t u_t] = 0$ )  $\Rightarrow \hat{\beta}_T \xrightarrow{p} \beta_0$ .

Then GLS is OLS applied to:

# Assignment Project Exam Help

•  $\ddot{y}_t = y_t - \rho_0 y_{t-1}$  etc. ~ "quasi-difference".

Consistency of Sies requires WCOder.com

So conditions for consistency of GLS are stronger than for OLS. Thus, the conditions for the consistency of GLS are stronger than those for the consistency of OLS.

#### GLS: alternative view

If sub  $u_t = y_t - x_t' \beta_0$  in regression model then:

## Assignment-Project(PExtam Help which is same as (re-written) transformed model in GLS.

### Also https://pow.cociem.com

$$y_t = \delta_0 y_{t-1} + x_t' \delta_1 + x_{t-1}' \delta_2 + \varepsilon_t,$$

### in whi Ather etist nat powcoder

$$\delta_2 + \delta_0 * \delta_1 = 0,$$

known as the common factor or COMFAC restrictions.

(c.f. response to violation of TS7.)

Alastair R. Hall

#### Breusch-Godfrey test for serial correlation

Assume errors follow AR(p) process that is,

### Assignment-Project Exam~Help

$$h \underset{\rho_{0,i}}{\text{thps}} : \stackrel{\rho q,i}{/} \bar{p} \underset{0, \text{ for at least one } I}{\text{east one } i} \bar{c} \underset{1}{\text{com}}$$

Add We Chat Powcoder

where  $R^2$  is from regression of  $e_t$  on  $x_t, e_{t-1}, e_{t-2}, \ldots, e_{t-p}$ .

Under  $H_0$ :  $LM_p \xrightarrow{d} \chi_p^2$ .

### Breusch-Godfrey test for serial correlation

Choice of *p*?

## Assignment Project Exam Help • may reflect sampling frequency of data

Interpreting in powcoder.com

- can be caused by AR(p) errors
- · Add We Chaired produceder

Empirical example: See Section 4.4.5 of Lecture Notes

#### Instrumental Variables

Consequences of  $E[x_t u_t] \neq 0$  (violation of *CS4* or *TS4*).

Assignment  $Project_1 Exam_{T^{-1}\sum_{t=1}^{t}x_tx_t'}$  Help

and diffeth will powcoder.com

### Add We Chat powcoder

$$T^{-1}\sum_{t=1}^{T}x_{t}u_{t} \stackrel{p}{\rightarrow} E[x_{t}u_{t}] = \mu \neq 0.$$

So using Slutsky's Theorem:  $\hat{\beta}_T \stackrel{p}{\to} \beta_0 + Q^{-1}\mu \neq \beta_0$ .

Alastair R. Hall

ECON 61001: Lecture 7

#### Instrumental Variables

So OLS is an inconsistent estimator of  $\beta_0$ .

### ssignment Project Exam Help

This can arise in three main ways in econometric models: <a href="https://powcoder.com">https://powcoder.com</a> reverse causality

- omitted variables eChat powcoder

Now consider an example of each.

### Example 1: Economic development and institutions

### Assignment Project Exam Help $ln[y_i] = \beta_0 + r_i \beta_1 + controls + u_i$

wherhttps://powcoder.com

y<sub>i</sub> is indome per dipita in developing country i

- r<sub>i</sub> is quality of institutions in developing country i

r is like de date Wie Chatev powooder

#### Example 2: returns to education

# Assignment Project Exam Help $ln[w_i] = \theta_1 + \theta_e e d_i + controls + u_i$

wher https://powcoder.com

- ed; is the number of years of education of individual i;
- ed; likely date Wite Chatmip Owy Coder "ability" that affect  $w_i$  and  $ed_i$ .

### Example 3: monetary policy reaction function

Clarida, Gali & Gertler (2000): Assignment, Project Exam, Help

- https://prowerouter.com
    $\pi_{t+1}$  = inflation in t+1

- $\underset{\leftarrow}{\text{Add}} \underset{\text{ditput gap in}}{\text{Mat powcoder}}$
- $u_t$  is error satisfying  $E[u_t \mid \Omega_t] = 0$

### Example 3: monetary policy reaction function

### A Sexpectation properties properties to the Lintroduce measurement error because:

#### where

#### OLS as MoM

To motivate IV estimation, we reinterpret OLS as a Method of Moments (MoM) estimator.

### Assignment Projected ExametiHelp

### https://powcoder.com

where  $u_t(\beta) = y_t - x_t'\overline{\beta}$  (and so  $u_t(\beta_0) = u_t$ ).

From Ais perspective extraction at the control of the perspective extraction at the perspective

- $E[x_t u_t(\beta_0)] = 0 \rightarrow \text{consistent estimator}.$
- $E[x_t u_t(\beta_0)] \neq 0 \rightarrow \text{inconsistent estimator.}$

#### IV estimation

Find a  $q \times 1$  vector  $z_t$  such that:

### Assignment Project Exam Help

• need  $q \ge k$  and  $z_t$  "sufficiently related to"  $x_t$  + certain other conditions discussed in the next lecture  $\sum_{z_t} \frac{1}{\sum_{t} \sum_{t}^{\infty}} \frac{1}{\sum_{t}^{\infty}} \frac{1}{\sum_{t}^{$ 

Use the population moment equation in  $(\sharp)$  as basis for estimation

→ Instrumental We Charling to powcoder

Discuss next time how to calculate IV estimator and what its properties are. Conclude by looking at choices of instruments in our three examples.

#### Example 1: Economic development and institutions

Acemoglu et al (2001):

## Assignment Project Exam Help

- $y_i$  is income per capita in developing country i
- · https://spowcoder.com

r is likely correlated with  $u_i$  due to reverse causality.

### Base estimation on the population morphocondition. Base estimation on the population morphocondition.

$$E[z_i u_i] = 0$$

where  $z_i = [1, \text{controls}, M_i]'$  and  $M_i$  is settler mortality in country i.

#### Example 2: returns to education

Angrist & Krueger (1991):

## Assignment Project Exam Help

- $w_i$  equals the weekly wage of individual i;
- · https://powcoderf.com;

ed; likely correlated with  $u_i$  due to omitted variables such as "ability thirdffetw; et ean at powcoder"

Base estimation on

$$E[z_i u_i] = 0$$

where  $z_i = [1, \text{controls}, Q_i]'$  and  $Q_i$  is quarter of birth of i.

### Example 3: monetary policy reaction function

Clarida, Gali & Gertler (2000): Assignment, Project Exam, Help

- https://prowerouter.com
    $\pi_{t+1}$  = inflation in t+1

- $\underset{\leftarrow}{\text{Add}} \underset{\text{ditput gap in}}{\text{Mat powcoder}}$
- $u_t$  is error satisfying  $E[u_t \mid \Omega_t] = 0$

#### Example 3: monetary policy reaction function

# Assignment Present Exam Help

 $r_{t} = c + \beta_{\pi} \pi_{t+1} + \beta_{y} y_{t+1} + \beta_{1} r_{t-1} + \beta_{2} r_{t-2} + \underbrace{\left(u_{t} - \beta_{\pi} v_{\pi,t+1} - \beta_{y} v_{y,t+1}\right)}_{}$ 

### https://powcoder.com

- $v_{(\cdot),t+1} = (\cdot)_{t+1} E[(\cdot)_{t+1}|\Omega_t],$
- Add We Chat powcoder

However:  $E[z_t e_t] = 0$  for any  $z_t \in \Omega_t$  (e.g. macro variables).

#### Further reading

# Assignment Project Exam Help

• Time series regression models with serial correlation,

https://pow/coderincom.il on "other conditions" in limit theorems than needed for the course), 20.5

### Add WeChat powcoder

- Testing for serial correlation 20.7
- IV 8.1, 8.2 and 8.3