

# Assignment Project Exam Help

ECON 61001: Review of OLS in simple linear regression

<https://powcoder.com>

Alastair R. Hall

The University of Manchester

Add WeChat powcoder

- Origins of OLS

Assignment Project Exam Help

- Simple linear regression model

- Intuition behind OLS

- Formal definition of OLS

- Other approaches and advantages of OLS

<https://powcoder.com>  
Add WeChat powcoder

Resources: any introductory econometrics text such as Wooldridge (2019) *Introductory Econometrics*.

The origin of Ordinary Least Squares (OLS) is shrouded in controversy.

## Assignment Project Exam Help

- The method is published by
- Adrien Legendre, a French mathematician, in 1805.

- Robert Adrain, an American, in 1808

- Carl Friedrich Gauss, a German, in 1809

• but Gauss claimed he had been using the method since 1795.

## Add WeChat powcoder

Stigler (1981): “The method of least squares is the automobile of modern statistical analysis” but who was the “Henry Ford of statistics”?



Assignment Project Exam Help

<https://powcoder.com>

Gauss was a very eminent mathematician whose key contributions include:

Add WeChat powcoder

- the Normal (or "Gaussian") distribution.
- development of Least Squares theory
- Gaussian elimination

# Assignment Project Exam Help

Stigler (1981) evaluates Gauss's claim using historical records and concludes

*Just as the automobile was not the product of one man of genius, so too the method of least squares is due to [...] at least two independent discoverers. Gauss may well have been the first of these, but he was no Henry Ford of statistics. If there was any single scientist who first put the method within the reach of the common man, it was Legendre. (Stigler, 1981, p.472)*

Suppose we are interesting in modeling the relationship between  $y_i$ , the annual salary of the CEO of a firm  $i$ , and  $x_i$  the average return on equity for the CEO's firm for the previous 3 years.

Assume simple linear regression model

$$y_i = \beta_{0,1} + \beta_{0,2}x_i + u_i = x_i'\beta_0 + u_i$$

where

- $u_i$  is the unobserved error term;
- $x_i' = (1, x_i)$ ;
- $\beta_0 = \begin{bmatrix} \beta_{0,1} \\ \beta_{0,2} \end{bmatrix}$  are unknown (regression) parameters.

# Assignment Project Exam Help

Under assumptions discussed in the lecture, we have

$$E[y_i | x_i] = \beta_{0,1} + \beta_{0,2}x_i$$

<https://powcoder.com>


So  $E[y_i | x_i]$  is linear function of  $x_i$  but weights (the regression parameters) are unknown.

Add WeChat powcoder

So collect sample on  $\{y_i, x_i\}$  and use this to estimate  $\beta_0$ .

Suppose our data consists of the following five observations on  $(y_i, x_i)$ :

$\{(1095, 14.1), (1001, 10.9), (1122, 23.5), (578, 5.9), (1145, 20.9)\}$ .




<https://powcoder.com>

Add WeChat powcoder



No single line passes through all the points.

Assignment Project Exam Help




<https://powcoder.com>

Add WeChat powcoder

Choose line that comes “closest” to fitting the scatter plot → issue of how to measure distance of actual  $y$  from value predicted by line,  $\beta_1 + \beta_2 x$ .

Define  $u_i(\beta) = y_i - \beta_1 - \beta_2 x_i$



## Add WeChat powcoder

Measure of distance from line must be non-negative: in OLS we measure this distance by  $\{u_i(\beta)\}^2$


OLS estimator of  $(\beta_1, \beta_2)$  is value that minimizes  $\sum_{i=1}^5 \{u_i(\beta)\}^2$ .

# OLS in the simple linear regression model

In this example the OLS line is:

$\hat{y}_i = -573.63 + 27.86x_i$

# Assignment Project Exam Help



Notice that  $\{u_i(\beta)\}^2$  is just one possible measure of distance.

For example, could also use  $|u_i(\beta)|$

# Assignment Project Exam Help

If choose  $(\beta_1, \beta_2)$  to minimize  $\sum_{i=1}^5 |u_i(\beta)| \rightarrow$  Least Absolute Deviation (LAD) regression

# <https://powcoder.com>

LAD actually proposed in 1757 by Roger Boscovich but OLS became the “automobile of modern statistical analysis” because:

# Add WeChat powcoder

- calculus of OLS is far easier
- OLS can be shown to have some desirable statistical properties (see lectures).