

# Assignment Project Exam Help

ECON 61001: *Lecture 5*

<https://powcoder.com>

Alastair R. Hall

The University of Manchester

## Add WeChat powcoder

- Large sample behaviour of OLS with time series data

Assignment Project Exam Help

- time series regression models

- concepts and conditions

- OLS as projection

- Non-spherical errors

Add WeChat powcoder

- Properties of OLS

- Generalized Least Squares

Two types of time series:

- **flow variable** – measured over an interval of time, for example monthly consumption expenditures;
- **stock variable** – measured at a moment in time, such as price or quantity of shares owned.

- Assume time series observed at regularly spaced intervals.

- Frequency at which time series is observed known as *sampling frequency*.

$$y_t = \beta_{0,1} + \beta_{0,2}y_{t-1} + \beta_{0,3}h_t + \beta_{0,4}h_{t-1} + u_t = x_t'\beta_0 + u_t$$

## Assignment Project Exam Help

- contains lagged “ $y$ ” and contemporaneous and lagged values of other variables (in this case “ $h$ ”).

Presence of lagged variables on the rhs of regression model  $\Rightarrow$  difference between the number of observations on the variables and the number of observations used in the estimation.

If start with  $T_*$  observations and  $p$  is the longest lag on rhs then **effective sample size** is  $T_* - p$  with the first  $p$  observations being used for conditioning.

For ease of notation, assume effective sample runs  $t = 1, 2, \dots, T$  and  $y_0, h'_0, \dots, y_{-p+1}, h'_{-p+1}$  are available for conditioning.

The sampling framework for time series is fundamentally different from that in analysis of cross-section data. Based on **stochastic process theory**.

## Assignment Project Exam Help

In stochastic process theory: time series  $v_t$  is viewed as evolving before we start observing it and continuing to evolve after we stop observing it, that is

<https://powcoder.com>

$$\dots, v_{-3}, v_{-2}, v_{-1}, v_0, \underbrace{v_1, v_2, v_3, \dots, v_T}_{\text{sample}}, v_{T+1}, v_{T+2}, \dots$$

## Add WeChat powcoder

The entire process  $\{v_t\}_{t=-\infty}^{\infty}$  is known as a *realization* of  $v_t$ .

**Key difference:** sample *once* leading to a particular realization of the series.

So, as the sample size grows, we see more of one realization of the process.

Does this allow us to uncover the underlying probability distribution of  $y_t$  as  $T \rightarrow \infty$ ?

Answer: Yes! Under certain conditions  $\rightarrow$  *stationarity* and *weak dependence*.

We distinguish two forms of stationarity: strong- and weak-stationarity.

The time series  $\{v_t\}_{t=-\infty}^{\infty}$  is said to be **strongly stationary** if the joint probability distribution function,  $F(\cdot)$ , of any subset of  $\{v_t\}$  satisfies:

$$F(v_{t_1}, v_{t_2}, \dots, v_{t_n}) = F(v_{t_1+c}, v_{t_2+c}, \dots, v_{t_n+c})$$

for any integer  $n$  and integer constant  $c$ .

# Add WeChat powcoder

- Also known as “strict” - stationarity

# Weak stationarity

The time series  $\{v_t\}_{t=-\infty}^{\infty}$  is said to be weakly stationary if for all  $t, s$  we have:

- (i)  $E[v_t] = \mu$ ; (independent of  $t$ )
- (ii)  $\text{Var}[v_t] = \Sigma$ ; (independent of  $t$ )
- (iii)  $\text{Cov}[v_t, v_s] = \Sigma_{t-s}$ . (depend only on  $t - s$ )

<https://powcoder.com>

- if  $v_t$  is scalar then  $\text{Cov}[v_t, v_s]$  is  $|t - s|^{\text{th}}$  autocovariance of  $v_t$ .

- if  $v_t$  is vector then  $\text{Cov}[v_t, v_t]$  is  $(t - s)^{\text{th}}$  autocovariance matrix of  $v_t$ :

- diagonal elements are autocovariances of  $v_{t,i}$ .
- off-diagonal elements are  $\text{Cov}[v_{t,i}, v_{s,j}]$ .
- $\text{Cov}[v_t, v_s] = \{\text{Cov}[v_s, v_t]\}'$ .



Weak dependence places restrictions on the **memory** of  $v_t$  that is, the relationship between  $v_t$  and  $v_{t-s}$  as  $s \rightarrow \infty$ .

If  $v_t$  is weakly stationary process then weak dependence implies that  $\text{Cov}[v_t, v_{t-s}] \rightarrow 0$  as  $s \rightarrow \infty$  and at a sufficiently fast rate.

<https://powcoder.com>

Can derive WLLN and CLT for (strong or weak) stationary and weak dependent series (subject to “certain other conditions” that are taken to hold without statement)

Note these conditions are sufficient and not necessary - see lecture notes.

**Weak Law of Large Numbers:** Let  $v_t$  be a stationary and weakly dependent time series with  $E[v_t] = \mu$  then, subject to certain other conditions, it follows that

Assignment Project Exam Help

$$T^{-1} \sum_{t=1}^T v_t \xrightarrow{P} \mu.$$

<https://powcoder.com>

**Central Limit Theorem:** Let  $v_t$  be a stationary and weakly dependent time series with  $E[v_t] = \mu$  and  $\text{Cov}[v_t, v_{t-j}] = \Gamma_j$  then, subject to certain other conditions, it follows that

Add WeChat powcoder

$$T^{-1/2} \sum_{t=1}^T (v_t - \mu) \xrightarrow{d} N(0, \Omega),$$

where  $\Omega = \sum_{-\infty}^{\infty} \Gamma_i = \Gamma_0 + \sum_{i=1}^{\infty} \{\Gamma_i + \Gamma_i'\}$ .

$\Omega$  is known as the **long run variance** of  $v_t$ .

Form of  $\Omega = \lim_{T \rightarrow \infty} \Omega_T$  comes from:

$$\Omega_T = \text{Var} \left[ T^{-1/2} \sum_{t=1}^T (v_t - \mu) \right] = T^{-1} \sum_{t=1}^T \sum_{s=1}^T \text{Cov}[v_t, v_s]$$

<https://powcoder.com>

$$= T^{-1} \sum_{t=1}^T \sum_{s=1}^T \Gamma_{t-s}, \text{ using stationarity,}$$

$$= \Gamma_0 + \sum_{i=1}^{T-1} \left( \frac{T-i}{T} \right) (\Gamma_i + \Gamma_{-i}).$$

$$\text{So } \lim_{T \rightarrow \infty} \Omega_T = \Gamma_0 + \sum_{i=1}^{\infty} \{\Gamma_i + \Gamma_{-i}\}$$

# Finite sample properties of OLS

Recall model with stochastic regressors.

Assumptions:

- SR1: true model is:  $y = (X'\beta_0 + u$

- SR2:  $X$  is stochastic.

- SR3:  $X$  is rank  $k$  with probability 1

- SR4:  $E[u|X] = 0$ .

- SR5:  $\text{Var}[u|X] = \sigma_0^2 I_T$ .

- SR6:  $u|X \sim \text{Normal}$ .

Argued OLS unbiased via:

$$\begin{aligned} E[\hat{\beta}_T] &= \beta_0 + E_X [E_{u|X} [(X'X)^{-1}X'u]] \\ &= \beta_0 + E_X [(X'X)^{-1}X'E_{u|X}[u]] = \beta_0 \end{aligned}$$

Nature of condition  $E[u|X] = 0$  in time series.

Whether it holds depends on “degree of **exogeneity**” of variables in

## Assignment Project Exam Help

- $x_t$  is said to be **contemporaneously exogenous** if  $E[u_t|x_t] = 0$   
 $\Rightarrow u_t$  and  $x_t$  are uncorrelated.

- $x_t$  is said to be **strictly exogenous** if  $E[u_t|\{x_t\}_{t=1}^T] = 0 \Rightarrow \{u_t\}_{t=1}^T$  and  $\{x_t\}_{t=1}^T$  are uncorrelated.

Since  $E[u_t|\{x_t\}_{t=1}^T] = E[u_t|X]$ ,  $E[u|X] = 0$  only holds if  $x_t$  is strictly exogenous.

Note: Assumption SR4 must fail if  $x_t$  contains lagged values of  $y$ , see Lecture Notes. In these cases, in general,  $E[\hat{\beta}_T] \neq \beta_0$ .

Assuming  $x_t = (1, x'_{2,t})'$  and  $x_{2,t}$  is function of (vector)  $h_t$  and  $y_{t-i}$ ,  $h_{t-j}$  (for  $i, j > 0$ ).

## Assignment Project Exam Help

- **Assumption TS1:**  $y_t = x'_t\beta_0 + u_t$ ,  $t = 1, 2, \dots, T$
- **Assumption TS2:** For  $(y_t, h'_t)$  is a weakly stationary, weakly dependent time series.
- **Assumption TS3:**  $E[x_t x'_t] = Q$ , a finite, positive definite matrix.
- **Assumption TS4:**  $E[u_t | x_t] = 0$  for all  $t = 1, 2, \dots, T$ .
- **Assumption TS5:**  $\text{Var}[u_t | x_t] = \sigma_0^2$  for all  $t = 1, 2, \dots, T$ .
- **Assumption TS6:** For all  $t \neq s$ ,  $E[u_t u_s | x_t, x_s] = 0$

<https://powcoder.com>

Add WeChat powcoder

Under these conditions can use essentially same arguments as for cross-section data to deduce the following results:

Assignment Project Exam Help

**Theorem** *If Assumptions TS1 - TS4 hold then  $\hat{\beta}_T$  is a consistent estimator for  $\beta_0$ .*

<https://powcoder.com>

**Theorem** *If Assumptions TS1 - TS6 hold then:*

$$T^{1/2}(\hat{\beta}_T - \beta_0) \xrightarrow{d} N(0, \sigma_0^2 Q^{-1}).$$

Add WeChat powcoder

All large sample inference procedures described in Lecture 4 go through.

# Dynamic completeness

Regression models can be interpreted as a statement about the conditional mean of  $y_t$  given an information set.

In cross-section data this information set contains information about the  $i^{th}$  sampling unit:  $E[y_i | x_i] = x_i' \beta_0$ .

In time series data, the relevant information set is not only the  $h_t$  but the history of both  $y$  and  $h$  that is

$$\mathcal{I}_t = \{ h_t, y_{t-1}, h_{t-1}, y_{t-2}, h_{t-2}, \dots, y_1, h_1 \}.$$

Therefore if regression model specifies conditional mean then we are really stating that we believe:

$$E[y_t | \mathcal{I}_t] = x_t' \beta_0,$$

- if this statement is true then model is said to be *dynamically complete*.



# Assignment Project Exam Help

It is convenient to introduce this property as a specific assumption.

- **Assumption TS7:**  $E[y_t | \mathcal{I}_t] = x_t' \beta_0$ .

<https://powcoder.com>

Note:

- Assumption TS7  $\Rightarrow$  Assumptions TS4 and TS6, see Lecture Notes

Add WeChat powcoder

Suppose wish to predict  $y_t$  given  $x_t$  using  $\tilde{y}_t = c(x_t)$ .

## Assignment Project Exam Help

Issue: choice of  $c(\cdot)$ .

If choose  $c(\cdot)$  to minimize

$$\text{MSE}(\tilde{y}_t) = E[y_t - \tilde{y}_t]^2$$

$$\rightarrow c_o(x_t) = E[y_t | x_t].$$

If  $c_o(\cdot)$  unknown then might restrict to class of linear forecasts

$y_t^{lp} = \alpha' x_t$  but then what should  $\alpha$  be?

Choice that minimizes MSE (over class of linear forecasts) is one associated with the linear projection of  $y_t$  on  $x_t$  which has the property

# Assignment Project Exam Help

$E[(y_t - \alpha' x_t)x_t] = 0.$   
<https://powcoder.com>

$\rightarrow \alpha = \{E[x_t x_t']\}^{-1} E[x_t y_t] \sim \text{population analogue to } \hat{\beta}_T.$

So OLS can be justified as estimator of weights in linear projection of  $y_t$  on  $x_t$  - but this does not justify using estimators to learn about how  $x_t$  affects  $y_t$ , for this we need to impose assumptions about the relationship between the variables.

# Add WeChat powcoder

Have developed large sample framework for inference in cross-section and time series data.

## Assignment Project Exam Help

Do assumptions always hold? No! In next part of course, we consider consequences of violations of assumptions about second moments of error term.

<https://powcoder.com>

Terminology (using fixed regressor model to simplify):

- $\text{Var}[u] = \sigma_0^2 I_T$  then  $u$  is said to have a spherical distribution.
- $\text{Var}[u] \neq \sigma_0^2 I_T$  then  $u$  is said to have a non-spherical distribution.

See Lecture notes Section 4.1 for origins of these terms.

# Assignment Project Exam Help

Specifically, we consider two examples of non-spherical errors:

- Cross-section data with  $\text{Var}[u_i|x_i] = \sigma_i^2$  - heteroscedasticity

<https://powcoder.com>

- Time series data with

- $\text{Var}[u_t|x_t] = \sigma_t^2$  - heteroscedasticity

Add WeChat powcoder

- $\text{Cov}[u_t, u_s|x_t, x_s] \neq 0$  - serial correlation

Recall that our model is:

$$y = X\beta_0 + u$$

where

- CA1: true model is:  $y = X\beta_0 + u$ .

- CA2:  $X$  is fixed in repeated samples.

- CA3:  $X$  is rank  $k$ .

- CA4:  $E[u] = 0$ .

- CA5-NS  $\text{Var}[u] = \Sigma$  where  $\Sigma$  is a  $T \times T$  positive definite matrix.

- CA6:  $u \sim \text{Normal}$ .

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

- $E[\hat{\beta}_T] = \beta_0$  as impose Assumptions CA1-CA4.

- $Var[\hat{\beta}_T] = (X'X)^{-1}X'E[uu']X(X'X)^{-1}$  and so

$$Var[\hat{\beta}_T] \neq (X'X)^{-1}X'\Sigma X(X'X)^{-1} \neq \sigma_0^2(X'X)^{-1}.$$

- $\hat{\beta}_T \sim N(\beta_0, (X'X)^{-1}X'\Sigma X(X'X)^{-1})$ .

Add WeChat powcoder

So inference procedures from Lectures 2, 3 and 4 are not valid because are based on wrong formula for  $Var[\hat{\beta}_T]$ .

Conditions of Gauss-Markov Theorem do not hold and so this result cannot be used to justify OLS is BLUE.

## Assignment Project Exam Help

Can we characterize the BLUE estimator? Yes!

It is the **The Generalized Least Squares (GLS)** estimator of  $\beta_0$  given by

$$\hat{\beta}_{GLS} = (X' \Sigma^{-1} X)^{-1} X' \Sigma^{-1} y$$

If Assumptions **CA1-CA4**, **CA5-VS** and **CV6** hold then

$$\hat{\beta}_{GLS} \sim N(\beta_0, (X' \Sigma^{-1} X)^{-1}).$$

But need  $\Sigma$  - what happens if  $\Sigma$  is unknown?



# Assignment Project Exam Help

- Notes: Sections 2.12, 3.3-3.5, 4.1, 4.2.

- Greene:

- time series/data Section 20.1
- conditions for limit theorems Section 20.2 and Section 20.4 (but more detail than in the course)
- OLS with non-spherical errors Sections 9.1 and 9.2 (finite sample part)

Add WeChat powcoder