

Assignment Project Exam Help

ECON3350/7350
Univariate Time Series - I

<https://powcoder.com>

Eric Eisenstat

The University of Queensland

Add WeChat powcoder

Tutorial 2

Stationarity

When a single realisation of observations is available, the aggregation of observations over time is important.

Definition

A stochastic process is **stationary** if the data generating process is such that the mean, variance and covariances are independent of time.

$$E(y_t) = \mu$$

$$Var(y_t) = E[(y_t - \mu)^2] = \sigma_y^2 = \gamma_0 \quad Cov(y_t, y_{t-k}) = E[(y_t - \mu)((y_{t-k} - \mu))] = \gamma_k$$

$$k = 1, 2, \dots$$

These conditions must be satisfied for all values of t .

At this initial stage we will only consider stationary processes and we will relax this assumption later in the course.

Autoregressive (AR) Models

To model the dependence in y_t upon its own past behaviour.

AR(1)

Assignment Project Exam Help

$$y_t = a_0 + a_1 y_{t-1} + \varepsilon_t$$

AR(2)

<https://powcoder.com>

$$y_t = a_0 + a_1 y_{t-1} + a_2 y_{t-2} + \varepsilon_t$$

AR(p)

Add WeChat powcoder

$$y_t = a_0 + a_1 y_{t-1} + \dots + a_p y_{t-p} + \varepsilon_t$$

$$y_t = a_0 + \sum_{i=1}^p a_i y_{t-i} + \varepsilon_t$$

Assignment Project Exam Help

$$y_t = \mu + \varepsilon_t + \beta_1 \varepsilon_{t-1}$$

- MA(1) <https://powcoder.com>

$$y_t = \mu + \varepsilon_t + \beta_1 \varepsilon_{t-1} + \beta_2 \varepsilon_{t-2}$$

- MA(q) Add WeChat powcoder

$$y_t = \mu + \varepsilon_t + \beta_1 \varepsilon_{t-1} + \dots + \beta_q \varepsilon_{t-q}$$

Assignment Project Exam Help

- ARMA(1,1)

$$y_t = a_0 + a_1 y_{t-1} + \varepsilon_t + \beta_1 \varepsilon_{t-1}$$

- ARMA(2,2)

<https://powcoder.com>

$$y_t = a_0 + a_1 y_{t-1} + a_2 y_{t-2} + \varepsilon_t + \beta_1 \varepsilon_{t-1} + \beta_2 \varepsilon_{t-2}$$

- ARMA(3,1)

Add WeChat powcoder

$$y_t = a_0 + a_1 y_{t-1} + a_2 y_{t-2} + a_3 y_{t-3} + \varepsilon_t + \beta_1 \varepsilon_{t-1}$$

Definitions

- $Var(y_t) = E[(y_t - \mu)^2] = \gamma_0$
- $Cov(y_t, y_{t-k}) = E[(y_t - \mu)(y_{t-k} - \mu)] = \gamma_k \quad k = 1, 2, \dots$

Definitions

Autocovariance Function: $\gamma_k, k = 1, 2, \dots$ If the process is stationary $\gamma_k = \gamma_{-k}$.

Autocorrelation Function (ACF): $\rho_k = \frac{\gamma_k}{\gamma_0}, k = 1, 2, \dots$

Correlogram or **SACF:** Plot of the sample autocorrelation function, r_k , against k .

Partial Autocorrelation Function (PACF)

Definition

The **partial autocorrelation function** (ϕ_{kk}) is given by the k th coefficients in the corresponding $AR(k)$ system of autoregressions.

$$AR(1) \quad y_t = a_0 + a_1 y_{t-1}; \quad \phi_{11} = a_1$$

$$AR(2) \quad y_t = a_0 + a_1 y_{t-1} + a_2 y_{t-2}; \quad \phi_{22} = a_2$$

\vdots

$$AR(k) \quad y_t = a_0 + a_1 y_{t-1} + a_2 y_{t-2} + \dots + a_k y_{t-k}; \quad \phi_{kk} = a_k$$

Add WeChat powcoder

We identify the DGP by plotting the sample ACF (SACF) and sample PACF (SPACF) together.

Using the Yule-Walker equations, we can work out the PACF from the ACF,

Assignment Project Exam Help

$\phi_{11} = \rho_1$
 $\phi_{22} = (\rho_2 - \rho_1^2) / (1 - \rho_1^2)$
 \vdots
<https://powcoder.com>

Add WeChat powcoder
 $\phi_{kk} = \frac{\rho_k - \sum_{j=1}^{k-1} \phi_{k-1,j} \rho_{k-j}}{1 - \sum_{j=1}^{k-1} \phi_{k-1,j} \rho_j}, k = 3, 4, 5, \dots$

where, $\phi_{k,j} = \phi_{k-1,j} - \phi_{kk} \phi_{k-1,k-j}, j = 1, 2, 3, \dots, k-1$