

Assignment Project Exam Help

ECON3350/7350

Deterministic and Stochastic Trends

<https://powcoder.com>

Eric Eisenstat

The University of Queensland

Add WeChat powcoder

Tutorial 5

Assignment Project Exam Help

$$y_t = a_0 + a_2 t + a_1 y_{t-1} + \varepsilon_t; |a_1| < 1$$

- y_t is trend stationary because if we take the trend out the new process is stationary.
- Implies a_2 is the average or long run growth rate.

<https://powcoder.com>
Add WeChat powcoder

Assignment Project Exam Help

- An alternative form of trend comes from $a_2 = 0$ and $a_1 = 1$

$$y_t = a_0 + a_2 t + a_1 y_{t-1} + \varepsilon_t$$

<https://powcoder.com>

$$y_t = a_0 + y_{t-1} + \varepsilon_t \text{ or } \Delta y_t = a_0 + \varepsilon_t$$

a_0 is a trend and can be interpreted as the average or long run growth rate..

Add WeChat powcoder

Consider the stationary AR(1)

Assignment Project Exam Help

$$y_t = a_1 y_{t-1} + \varepsilon_t \text{ with } |a_1| < 1$$
$$= a_1^t y_0 + \varepsilon_t + a_1 \varepsilon_{t-1} + \cdots + a_1^{t-1} \varepsilon_1$$

The effect of shocks or innovations dies out since $a_1^t \rightarrow 0$ as t increases.
Contrast this with the **Random Walk Model**

Add WeChat powcoder

$$y_t = y_{t-1} + \varepsilon_t \text{ or } \Delta y_t = \varepsilon_t$$
$$= y_0 + \sum_{i=0}^t \varepsilon_{t-i}$$

All past shocks have a permanent effect.



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Random Walk with Drift

The original model

$$y_t = a_0 + y_{t-1} + \varepsilon_t \text{ or } \Delta y_t = a_0 + \varepsilon_t$$

The expected change is $E(\Delta y_t) = a_0$. If $a_0 > 0$, y_t will tend to drift upwards. Hence this is called the **Random Walk with Drift**.

Substituting back

$$y_t = a_0 + y_{t-1} + \varepsilon_t$$

$$= a_0 + \mu + y_{t-2} + \varepsilon_{t-1} + \varepsilon_t$$

$$= a_0 t + y_0 + \varepsilon_t + \varepsilon_{t-1} + \cdots \varepsilon_1$$

Thus

$$E(y_t | y_0) = y_0 + a_0 t$$

And, the variance also has a trend in it.

- $y_t \sim I(1)$ integrated of order 1.
- An $AR(p)$ for $y_t \sim I(1)$ implies an $AR(p-1)$ for $\Delta y_t \sim I(0)$.
- Difference of y_t , $\Delta y_t \sim I(0)$. Thus y_t is **difference stationary**.
- Generally, if y_t has d unit roots,

$$y_t \sim I(d),$$

$$\Delta y_t \sim I(d-1),$$

$$\Delta^d y_t \sim I(0)$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

$$\Delta y_t = \alpha_0 + \alpha_1 t + \alpha_2 y_{t-1} + \sum_{i=1}^{p-1} \beta_i \Delta y_{t-i} + \varepsilon_t \quad (1)$$

Assignment Project Exam Help


$$y_t = \alpha_0 + \alpha_1 t + \alpha_2 y_{t-1} + \sum_{i=1}^{p-1} \beta_i \Delta y_{t-i} + \varepsilon_t \quad (2)$$

https://powcoder.com

$$\Delta y_t = \alpha_0 + \alpha_1 t + \alpha_2 y_{t-1} + \sum_{i=1}^{p-1} \beta_i \Delta y_{t-i} + \varepsilon_t \quad (3)$$

Add WeChat powcoder

- Use testing procedure in the next slide and Tables A and B pp 488-489 Enders



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

- Tests of unit roots, e.g., $H_0: y_t \sim I(1)$ against $H_1: y_t \sim I(0)$; often suffer from low power. That is, the probability they lead us to reject the null is low. This may lead us to conclude there are unit roots where there are not.
- One way to circumvent this problem is to test the null there is no unit root against the **alternative** that there is a unit root.

That is, test

Add WeChat powcoder
 $H_0: y_t \sim I(0)$ against $H_1: y_t \sim I(1)$

Assignment Project Exam Help

The Kwiatkowski–Phillips–Schmidt–Shin (KPSS) tests test is one such test. The steps are:

- 1 Estimate $y_t = a_0 + \varepsilon_t$ and save the residuals, e_t
- 2 Compute $S_t = \sum_{s=1}^t e_s$ for $t = 1, 2, \dots, T$.

- 3 Compute $KPSS = T^{-2} \sum_{t=1}^T \frac{S_t^2}{\hat{\sigma}^2}$

- 4 Compare this to the critical values. - See Table

<https://powcoder.com>

Add WeChat powcoder

If we assume the process is trend stationary under the null, the test becomes:

$$H_0 : y_t \sim \text{trend stationary against } H_1 : y_t \sim I(1)$$

The steps are:

- 1 Estimate $y_t = a_0 + a_2 t + \varepsilon_t$ and save the residuals, e_t
- 2 Compute $S_t = \sum_{s=1}^t e_s$ for $t = 1, 2, \dots, T$.
- 3 Compute $KPSS = T^{-2} \sum_{t=1}^T \frac{e_t^2}{\hat{\sigma}^2}$
- 4 Compare this to the critical values. - See Table

Identification of $ARIMA(p, d, q)$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder




Figure T-1: Schematic representation of the Box-Jenkins methodology for time series modeling.