



MONASH  
University

MONASH  
BUSINESS  
SCHOOL

ETW3420:

Assignment Project Exam Help

Principles of

Forecasting and

Applications

Add WeChat powcoder

Topic 7: Regression Models

Dr. Jason Ng

1 The linear model with time series

2 Residual diagnostics

3 Some useful predictors for linear models

4 Selecting predictors and forecast evaluation

5 Forecasting with regression

6 Matrix formulation

7 Correlation, causation and forecasting

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

## Multiple regression and forecasting

$$y_t = \beta_0 + \beta_1 x_{1,t} + \beta_2 x_{2,t} + \cdots + \beta_k x_{k,t} + \varepsilon_t.$$

■  $y_t$  is the variable we want to predict: the “response” variable

■ Each  $x_{j,t}$  is numerical and is called a “predictor”. They are usually assumed to be known for all past and future times.


■ The coefficients  $\beta_1, \dots, \beta_k$  measure the effect of each predictor after taking account of the effect of all other predictors in the model.

That is, the coefficients measure the **marginal effects**.


■  $\varepsilon_t$  is a white noise error term

## Example: US consumption expenditure

```
autoplot(uschange[,c("Consumption","Income")]) +  
  ylab("% change") + xlab("Year")
```



## Example: US consumption expenditure



## Example: US consumption expenditure

```
tslm(Consumption ~ Income, data=uschange) %>% summary
```


```
##  
## Call:  
## tslm(formula = Consumption ~ Income, data = uschange)  
##  
## Residuals:  
##      Min       1Q   Median       3Q      Max   
## -2.40345 -0.31316  0.02158  0.29978  1.45157   
##  
## Coefficients:  
##              Estimate Std. Error t value Pr(>|t|)      
## (Intercept)  0.51510    0.05169   9.789 < 2e-16 ***  
## Income      0.28960    0.04444   6.515 1.58e-08 ***  
## ---  
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  
##  
## Residual standard error: 0.6026 on 185 degrees of freedom  
## Multiple R-squared:  0.159, Adjusted R-squared:  0.1545   
## F-statistic: 34.98 on 1 and 185 DF, p-value: 1.577e-08
```

## Example: US consumption expenditure


Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



## Example: US consumption expenditure





## Example: US consumption expenditure

```
fit.consMR <- tslm(  
  Consumption ~ Income + Production + Unemployment + Savings,  
  data=uschange)  
summary(fit.consMR)
```

```
##  
## Call:  
## tslm(formula = Consumption ~ Income + Production + Unemployment +  
##       Savings, data = uschange)  
##  
## Residuals:  
##      Min       1Q   Median       3Q      Max   
## -0.88296 -0.17638 -0.03679  0.15251  1.20553   
##  
## Coefficients:  
##              (Intercept)      Income      Production      Unemployment      Savings        
##               0.26729      0.71449      0.04589     -0.20477     -0.04527   
##               0.02721      0.04219      0.02588      0.10550      0.00278   
##               7.1841e-11      16.934      1.773     -1.941     -16.287   
##               1.68e-11      < 2e-16      0.0778      0.0538      < 2e-16   
##               ***                ***                .                .                ***   
## ---
```

```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  
##  
## Residual standard error: 0.3286 on 182 degrees of freedom  
## Multiple R-squared:  0.754, Adjusted R-squared:  0.7486   
## F-statistic: 139.5 on 4 and 182 DF,  p-value: < 2.2e-16
```


Assignment Project Exam Help

<https://powcoder.com>


Add WeChat powcoder

## Example: US consumption expenditure

Percentage change in US consumption expenditure




## Example: US consumption expenditure





## Example: US consumption expenditure

Residuals from Linear regression model



<https://powcoder.com>



Add WeChat powcoder

1 The linear model with time series

2 Residual diagnostics

3 Some useful predictors for linear models

4 Selecting predictors and forecast evaluation

5 Forecasting with regression

6 Matrix formulation

7 Correlation, causation and forecasting

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

# Assignment Project Exam Help

For forecasting purposes, we require the following assumptions:

- $\varepsilon_t$  are uncorrelated and zero mean
- $\varepsilon_t$  are uncorrelated with each  $x_{j,t}$ .

Add WeChat powcoder

# Assignment Project Exam Help

For forecasting purposes, we require the following assumptions:

- $\varepsilon_t$  are uncorrelated and zero mean
- $\varepsilon_t$  are uncorrelated with each  $x_{j,t}$ .

It is **useful** to also have  $\varepsilon_t \sim N(0, \sigma^2)$  when producing prediction intervals or doing statistical tests.

# Assignment Project Exam Help

Useful for spotting outliers and whether the linear model was appropriate.

- Scatterplot of residuals  $\epsilon_t$  against each predictor  $x_{jt}$ .
- Scatterplot residuals against the fitted values  $\hat{y}_t$
- Expect to see scatterplots resembling a horizontal band with no values too far from the band and no pattern, such as curvature or increasing spread.



# Assignment Project Exam Help

- If a plot of the residuals vs any predictor in the model shows a pattern, then the relationship is nonlinear.
- If a plot of the residuals vs any predictor **not** in the model shows a pattern, then the predictor should be added to the model.
- If a plot of the residuals vs fitted values shows a pattern, then there is heteroscedasticity in the errors. (Could try a transformation.)

<https://powcoder.com>

Add WeChat powcoder

## Breusch-Godfrey test

OLS regression:

$$y_t = \beta_0 + \beta_1 x_{t,1} + \dots + \beta_k x_{t,k} + u_t$$

Auxiliary regression:

$$\hat{u}_t = \beta_0 + \beta_1 x_{t,1} + \dots + \beta_k x_{t,k} + \rho_1 \hat{u}_{t-1} + \dots + \rho_p \hat{u}_{t-p} + \varepsilon_t$$

If  $R^2$  statistic is calculated for this model, then

$$(T - p)R^2 \sim \chi_p^2,$$

when there is no serial correlation up to lag  $p$  and  $T$  = length of series.

- Breusch-Godfrey test better than Ljung-Box for regression models.

# Assignment Project Exam Help

```
## Breusch-Godfrey test for serial correlation of order up to 8
##
## data: Residuals from linear regression model
## LM test = 14.874, df = 8, p-value = 0.06163
```

If the model fails the Breusch-Godfrey test ...

- The forecasts are not wrong, but have higher variance than they need to.
- There is information in the residuals that we should exploit.
- This is done with a regression model with ARMA errors.

1 The linear model with time series

2 Residual diagnostics

3 Some useful predictors for linear models

4 Selecting predictors and forecast evaluation

5 Forecasting with regression

6 Matrix formulation

7 Correlation, causation and forecasting

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

# Assignment Project Exam Help

Linear trend

<https://powcoder.com>

- $t = 1, 2, \dots, T$
- Strong assumption that trend will continue.

Add WeChat powcoder

## Dummy variables

If a categorical variable takes only two values (e.g., 'Yes' or 'No'), then an equivalent numerical variable can be constructed taking value 1 if yes and 0 if no. This is called a **dummy variable**.

	A	B
1	Yes	1
2	Yes	1
3	No	0
4	Yes	1
5	No	0
6	No	0
7	Yes	1
8	Yes	1
9	No	0
10	No	0
11	No	0
12	No	0
13	Yes	1
14	No	0

## Dummy variables

If there are more than two categories, then the variable can be coded using several dummy variables (one fewer than the total number of categories).

	A	B	C	D	E
1	Monday	1	0	0	0
2	Tuesday	0	1	0	0
3	Wednesday	0	0	1	0
4	Thursday	0	0	0	1
5	Friday	0	0	0	0
6	Monday	1	0	0	0
7	Tuesday	0	1	0	0
8	Wednesday	0	0	1	0
9	Thursday	0	0	0	1
10	Friday	0	0	0	0
11	Monday	1	0	0	0
12	Tuesday	0	1	0	0
13	Wednesday	0	0	1	0
14	Thursday	0	0	0	1
15	Friday	0	0	0	0

## Beware of the dummy variable trap!

# Assignment Project Exam Help

- Using one dummy for each category gives too many dummy variables!
- The regression will then be singular and inestimable.
- Either omit the constant, or omit the dummy for one category.
- The coefficients of the dummies are relative to the omitted category.

<https://powcoder.com>

Add WeChat powcoder



### Seasonal dummies

- For quarterly data: use 3 dummies
- For monthly data: use 11 dummies
- For daily data: use 6 dummies
- What to do with weekly data?

<https://powcoder.com>

Add WeChat powcoder

# Uses of dummy variables

## Seasonal dummies

- For quarterly data: use 3 dummies
- For monthly data: use 11 dummies
- For daily data: use 6 dummies
- What to do with weekly data?

## Outliers

- If there is an outlier, you can use a dummy variable (taking value 1 for that observation and 0 elsewhere) to remove its effect.

# Uses of dummy variables

## Seasonal dummies

- For quarterly data: use 3 dummies
- For monthly data: use 11 dummies
- For daily data: use 6 dummies
- What to do with weekly data?

## Outliers


- If there is an outlier, you can use a dummy variable (taking value 1 for that observation and 0 elsewhere) to remove its effect.

## Public holidays

- For daily data: if it is a public holiday, dummy=1, otherwise dummy=0.

## Beer production revisited

Australian quarterly beer production



# Assignment Project Exam Help

Regression model

$$y_t = \beta_0 + \beta_1 t + \beta_2 d_{2,t} + \beta_3 d_{3,t} + \beta_4 d_{4,t} + \epsilon_t$$

- $d_{i,t} = 1$  if  $t$  is quarter  $i$  and 0 otherwise.

Add WeChat powcoder

## Beer production revisited

```
fit.beer <- tslm(beer ~ trend + season)
summary(fit.beer)
```

# Assignment Project Exam Help

```
## Call:
## tslm(formula = beer ~ trend + season)
```

```
##
```

```
## Residuals:
```

```
##      Min       1Q   Median       3Q      Max
```

```
## -42.500  -7.599   -0.459   7.901  21.780
```

```
##
```

```
## Coefficients:
```

```
##              Estimate Std. Error t value Pr(>|t|)
```

```
## (Intercept)  441.80044    3.72353  118.333  < 2e-16 ***
```

```
## trend        -6.34627     0.66571   -9.541  2.71e-06 ***
```

```
## season2      -34.65913     5.96832   -5.813  9.10e-03 ***
```

```
## season3      -17.82164     4.02249   -4.430  3.45e-05 ***
```

```
## season4       72.79641     4.02305   18.095  < 2e-16 ***
```

```
## ---
```

```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
##
```


```
## Residual standard error: 12.23 on 69 degrees of freedom
```

```
## Multiple R-squared:  0.9243, Adjusted R-squared:  0.9199
```


```
## F-statistic: 210.7 on 4 and 69 DF, p-value: < 2.2e-16
```

<https://powcoder.com>

Add WeChat powcoder



## Beer production revisited






# Beer production revisited


Assignment Project Exam Help

<https://powcoder.com>



Add WeChat powcoder

## Beer production revisited



Periodic seasonality can be handled using pairs of Fourier terms:

$$s_k(t) = \sin\left(\frac{2\pi kt}{n}\right) \quad c_k(t) = \cos\left(\frac{2\pi kt}{n}\right)$$

$$y_t = a + bt + \sum_{k=1}^K [\alpha_k s_k(t) + \beta_k c_k(t)] + \varepsilon_t$$

- Every periodic function can be approximated by sums of sin and cos terms for large enough  $K$ .
- Choose  $K$  by minimizing AICc.
- Called "harmonic regression"

```
fit <- tslm(y ~ trend + fourier(y, K))
```

# Harmonic regression: beer production

```
fourier.beer <- tslm(beer ~ trend + fourier(beer, K=2))  
summary(fourier.beer)
```

# Assignment Project Exam Help

```
## Call:
## tslm(formula = beer ~ trend + fourier(beer, K = 2))
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -42.500  -7.599  -0.459   7.901  21.780
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   446.87920     2.87321 155.533 < 2e-16 ***
## trend         -0.34027     0.06151  -5.111 2.78e-06 ***
## fourier(beer, k = 2)S1-4   8.31082     2.01125   4.130 3.45e-05 ***
## fourier(beer, K = 2)C1-4  53.72807     2.01125  26.714 < 2e-16 ***
## fourier(beer, K = 2)C2-4  13.98958     1.42256   9.834 9.26e-15 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 12.23 on 69 degrees of freedom
## Multiple R-squared:  0.9243, Adjusted R-squared:  0.9199
## F-statistic: 210.7 on 4 and 69 DF, p-value: < 2.2e-16
```

<https://powcoder.com>

Add WeChat powcoder

# Assignment Project Exam Help

- With Fourier terms, we often need fewer predictors than with dummy variables, especially when  $m$  is large.
- This makes them useful for weekly data, for example, where  $m = 52$ .
- For short seasonal periods (e.g., quarterly data), there is little advantage in using Fourier terms over seasonal dummy variables.

### Spikes

Assignment Project Exam Help

- Equivalent to a dummy variable for handling an outlier ; account for the effect which lasts for only one period.

<https://powcoder.com>

Add WeChat powcoder

### Spikes

■ Equivalent to a dummy variable for handling an outlier : account for the effect which lasts for only one period.

### Steps

- Variable takes value 0 before the intervention and 1 afterwards.

<https://powcoder.com>  
Add WeChat powcoder

### Spikes

- Equivalent to a dummy variable for handling an outlier : account for the effect which lasts for only one period.

### Steps

- Variable takes value 0 before the intervention and 1 afterwards.

### Change of slope

- Variables take values 0 before the intervention and values  $\{1, 2, 3, \dots\}$  afterwards; (See piecewise linear trend)



# Assignment Project Exam Help

For monthly data

- Christmas: always in December so part of monthly seasonal effect
- Easter: use a dummy variable  $v_t = 1$  if any part of Easter is in that month,  $v_t = 0$  otherwise.
- Ramadan and Chinese new year similar.

Assignment Project Exam Help

With monthly data, if the observations vary depending on how many different types of days in the month, then trading day predictors can be useful.

<https://powcoder.com>

$z_1$  = # Mondays in month;

$z_2$  = # Tuesdays in month;

Add WeChat powcoder

$z_7$  = # Sundays in month.

# Assignment Project Exam Help

Lagged values of a predictor

Example:  $x$  is advertising which has a delayed effect

<https://powcoder.com>

$x_1$  = advertising for previous month;

$x_2$  = advertising for two months previously;

Add WeChat powcoder

$x_m$  = advertising for  $m$  months previously.

# Assignment Project Exam Help

Piecewise linear trend with bend at  $\tau$

<https://powcoder.com>

$x_{1,t} = t$   
 $x_{2,t} = \begin{cases} 0 & t < \tau \\ t - \tau & t \geq \tau \end{cases}$   
Add WeChat powcoder

- $\tau$  is the “knot” or point in time at which the line should bend.
- By fitting a piecewise linear trend which bends at some point in time, the nonlinear trend can be constructed via a series of linear pieces.
- If the associated coefficients of  $x_{1,t}$  and  $x_{2,t}$  are  $\beta_1$  and  $\beta_2$ , then,
  - $\beta_1$  = slope of trend before time  $\tau$ .
  - $\beta_1 + \beta_2$  = slope of trend after time  $\tau$ .
- Additional trends can be included in the relationship by adding further variables of the above form.

# Assignment Project Exam Help

Quadratic or higher order trend

<https://powcoder.com>

$$x_{1,t} = t, \quad x_{2,t} = t^2, \quad \dots$$

Add WeChat powcoder

# Assignment Project Exam Help

Quadratic or higher order trend

<https://powcoder.com>


$$x_{1,t} = t, \quad x_{2,t} = t^2, \quad \dots$$

NOT RECOMMENDED!

Add WeChat powcoder

## Example: Boston marathon winning times


```
autoplot(marathon) +  
  xlab("Year") + ylab("Winning times in minutes")
```



```
fit.lin <- tslm(marathon ~ trend)
```




## Example: Boston marathon winning times



## Example: Boston marathon winning times

```
autoplot(residuals(fit.lin)) +  
  xlab("Year") + ylab("Residuals from a linear trend")
```



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

## Example: Boston marathon winning times

```
# Linear trend
fit.lin <- tslm(marathon ~ trend)
fcasts.lin <- forecast(fit.lin, h=10)

# Exponential trend
fit.exp <- tslm(marathon ~ trend, lambda = 0)
fcasts.exp <- forecast(fit.exp, h=10)


# Piecewise linear trend
t.break1 <- 1940
t.break2 <- 1980
t <- time(marathon)
t1 <- ts(pmax(0, t-t.break1), start=1897)
t2 <- ts(pmax(0, t-t.break2), start=1897)
fit.pw <- tslm(marathon ~ t + 1 + t2)
t.new <- t[length(t)] + seq(10)
t1.new <- t1[length(t1)] + seq(10)
t2.new <- t2[length(t2)] + seq(10)
newdata <- cbind(t=t.new, t1=t1.new, t2=t2.new) %>%
  as.data.frame
fcasts.pw <- forecast(fit.pw, newdata = newdata)
```

Assignment Project Exam Help


<https://powcoder.com>

Add WeChat powcoder

## Example: Boston marathon winning times




## Example: Boston marathon winning times



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder





# Assignment Project Exam Help

A spline is a continuous function  $f(x)$  interpolating all points  $(\kappa_j, y_j)$  for  $j = 1, \dots, K$  and consisting of polynomials between each consecutive pair of 'knots'  $\kappa_j$  and  $\kappa_{j+1}$ .

Add WeChat powcoder

# Assignment Project Exam Help

A spline is a continuous function  $f(x)$  interpolating all points  $(\kappa_j, y_j)$  for  $j = 1, \dots, K$  and consisting of polynomials between each consecutive pair of 'knots'  $\kappa_j$  and  $\kappa_{j+1}$ .

- Parameters constrained so that  $f(x)$  is continuous.
- Further constraints imposed to give continuous derivatives.

# Assignment Project Exam Help

- Let  $\kappa_1 < \kappa_2 < \dots < \kappa_K$  be "knots" in interval  $(a, b)$ .
- Let  $\chi_1 = \chi$ ,  $\chi_j = (\chi - \kappa_{j-1})_+$  for  $j = 2, \dots, K+1$ .
- Then the regression is piecewise linear with bends at the knots.

Add WeChat powcoder

# Assignment Project Exam Help

- Let  $x_1 = x$ ,  $x_2 = x^2$ ,  $x_3 = x^3$ ,  $x_j = (x - \kappa_{j-3})_+^3$  for  $j = 4, \dots, K + 3$ .
- Then the regression is piecewise cubic, but smooth at the knots.
- Choice of knots can be difficult and arbitrary.
- Automatic knot selection algorithms very slow.

## Example: Boston marathon winning times

```
# Spline trend
library(splines)
t <- time(marathon)
fit.splines <- lm(marathon ~ ns(t, df=6))
summary(fit.splines)


##
## Call:
## lm(formula = marathon ~ ns(t, df = 6))
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -13.0028  -2.5722   0.0122   2.1242  21.5681
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    118.447      2.036   58.448 < 2e-16 ***
## ns(t, df = 1)    -6.948      2.688   -2.584  0.011 *
## ns(t, df = 6)2  -28.856      3.416  -8.448 1.16e-13 ***
## ns(t, df = 6)3  -35.081      3.045 -11.522 < 2e-16 ***
## ns(t, df = 6)4  -32.563      2.652 -12.279 < 2e-16 ***
## ns(t, df = 6)5  -64.847      5.322 -12.184 < 2e-16 ***
## ns(t, df = 6)6  -21.002      2.403  -8.741 2.46e-14 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 4.834 on 113 degrees of freedom
## Multiple R-squared:  0.8418, Adjusted R-squared:  0.8334
```

Assignment Project Exam Help

<https://powcoder.com>


Add WeChat powcoder

## Example: Boston marathon winning times



A slightly different type of spline is provided by `splinef`

```
fc <- splinef(marathon)
autoplot(fc)
```



# Assignment Project Exam Help

- Cubic smoothing splines (rather than cubic regression splines).
- Still piecewise cubic, but with many more knots (one at each observation).
- Coefficients constrained to prevent the curve becoming too “wiggly”.
- Degrees of freedom selected automatically.
- Equivalent to ARIMA(0,2,2) and Holt's method.

<https://powcoder.com>

Add WeChat powcoder



1 The linear model with time series

2 Residual diagnostics

3 Some useful predictors for linear models

4 Selecting predictors and forecast evaluation

5 Forecasting with regression

6 Matrix formulation

7 Correlation, causation and forecasting

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

- When there are many predictors, how should we choose which ones to use?
- We need a way of comparing two competing models.

<https://powcoder.com>

Add WeChat powcoder

- When there are many predictors, how should we choose which ones to use?
- We need a way of comparing two competing models.

### What not to do:

- Plot  $y$  against a particular predictor ( $x_j$ ) and if it shows no noticeable relationship, drop it.
- Do a multiple linear regression on all the predictors and disregard all variables whose  $p$  values are greater than 0.05.
- Maximize  $R^2$  or minimize MSE

Computer output for regression will always give the  $R^2$  value. This is a useful summary of the model.

- It is equal to the square of the correlation between  $y$  and  $\hat{y}$ .
- It is often called the “coefficient of determination”
- It can also be calculated as follows:

$$R^2 = \frac{\sum(\hat{y}_t - \bar{y})^2}{\sum(y_t - \bar{y})^2}$$

- It is the proportion of variance accounted for (explained) by the predictors.

## Comparing regression models

However ...

- $R^2$  does not allow for “degrees of freedom”.
- Adding any variable tends to increase the value of  $R^2$ , even if that variable is irrelevant.

<https://powcoder.com>

Add WeChat powcoder

## Comparing regression models

However ...

- $R^2$  does not allow for “degrees of freedom”.
- Adding any variable tends to increase the value of  $R^2$ , even if that variable is irrelevant.

To overcome this problem, we can use *adjusted*  $R^2$ :

$$\bar{R}^2 = 1 - (1 - R^2) \frac{T - 1}{T - k - 1}$$

where  $k$  = no. predictors and  $T$  = no. observations.

## Comparing regression models

However ...

- $R^2$  does not allow for “degrees of freedom”.
- Adding any variable tends to increase the value of  $R^2$ , even if that variable is irrelevant.

To overcome this problem, we can use *adjusted*  $R^2$ :

$$\bar{R}^2 = 1 - (1 - R^2) \frac{T - 1}{T - k - 1}$$

where  $k$  = no. predictors and  $T$  = no. observations.

**Maximizing  $\bar{R}^2$  is equivalent to minimizing  $\hat{\sigma}^2$ .**

$$\hat{\sigma}^2 = \frac{1}{T - k - 1} \sum_{t=1}^T \varepsilon_t^2$$

# Assignment Project Exam Help

### Cross-validation for regression

(Assuming future predictors are known)

- Select one observation for test set and use remaining observations in training set. Compute error on test observation.
- Repeat using each possible observation as the test set.
- Compute accuracy measure over all errors.



### Traditional evaluation



The diagram shows a horizontal timeline with a series of blue dots representing data points. A label 'Training data' is placed above the timeline, pointing to the first 10 dots. A label 'Test data' is placed above the timeline, pointing to the last dot. An arrow labeled 'time' points to the right at the end of the timeline.

Assignment Project Exam Help

<https://powcoder.com>


Add WeChat powcoder

# Cross-validation

## Traditional evaluation

Assignment Project Exam Help

## Leave-one-out cross-validation



# Assignment Project Exam Help

Leave-one-out cross-validation for regression can be carried out using the following steps.


- Remove observation  $t$  from the data set, and fit the model using the remaining data. Then compute the error ( $e_t^* = y_t - \hat{y}_t$ ) for the omitted observation.
- Repeat step 1 for  $t = 1, \dots, T$ .
- Compute the MSE from  $\{e_1^*, \dots, e_T^*\}$ . We shall call this the CV.

The best model is the one with minimum CV.

# Cross-validation

## Five-fold cross-validation

■ 20 observations. 4 test observations per fold




<https://powcoder.com>

Add WeChat powcoder

# Cross-validation


## Five-fold cross-validation

■ 20 observations. 4 test observations per fold



## Ten-fold cross-validation

■ 20 observations. 2 test observations per fold



# Assignment Project Exam Help

### Ten-fold cross-validation

- Randomly split data into 10 parts.
- Select one part for test set, and use remaining parts as training set. Compute accuracy measures on test observations.
- Repeat for each of 10 parts.
- Average over all measures.

<https://powcoder.com>

Add WeChat powcoder

# Assignment Project Exam Help

$$AIC = -2 \log(L) + 2(k + 2)$$

where  $L$  is the likelihood and  $k$  is the number of predictors in the model

<https://powcoder.com>

Add WeChat powcoder

# Assignment Project Exam Help

$$AIC = -2 \log(L) + 2(k + 2)$$

where  $L$  is the likelihood and  $k$  is the number of predictors in the model

<https://powcoder.com>

- This is a *penalized likelihood* approach.
- Minimizing the AIC gives the best model for prediction.
- AIC penalizes terms more heavily than  $R^2$ .
- Minimizing the AIC is asymptotically equivalent to minimizing MSE via leave-one-out cross-validation.

Add WeChat powcoder



# Assignment Project Exam Help

For small values of  $T$ , the AIC tends to select too many predictors, and so a bias-corrected version of the AIC has been developed.

<https://powcoder.com>

$$AIC_C = AIC + \frac{2(k+2)(k+3)}{T-k-3}$$

As with the AIC, the  $AIC_C$  should be minimized.

Add WeChat powcoder

# Assignment Project Exam Help

$$\text{BIC} = -2 \log(L) + (k + 2) \log(T)$$

where  $L$  is the likelihood and  $k$  is the number of predictors in the model.

<https://powcoder.com>

Add WeChat powcoder

# Assignment Project Exam Help

$$\text{BIC} = -2 \log(L) + (k + 2) \log(T)$$

where  $L$  is the likelihood and  $k$  is the number of predictors in the model.

<https://powcoder.com>

- BIC penalizes terms more heavily than AIC
- Also called SBIC and SC.
- Minimizing BIC is asymptotically equivalent to leave- $v$ -out cross-validation when  $v = T[1 - 1/(\log(T) - 1)]$ .

Add WeChat powcoder

### Best subsets regression

- Fit all possible regression models using one or more of the predictors.
- Choose the best model based on one of the measures of predictive ability (CV, AIC, AICc).

Add WeChat powcoder

### Best subsets regression

- Fit all possible regression models using one or more of the predictors.
- Choose the best model based on one of the measures of predictive ability (CV, AIC, AICc).

### Warning!

- If there are a large number of predictors, this is not possible.
- For example, 44 predictors leads to 18 trillion possible models!

## Choosing regression variables

### Backwards stepwise regression

- Start with a model containing all variables.
- Try subtracting one variable at a time. Keep the model if it has lower CV or AICc.
- Iterate until no further improvement.

<https://powcoder.com>

Add WeChat powcoder

# Choosing regression variables

## Backwards stepwise regression

- Start with a model containing all variables.
- Try subtracting one variable at a time. Keep the model if it has lower CV or AICc.
- Iterate until no further improvement.

## Notes

- Stepwise regression is not guaranteed to lead to the best possible model.
- Inference on coefficients of final model will be wrong. Any procedure involving selecting predictors first will invalidate the assumptions behind the p-values.

# Cross-validation

```
tslm(Consumption ~ Income + Production + Unemployment + Savings,  
      data=uschange) %>% CV()
```

```
##           CV           AIC           AICc           BIC           AdjR2  
## 0.1163477 -409.2986258 -408.8313631 -389.5113481 0.7485856
```

```
tslm(Consumption ~ Income + Production + Unemployment,  
      data=uschange) %>% CV()
```

```
##           CV           AIC           AICc           BIC           AdjR2  
## 0.2776928 -243.1635677 -242.8320760 -227.0080246 0.3855438
```

```
tslm(Consumption ~ Income + Production + Savings,  
      data=uschange) %>% CV()
```

```
##           CV           AIC           AICc           BIC           AdjR2  
## 0.1178681 -407.4669279 -407.1354362 -391.3113848 0.7447840
```

```
tslm(Consumption ~ Income + Unemployment + Savings,  
      data=uschange) %>% CV()
```

```
##           CV           AIC           AICc           BIC           AdjR2  
## 0.1160223 -408.0941325 -407.7626408 -391.9385894 0.7456386
```



1 The linear model with time series

2 Residual diagnostics

3 Some useful predictors for linear models

4 Selecting predictors and forecast evaluation

5 Forecasting with regression

6 Matrix formulation

7 Correlation, causation and forecasting

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

- Ex-ante forecasts are made using only information available in advance.

■ For example, ex-ante forecasts for the percentage change in US consumption for quarters following the end of the sample, should only use information that was available up to and including 2016 Q3.

■ These are genuine forecasts made in advance using whatever information is available at the time.

- Require forecasts of predictors

## Ex-ante versus ex-post forecasts

- *Ex post forecasts* are made using later information on the predictors.

- For example, ex-post forecasts of consumption may use the actual observations of the predictors, once these have been observed.

- These are not genuine forecasts.


- useful for studying behaviour of forecasting models.

- trend, seasonal and calendar variables are all known in advance, so these don't need to be forecast. In such cases, there is no difference between ex post and ex ante forecasts.

## Example: Beer production

```
beer2 <- window(ausbeer, start=1992)
fit.beer <- tslm(beer2 ~ trend + season)
fcst <- forecast(fit.beer)
autoplot(fcst) +
  ggtitle("Forecasts of beer production using regression") +
  xlab("Year") + ylab("megalitres")
```

<https://powcoder.com>



## Example: US Consumption

```
fit.consBest <- tslm(
  Consumption ~ Income + Savings + Unemployment,
  data = uschange)
h <- 4
newdata <- data.frame(
  Income = c(1, 1, 1, 1),
  Savings = c(0.5, 0.5, 0.5, 0.5),
  Unemployment = c(0, 0, 0, 0))
fcst.up <- forecast(fit.consBest, newdata = newdata)
newdata <- data.frame(
  Income = rep(-1, h),
  Savings = rep(-0.5, h),
  Unemployment = rep(0, h))
fcst.down <- forecast(fit.consBest, newdata = newdata)
```


Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

## Example: US Consumption

```
autoplot(uschange[, 1]) +  
  ylab("% change in US consumption") +  
  autolayer(feast.up, PI = TRUE, series = "increase") +  
  autolayer(feast.down, PI = TRUE, series = "decrease") +  
  guides(colour = guide_legend(title = "Scenario"))
```



## Building a predictive regression model

- If getting forecasts of predictors is difficult, you can use lagged predictors instead.

$$y_t = \beta_0 + \beta_1 x_{1,t-h} + \dots + \beta_k x_{k,t-h} + \varepsilon_t$$

- A different model for each forecast horizon  $h$ .
- Including lagged values of the predictors does not only make the model operational for easily generating forecasts, it also makes it intuitively appealing.
- For example, the effect of a policy change with the aim of increasing production may not have an instantaneous effect on consumption expenditure. It is most likely that this will happen with a lagging effect.

1 The linear model with time series

2 Residual diagnostics

3 Some useful predictors for linear models

4 Selecting predictors and forecast evaluation

5 Forecasting with regression

6 Matrix formulation

7 Correlation, causation and forecasting

# Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



$$y_t = \beta_0 + \beta_1 X_{1,t} + \beta_2 X_{2,t} + \dots + \beta_k X_{k,t} + \varepsilon_t$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

$$y_t = \beta_0 + \beta_1 x_{1,t} + \beta_2 x_{2,t} + \dots + \beta_k x_{k,t} + \varepsilon_t$$

Let  $\mathbf{y} = (y_1, \dots, y_T)'$ ,  $\boldsymbol{\varepsilon} = (\varepsilon_1, \dots, \varepsilon_T)'$ ,  $\boldsymbol{\beta} = (\beta_0, \beta_1, \dots, \beta_k)'$  and

$$\mathbf{X} = \begin{bmatrix} 1 & x_{1,1} & x_{2,1} & \dots & x_{k,1} \\ 1 & x_{1,2} & x_{2,2} & \dots & x_{k,2} \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & x_{1,T} & x_{2,T} & \dots & x_{k,T} \end{bmatrix}.$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

$$y_t = \beta_0 + \beta_1 x_{1,t} + \beta_2 x_{2,t} + \dots + \beta_k x_{k,t} + \varepsilon_t$$

Let  $\mathbf{y} = (y_1, \dots, y_T)'$ ,  $\boldsymbol{\varepsilon} = (\varepsilon_1, \dots, \varepsilon_T)'$ ,  $\boldsymbol{\beta} = (\beta_0, \beta_1, \dots, \beta_k)'$  and

$$\mathbf{X} = \begin{bmatrix} 1 & x_{1,1} & x_{2,1} & \dots & x_{k,1} \\ 1 & x_{1,2} & x_{2,2} & \dots & x_{k,2} \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & x_{1,T} & x_{2,T} & \dots & x_{k,T} \end{bmatrix}.$$

Then

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\varepsilon}.$$

Least squares estimation

Minimize:  $\|y - X\beta\|^2$

# Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Least squares estimation

Minimize:  $\|y - X\beta\|^2$  ( $y - X\beta$ )

Differentiate wrt  $\beta$  gives

<https://powcoder.com>

Add WeChat powcoder

Least squares estimation

Minimize:  $\|y - X\beta\|^2$  ( $y - X\beta$ )

Differentiate wrt  $\beta$  gives

[\$\beta = \(X'X\)^{-1}X'y\$](https://powcoder.com)

(The “normal equation”.)

Add WeChat powcoder

Least squares estimation

Minimize:  $\|y - X\beta\|^2$

Differentiate wrt  $\beta$  gives

$$\beta = (X'X)^{-1}X'y$$

(The “normal equation”.)

$$\hat{\sigma}^2 = \frac{1}{T - k - 1} (y - X\hat{\beta})'(y - X\hat{\beta})$$

**Note:** If you fall for the dummy variable trap,  $(X'X)$  is a singular matrix.

# Assignment Project Exam Help

If the errors are iid and normally distributed, then

$$\mathbf{y} \sim N(\mathbf{X}\beta, \sigma^2\mathbf{I}).$$

<https://powcoder.com>

Add WeChat powcoder



# Assignment Project Exam Help

If the errors are iid and normally distributed, then

$$\mathbf{y} \sim N(\mathbf{X}\beta, \sigma^2 \mathbf{I}).$$

So the likelihood is

$$L = \frac{1}{\sigma^T (2\pi)^{T/2}} \exp \left( -\frac{1}{2\sigma^2} (\mathbf{y} - \mathbf{X}\beta)' (\mathbf{y} - \mathbf{X}\beta) \right)$$

Add WeChat powcoder

# Assignment Project Exam Help

If the errors are iid and normally distributed, then

$$\mathbf{y} \sim N(\mathbf{X}\beta, \sigma^2 \mathbf{I}).$$

So the likelihood is

$$L = \frac{1}{\sigma^T (2\pi)^{T/2}} \exp \left( -\frac{1}{2\sigma^2} (\mathbf{y} - \mathbf{X}\beta)' (\mathbf{y} - \mathbf{X}\beta) \right)$$

which is maximized when  $(\mathbf{y} - \mathbf{X}\beta)' (\mathbf{y} - \mathbf{X}\beta)$  is minimized

<https://powcoder.com>  
Add WeChat powcoder

# Assignment Project Exam Help

If the errors are iid and normally distributed, then

$$\mathbf{y} \sim N(\mathbf{X}\beta, \sigma^2\mathbf{I}).$$

So the likelihood is

$$L = \frac{1}{\sigma^T(2\pi)^{T/2}} \exp\left(-\frac{1}{2\sigma^2}(\mathbf{y} - \mathbf{X}\beta)'(\mathbf{y} - \mathbf{X}\beta)\right)$$

which is maximized when  $(\mathbf{y} - \mathbf{X}\beta)'(\mathbf{y} - \mathbf{X}\beta)$  is minimized

So MLE = OLS.

## Multiple regression forecasts

### Optimal forecasts

$$\hat{y}^* = E(y^* | \mathbf{y}, \mathbf{X}, \mathbf{x}^*) = \mathbf{x}^* \hat{\beta} = \mathbf{x}^* (\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}'\mathbf{y}$$

where  $\mathbf{x}^*$  is a row vector containing the values of the predictors for the forecasts (in the same format as  $\mathbf{X}$ ).

<https://powcoder.com>

Add WeChat powcoder

## Multiple regression forecasts

### Optimal forecasts

$$\hat{y}^* = E(y^* | \mathbf{X}, \mathbf{x}^*) = \mathbf{x}^* \hat{\beta} = \mathbf{x}^* (\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}'\mathbf{y}$$

where  $\mathbf{x}^*$  is a row vector containing the values of the predictors for the forecasts (in the same format as  $\mathbf{X}$ ).

### Forecast variance

$$\text{Var}(y^* | \mathbf{X}, \mathbf{x}^*) = \sigma^2 \left[ 1 + \mathbf{x}^* (\mathbf{X}'\mathbf{X})^{-1} (\mathbf{x}^*)' \right]$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

# Multiple regression forecasts

## Optimal forecasts

$$\hat{y}^* = E(y^* | \mathbf{X}, \mathbf{x}^*) = \mathbf{x}^* \hat{\beta} = \mathbf{x}^* (\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}'\mathbf{y}$$

where  $\mathbf{x}^*$  is a row vector containing the values of the predictors for the forecasts (in the same format as  $\mathbf{X}$ ).

## Forecast variance

$$\text{Var}(y^* | \mathbf{X}, \mathbf{x}^*) = \sigma^2 \left[ 1 + \mathbf{x}^* (\mathbf{X}'\mathbf{X})^{-1} (\mathbf{x}^*)' \right]$$

- This ignores any errors in  $\mathbf{x}^*$ .
- 95% prediction intervals assuming normal errors:

$$\hat{y}^* \pm 1.96 \sqrt{\text{Var}(y^* | \mathbf{X}, \mathbf{x}^*)}$$

Fitted values

Assignment Project Exam Help

where  $\hat{y} = X(X'X)^{-1}X'y = Hy$  is the “hat matrix”.

<https://powcoder.com>

Add WeChat powcoder

## Multiple regression forecasts

### Fitted values

$$\hat{y} = X\hat{\beta} = X(X'X)^{-1}X'y = Hy$$

where  $H = X(X'X)^{-1}X'$  is the “hat matrix”.

### Leave one-out residuals

Let  $h_1, \dots, h_T$  be the diagonal values of  $H$ , then the cross-validation statistic is

$$CV = \frac{1}{T} \sum_{t=1}^T [e_t / (1 - h_t)]^2,$$

where  $e_t$  is the residual obtained from fitting the model to all  $T$  observations.



1 The linear model with time series

2 Residual diagnostics

3 Some useful predictors for linear models

4 Selecting predictors and forecast evaluation

5 Forecasting with regression

6 Matrix formulation

7 Correlation, causation and forecasting

# Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

# Assignment Project Exam Help

- When  $x$  is useful for predicting  $y$  it is not necessarily causing  $y$ .
- e.g., predict number of drownings  $y$  using number of ice creams sold  $x$
- Correlations are useful for forecasting, even when there is no causality.
- Better models usually involve causal relationships (e.g., temperature  $x$  and people  $z$  to predict drownings  $y$ ).

# Assignment Project Exam Help

In regression analysis, multicollinearity occurs when

- Two predictors are highly correlated (i.e., the correlation between them is close to  $\pm 1$ ).
- A linear combination of some of the predictors is highly correlated with another predictor.
- A linear combination of one subset of predictors is highly correlated with a linear combination of another subset of predictors.

<https://powcoder.com>

Add WeChat powcoder

If multicollinearity exists...

- the numerical estimates of coefficients may be wrong (worse in Excel than in a statistics package)
- don't rely on the p-values to determine significance.
- there is no problem with model *predictions* provided the predictors used for forecasting are within the range used for fitting.
- omitting variables can help.
- combining variables can help.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

# Assignment Project Exam Help

Things to watch for

- *Outliers*: observations that produce large residuals.
- *Influential observations*: removing them would markedly change the coefficients. (Often outliers in the  $x$  variable).
- *Lurking variable*: a predictor not included in the regression but which has an important effect on the response.
- Points should not normally be removed without a good explanation of why they are different.

<https://powcoder.com>

Add WeChat powcoder