



MONASH
University

MONASH
BUSINESS
SCHOOL

ETW3420:

Assignment Project Exam Help

Principles of

Forecasting and

Applications

Add WeChat powcoder

Topic 6: ARIMA Models

Dr. Jason Ng

1 Introduction

2 Stationarity and differencing

3 Non-seasonal ARIMA models

4 Model identification

5 Estimation and order selection

6 ARIMA modeling in R

7 Forecasting

8 Seasonal ARIMA models

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

- ARIMA models provide another approach to time series forecasting.
- Exponential smoothing and ARIMA models are the two most widely used approaches to time series forecasting, and provide complementary approaches to the problem.
- While exponential smoothing models are based on a description of the trend and seasonality in the data, ARIMA models aim to describe the autocorrelations in the data.
- Before we introduce ARIMA models, we must first discuss the concept of stationarity and the technique of differencing time series.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

1 Introduction

2 Stationarity and differencing

3 Non-seasonal ARIMA models

4 Model identification

5 Estimation and order selection

6 ARIMA modeling in R

7 Forecasting

8 Seasonal ARIMA models

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

Definition

If $\{y_t\}$ is a stationary time series, then for all s , the distribution of (y_t, \dots, y_{t+s}) does not depend on t .

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

Definition

If $\{y_t\}$ is a stationary time series, then for all s , the distribution of (y_t, \dots, y_{t+s}) does not depend on t .

A **stationary series** is:

- roughly horizontal
- constant variance
- no patterns predictable in the long-term

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

Statistical Properties of Stationary Series

Statistically, a time series $\{y_t\}$ is stationary if:

<https://powcoder.com>


$$E(y_t) = \mu < \infty \text{ for all } t$$

$$\text{Var}(y_t) = \gamma_0 < \infty \text{ for all } t$$


$$\text{cov}(y_t, y_{t-j}) = \gamma_j < \infty \text{ for all } t$$

Add WeChat powcoder


Stationary?



Stationary?




Stationary?




Stationary?

Sales of new one-family houses, USA




Stationary?

Price of a dozen eggs in 1993 dollars




Stationary?



Stationary?

Australian quarterly beer production



Assignment Project Exam Help

Definition

If $\{y_t\}$ is a stationary time series, then for all s , the distribution of (y_t, \dots, y_{t+s}) does not depend on t .

Add WeChat powcoder

Assignment Project Exam Help

Definition

If $\{y_t\}$ is a stationary time series, then for all s , the distribution of (y_t, \dots, y_{t+s}) does not depend on t .


- Transformations help to **stabilize the variance**.
- For ARIMA modelling, we also need to **stabilize the mean**.

Assignment Project Exam Help


Identifying non-stationary series

- Time plot.
- The ACF of stationary data drops to zero relatively quickly
- The ACF of non-stationary data decreases slowly.
- For non-stationary data, the value of r_1 is often large and positive.


Example: Dow-Jones index




Example: Dow-Jones index



Example: Dow-Jones index



Example: Dow-Jones index



Assignment Project Exam Help

- Differencing helps to **stabilize the mean**.
- The differenced series is the *change* between each observation in the original series. $y'_t = y_t - y_{t-1}$.
- The differenced series will have only $T - 1$ values since it is not possible to calculate a difference y'_1 for the first observation.

<https://powcoder.com>

Add WeChat powcoder

Occasionally the differenced data will not appear stationary and it may be necessary to difference the data a second time:

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Occasionally the differenced data will not appear stationary and it may be necessary to difference the data a second time:

$$\begin{aligned}y_t'' &= y_t' - y_{t-1}' \\&= (y_t - y_{t-1}) - (y_{t-1} - y_{t-2}) \\&= y_t - 2y_{t-1} + y_{t-2}.\end{aligned}$$

Add WeChat powcoder

Occasionally the differenced data will not appear stationary and it may be necessary to difference the data a second time:

$$\begin{aligned}y_t'' &= y_t' - y_{t-1}' \\&= (y_t - y_{t-1}) - (y_{t-1} - y_{t-2}) \\&= y_t - 2y_{t-1} + y_{t-2}.\end{aligned}$$

- y_t' will have $T-2$ values

- In practice, it is almost never necessary to go beyond second-order differences.

Assignment Project Exam Help

A seasonal difference is the difference between an observation and the corresponding observation from the previous year.

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

A seasonal difference is the difference between an observation and the corresponding observation from the previous year.

$y'_t = y_t - y_{t-m}$
<https://powcoder.com>

where m = number of seasons.

Add WeChat powcoder

Assignment Project Exam Help

A seasonal difference is the difference between an observation and the corresponding observation from the previous year.

$y'_t = y_t - y_{t-m}$
<https://powcoder.com>

where m = number of seasons.

- For monthly data $m = 12$.
- For quarterly data $m = 4$.

Add WeChat powcoder


Electricity production

```
usmelec %>% autoplot()
```

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder




Electricity production

```
usmelec %>% log() %>% autoplot()
```

Assignment Project Exam Help


<https://powcoder.com>

Add WeChat powcoder




Electricity production

```
usmelec %>% log() %>% diff(lag=12) %>%  
autoplot()
```



Electricity production

```
usmelec %>% log() %>% diff(lag=12) %>%  
diff(lag=1) %>% autoplot()
```



Electricity production

- Seasonally differenced series is closer to being stationary.
- Remaining non-stationarity can be removed with further first difference.

If $y'_t = y_t - y_{t-12}$ denotes seasonally differenced series, then twice-differenced series is given by:

$$y_t^* = y'_t - y'_{t-1}$$

$$\begin{aligned} &= (y_t - y_{t-12}) - (y_{t-1} - y_{t-13}) \\ &= y_t - y_{t-1} - y_{t-12} + y_{t-13} . \end{aligned}$$

When both seasonal and first differences are applied...

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

When both seasonal and first differences are applied...

- it makes no difference which is done first—the result will be the same.
- If seasonality is strong, we recommend that seasonal differencing be done first because sometimes the resulting series will be stationary and there will be no need for further first difference.

When both seasonal and first differences are applied...

- it makes no difference which is done first—the result will be the same.
- If seasonality is strong, we recommend that seasonal differencing be done first because sometimes the resulting series will be stationary and there will be no need for further first difference.

It is important that if differencing is used, the differences are interpretable.

Assignment Project Exam Help

- first differences are the change between **one observation and the next**;
- seasonal differences are the change between **one year to the next**.

<https://powcoder.com>
Add WeChat powcoder

Assignment Project Exam Help

- first differences are the change between **one observation and the next**;
- seasonal differences are the change between **one year to the next**.

But taking lag 3 differences for yearly data, for example, results in a model which cannot be sensibly interpreted.

Assignment Project Exam Help

Statistical tests to determine the required order of differencing.

- 1 Augmented Dickey Fuller test: null hypothesis is that the data are non-stationary.
- 2 Kwiatkowski-Phillips-Schmidt-Shin (KPSS) test: null hypothesis is that the data are stationary. (Will use this for our analysis)
- 3 Other tests available for seasonal data.

```
library(urca)
summary(ur.kpss(goog))
```

Assignment Project Exam Help

```
##
## #####
## # KPSS Unit Root Test #
## #####
##
## Test is of type: mu with 7 lags.
##
## Value of test-statistic is: 10.7203
##
## Critical value for a significance level of:
##          10pct  5pct  2.5pct  1pct
## critical values 0.347 0.463  0.574 0.739
```

<https://powcoder.com>

Add WeChat powcoder

- Test statistic is much larger than the 1% critical value, indicating that the null hypothesis is rejected, i.e. data are not stationary.


```
goog %>% diff() %>% ur.kpss() %>% summary()
```

Assignment Project Exam Help

```
## #####
```

```
## # KPSS Unit Root Test #
```

```
## #####
```

```
## https://powcoder.com
```

```
## Test is of type: m1 with 7 lags.
```

```
##
```

```
## Value of test-statistic is: 0.0324
```

```
## Add WeChat powcoder
```

```
## Critical value for a significance level of:
```

```
##          10pct  5pct 2.5pct  1pct
```

```
## critical values 0.347 0.463 0.574 0.739
```

- Conclude that the differenced data are stationary.

Assignment Project Exam Help

This process of using a sequence of KPSS tests to determine the appropriate number of first differences is carried out by the function `ndiffs()`.

<https://powcoder.com>

```
ndiffs(goog)
```

Add WeChat powcoder

```
## [1] 1
```

Automatic selection of seasonal difference

- STL decomposition: $y_t = T_t + S_t + R_t$
- Seasonal strength $F_s = \max(0, 1 - \frac{\text{Var}(R_t)}{\text{Var}(T_t + R_t)})$
- If $F_s > 0.64$, do one seasonal difference. Otherwise, no seasonal difference required.
- Use the `nsdiffs()` function to test.

<https://powcoder.com>

Add WeChat powcoder

Automatic selection of seasonal difference

- STL decomposition: $y_t = T_t + S_t + R_t$
- Seasonal strength $F_s = \max(0, 1 - \frac{\text{Var}(R_t)}{\text{Var}(T_t + R_t)})$
- If $F_s > 0.64$, do one seasonal difference. Otherwise, no seasonal difference required.
- Use the `nsdiffs()` function to test.

```
usmelec %>% log() %>% nsdiffs()
```

```
## [1] 1
```

```
usmelec %>% log() %>% diff(lag=12) %>% ndiffs()
```

```
## [1] 1
```

- These functions suggest performing both a seasonal and first difference

Assignment Project Exam Help

<https://powcoder.com>

For the visitors series, find an appropriate differencing (after transformation if necessary) to obtain stationary data.

Add WeChat powcoder

A very useful notational device is the backward shift operator, B , which is used as follows:

$$By_t = y_{t-1}.$$

<https://powcoder.com>

Add WeChat powcoder

A very useful notational device is the backward shift operator, B , which is used as follows:

$$By_t = y_{t-1} .$$

In other words, B , operating on y_t , has the effect of shifting the data back one period.

Add WeChat powcoder

A very useful notational device is the backward shift operator, B , which is used as follows:

$$By_t = y_{t-1}.$$

In other words, B , operating on y_t , has the effect of shifting the data back one period. Two applications of B to y_t shifts the data back two periods:

$$B^2 y_t = y_{t-2}.$$

A very useful notational device is the backshift operator, B , which is used as follows:

$$By_t = y_{t-1}.$$

In other words, B , operating on y_t , has the effect of shifting the data back one period. Two applications of B to y_t shifts the data back two periods:

$$B^2 y_t = y_{t-2}.$$

For monthly data, if we wish to shift attention to “the same month last year,” then B^{12} is used, and the notation is $B^{12}y_t = y_{t-12}$.

Assignment Project Exam Help

The backward shift operator is convenient for describing the process of *differencing*.

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

The backward shift operator is convenient for describing the process of *differencing*. A first difference can be written as

$$y'_t \equiv y_t - y_{t-1} \equiv y_t - By_t \equiv (1 - B)y_t$$

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

The backward shift operator is convenient for describing the process of *differencing*. A first difference can be written as

$$y'_t \equiv y_t - y_{t-1} \equiv y_t - By_t \equiv (1 - B)y_t$$

Note that a first difference is represented by $(1 - B)$.

Add WeChat powcoder

Assignment Project Exam Help

The backward shift operator is convenient for describing the process of *differencing*. A first difference can be written as

$$y'_t = y_t - y_{t-1} = y_t - By_t = (1 - B)y_t$$

Note that a first difference is represented by $(1 - B)$.

Similarly, if second-order differences (i.e., first differences of first differences) have to be computed, then

$$y''_t = y_t - 2y_{t-1} + y_{t-2} = (1 - B)^2 y_t.$$

Assignment Project Exam Help

- Second-order difference is denoted $(1 - B)^2$
- Second-order difference is not the same as a second difference, which would be denoted $1 - B^2$;
- In general, a d th-order difference can be written as

$$(1 - B)^d y_t.$$

- A seasonal difference followed by a first difference can be written as

$$(1 - B)(1 - B^m)y_t.$$

Assignment Project Exam Help

The “backshift” notation is convenient because the terms can be multiplied together to see the combined effect.

<https://powcoder.com>

$$(1 - B)(1 - B^m)y_t = (1 - B - B^m + B^{m+1})y_t$$

Add WeChat $y_t - y_{t-1} - y_{t-m} + y_{t-m-1}$ powcoder

Assignment Project Exam Help

The “backshift” notation is convenient because the terms can be multiplied together to see the combined effect.

<https://powcoder.com>

$$(1 - B)(1 - B^m)y_t = (1 - B - B^m + B^{m+1})y_t$$

Add WeChat $y_t - y_{t-1} - y_{t-m} + y_{t-m-1}$ powcoder

For monthly data, $m = 12$ and we obtain the same result as earlier.

1 Introduction

2 Stationarity and differencing

3 Non-seasonal ARIMA models

4 Model identification

5 Estimation and order selection

6 ARIMA modeling in R

7 Forecasting

8 Seasonal ARIMA models

Assignment Project Exam Help



<https://powcoder.com>

Add WeChat powcoder

Autoregressive models

Autoregressive (AR) models:

$y_t = \phi_1 y_{t-1} + \phi_2 y_{t-2} + \dots + \phi_p y_{t-p} + \varepsilon_t$,
where ε_t is white noise. This is a multiple regression with **lagged values** of y_t as predictors.




AR(1) model

$$y_t = 2 - 0.8y_{t-1} + \varepsilon_t$$

$$\varepsilon_t \sim N(0, 1), \quad T = 100$$

AR(1)



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

$$y_t = c + \phi_1 y_{t-1} + \varepsilon_t$$


- When $\phi_1 = 0$, y_t is equivalent to WN
- When $\phi_1 = 1$ and $c = 0$, y_t is equivalent to a RW
- When $\phi_1 = 1$ and $c \neq 0$, y_t is equivalent to a RW with drift
- When $\phi_1 < 0$, y_t tends to oscillate between positive and negative values.

AR(2) model

$$y_t = 8 + 1.3y_{t-1} - 0.7y_{t-2} + \varepsilon_t$$

$$\varepsilon_t \sim N(0, 1), \quad T = 100$$

AR(2)



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Stationarity conditions

We normally restrict autoregressive models to stationary data, and then some constraints on the values of the parameters are required.

Assignment Project Exam Help

General condition for stationarity

Complex roots of $1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p$ lie outside the unit circle on the complex plane.

<https://powcoder.com>
Add WeChat powcoder

Stationarity conditions

We normally restrict autoregressive models to stationary data, and then some constraints on the values of the parameters are required.

General condition for stationarity

Complex roots of $1 - \phi_1 z - \phi_2 z^2 - \dots - \phi_p z^p$ lie outside the unit circle on the complex plane.



- For $p = 1$: $-1 < \phi_1 < 1$.
- For $p = 2$:
 $-1 < \phi_2 < 1$ $\phi_2 + \phi_1 < 1$ $\phi_2 - \phi_1 < 1$.
- More complicated conditions hold for $p \geq 3$.
- Estimation software takes care of this.

Moving Average (MA) models

Moving Average (MA) models:

$y_t = \mu + \epsilon_t + \theta_1 \epsilon_{t-1} + \theta_2 \epsilon_{t-2} + \dots + \theta_q \epsilon_{t-q}$,
where ϵ_t is white noise. This is a multiple regression with **past errors** as predictors. Don't confuse this with moving average smoothing.

<https://powcoder.com>




MA(1) model

$$y_t = 20 + \varepsilon_t + 0.8\varepsilon_{t-1}$$

$\varepsilon_t \sim N(0, 1), T = 100$

MA(1)



Assignment Project Exam Help

<https://powcoder.com>


Add WeChat powcoder

MA(2) model

$$y_t = \varepsilon_t - \varepsilon_{t-1} + 0.8\varepsilon_{t-2}$$

$\varepsilon_t \sim N(0, 1), T = 100$

MA(2)



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

It is possible to write any stationary AR(p) process as an MA(∞) process.

Example: AR(1).

$$\begin{aligned} y_t &= \phi_1 y_{t-1} + \varepsilon_t \\ &= \phi_1(\phi_1 y_{t-2} + \varepsilon_{t-1}) + \varepsilon_t \end{aligned}$$

$$= \phi_1^2 y_{t-2} + \phi_1 \varepsilon_{t-1} + \varepsilon_t$$

$$= \phi_1^3 y_{t-3} + \phi_1^2 \varepsilon_{t-2} + \phi_1 \varepsilon_{t-1} + \varepsilon_t$$

...

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

It is possible to write any stationary AR(p) process as an MA(∞) process.

Example: AR(1).

$$\begin{aligned}y_t &= \phi_1 y_{t-1} + \varepsilon_t \\&= \phi_1(\phi_1 y_{t-2} + \varepsilon_{t-1}) + \varepsilon_t\end{aligned}$$

$$= \phi_1^2 y_{t-2} + \phi_1 \varepsilon_{t-1} + \varepsilon_t$$

$$= \phi_1^3 y_{t-3} + \phi_1^2 \varepsilon_{t-2} + \phi_1 \varepsilon_{t-1} + \varepsilon_t$$

...

Provided $-1 < \phi_1 < 1$:

$$y_t = \varepsilon_t + \phi_1 \varepsilon_{t-1} + \phi_1^2 \varepsilon_{t-2} + \phi_1^3 \varepsilon_{t-3} + \dots$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

- Any $MA(q)$ process can be written as an $AR(\infty)$ process if we impose some constraints on the MA parameters.
- Then the MA model is called “invertible”.
- Invertible models have some mathematical properties that make them easier to use in practice.
- Invertibility of an ARMA model is equivalent to forecastability of an ETS model.

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

General condition for invertibility

Complex roots of $1 + \theta_1 z + \theta_2 z^2 + \dots + \theta_q z^q$ lie outside the unit circle on the complex plane.

<https://powcoder.com>

Add WeChat powcoder

General condition for invertibility

Complex roots of $1 + \theta_1 z + \theta_2 z^2 + \dots + \theta_q z^q$ lie outside the unit circle on the complex plane.

■ For $q = 1$: $-1 < \theta_1 < 1$.

■ For $q = 2$:

$$-1 < \theta_2 < 1, \quad \theta_2 + \theta_1 > -1, \quad \theta_1 - \theta_2 < 1$$

■ More complicated conditions hold for $q \geq 3$.

■ Estimation software takes care of this.

Assignment Project Exam Help

Autoregressive Moving Average models:

$$y_t = c + \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p}$$

$$+ \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q} + \varepsilon_t$$

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

Autoregressive Moving Average models:

$$y_t = c + \phi_1 y_{t-1} + \cdots + \phi_p y_{t-p}$$

$$+ \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q} + \varepsilon_t$$

<https://powcoder.com>

- Predictors include both **lagged values of y_t and lagged errors.**

- Conditions on AR coefficients ensure stationarity.

- Conditions on MA coefficients ensure invertibility.

Add WeChat powcoder

Assignment Project Exam Help

Autoregressive Integrated Moving Average models:

$$y'_t = c + \phi_1 y'_{t-1} + \dots + \phi_p y'_{t-p} + \theta_1 \varepsilon_{t-1} + \dots + \theta_q \varepsilon_{t-q} + \varepsilon_t.$$

- Combine ARIMA model with **differencing**

- $y_t = (1 - B)^d y_t$ follows an ARIMA model.

Autoregressive Integrated Moving Average models

ARIMA(p, d, q) model

AR: p = order of the autoregressive part

I: d = degree of first differencing involved

MA: q = order of the moving average part.

- White noise model: ARIMA(0,0,0)
- Random walk: ARIMA(0,1,0) with no constant
- Random walk with drift: ARIMA(0,1,0) with const.
- AR(p): ARIMA($p,0,0$)
- MA(q): ARIMA(0,0, q)

- ARMA model:

$$y_t = c + \phi_1 B y_t + \dots + \phi_p B^p y_t + \varepsilon_t + \theta_1 B \varepsilon_t + \dots + \theta_q B^q \varepsilon_t$$

or $(1 - \phi_1 B - \dots - \phi_p B^p) y_t = c + (1 + \theta_1 B + \dots + \theta_q B^q) \varepsilon_t$

- ARIMA(1,1,1) model:

$$(1 - \phi_1 B) (1 - B) y_t = c + (1 + \theta_1 B) \varepsilon_t$$

AR(1) First difference MA(1)

Add WeChat powcoder

Backshift notation for ARIMA

- ARMA model:

$$y_t = c + \phi_1 B y_t + \dots + \phi_p B^p y_t + \varepsilon_t + \theta_1 B \varepsilon_t + \dots + \theta_q B^q \varepsilon_t$$

or $(1 - \phi_1 B - \dots - \phi_p B^p) y_t = c + (1 + \theta_1 B + \dots + \theta_q B^q) \varepsilon_t$

- ARIMA(1,1,1) model:

$$(1 - \phi_1 B) (1 - B) y_t = c + (1 + \theta_1 B) \varepsilon_t$$

AR(1) First difference MA(1)

Add WeChat powcoder

Written out:

$$y_t = c + y_{t-1} + \phi_1 y_{t-1} - \phi_1 y_{t-2} + \theta_1 \varepsilon_{t-1} + \varepsilon_t$$

Intercept form

$$(1 - \phi_1 B - \dots - \phi_p B^p) y'_t = c + (1 + \theta_1 B + \dots + \theta_q B^q) \varepsilon_t$$

Mean form (Parameterisation in R)

$$(1 - \phi_1 B - \dots - \phi_p B^p) (y_t - \mu) = (1 + \theta_1 B + \dots + \theta_q B^q) \varepsilon_t$$

- $y'_t = (1 - B)^d y_t$
- μ is the mean of y'_t .
- Set $c = \mu(1 - \phi_1 - \dots - \phi_p)$ to convert from mean form to intercept form


US personal consumption

- Figure below shows quarterly percentage changes in US consumption expenditure.

- Although it is a quarterly series, there does NOT appear to be a seasonal pattern.

- Hence, fit a non-seasonal ARIMA model.

US consumption



US personal consumption

```
(fit <- auto.arima(uschange[, "Consumption"])
```

```
## Series: uschange[, "Consumption"]
```

```
## ARIMA(2,0,2) with non-zero mean
```

```
##
```

```
## Coefficients:
```

```
##          ar1          ar2          ma1          ma2          mean
```

```
##      1.3908    -0.5313    -1.1800     0.5584     0.7463
```

```
## s.e.   0.2553     0.2078     0.2381     0.1403     0.0845
```

```
##
```

```
## sigma^2 estimated as 0.3511: log likelihood=-165.14
```

```
## AIC=342.28   AICc=342.75   BIC=361.67
```

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

US personal consumption

```
(fit <- auto.arima(uschange[, "Consumption"]))
```

```
## Series: uschange[, "Consumption"]  
## ARIMA(2,0,2) with non-zero mean  
##  
## Coefficients:  
##          ar1          ar2          ma1          ma2          mean  
##      1.3908    -0.5813    -1.1800    0.5584    0.7463  
## s.e.   0.2553     0.2078     0.2381    0.1403    0.0845  
##  
## sigma^2 estimated as 0.351: log likelihood=-165.14  
## AIC=342.28   AICc=342.75   BIC=361.67  
##
```

ARIMA(2,0,2) model:


$$y_t = c + 1.391y_{t-1} - 0.581y_{t-2} - 1.180\varepsilon_{t-1} + 0.558\varepsilon_{t-2} + \varepsilon_t,$$

where $c = 0.746 \times (1 - 1.391 + 0.581) = 0.142$ and ε_t is white noise with a standard deviation of $0.593 = \sqrt{0.351}$.

US personal consumption

```
fit %>% forecast(h=10) %>% autoplot(include=80)
```

Forecasts from ARIMA(2,0,2) with non-zero mean



Understanding ARIMA models

- If $c = 0$ and $d = 0$, the long-term forecasts will go to zero.
- If $c = 0$ and $d = 1$, the long-term forecasts will go to a non-zero constant determined by the last few observations.
- If $c = 0$ and $d = 2$, the long-term forecasts will follow a straight line with intercept and slope determined by the last few observations.
- If $c \neq 0$ and $d = 0$, the long-term forecasts will go to the mean of the data.
- If $c \neq 0$ and $d = 1$, the long-term forecasts will follow a straight line with slope equal to the mean of the differenced data.
- If $c \neq 0$ and $d = 2$, the long-term forecasts will follow a quadratic trend.

Understanding ARIMA models

Forecast variance and d

- The higher the value of d , the more rapidly the prediction intervals increase in size.
- For $d = 0$, the long-term forecast standard deviation will go to the standard deviation of the historical data.

Cyclic behaviour

- For cyclic forecasts, $p \geq 2$ and some restrictions on coefficients are required.
- If $p = 2$, we need $\phi_1^2 + 4\phi_2 < 0$. Then average cycle of length

$$(2\pi) / \left[\arccos(-\phi_1(1 - \phi_2)/(4\phi_2)) \right].$$

1 Introduction

2 Stationarity and differencing

3 Non-seasonal ARIMA models

4 Model identification

5 Estimation and order selection

6 ARIMA modeling in R

7 Forecasting

8 Seasonal ARIMA models

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

- Question: How do we identify an ARIMA model? That is, how do we identify the values of p, d, q ?

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

- Question: How do we identify an ARIMA model? That is, how do we identify the values of p, d, q ?

<https://powcoder.com>

- Rely on the sample ACF and sample PACF plots to determine appropriate values for p and q .
- To do this, will need to know the theoretical ACF and PACF of some common AR and MA models.

Add WeChat powcoder

Partial autocorrelations

Partial autocorrelations measure relationship

between y_t and y_{t-k} , when the effects of other time lags —

1, 2, 3, ..., $k-1$ — are removed.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Partial autocorrelations

Partial autocorrelations measure relationship

between y_t and y_{t-k} , when the effects of other time lags —
1, 2, 3, ..., $k-1$ — are removed.

α_k = k th partial autocorrelation coefficient

= equal to the estimate of ϕ_k in regression

$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \cdots + \phi_k y_{t-k}.$$

Add WeChat powcoder

Partial autocorrelations

Partial autocorrelations measure relationship

between y_t and y_{t-k} , when the effects of other time lags —

1, 2, 3, ..., $k-1$ — are removed.


α_k = k th partial autocorrelation coefficient

= equal to the estimate of ϕ_k in regression


$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \cdots + \phi_k y_{t-k}.$$

- Varying number of terms on RHS gives α_k for different values of k .
- There are more efficient ways of calculating α_k .
- $\alpha_1 = \rho_1$
- same critical values of $\pm 1.96/\sqrt{T}$ as for ACF.

Example: US consumption



Example: US consumption






Assignment Project Exam Help

$$\begin{aligned} \rho_k &= \phi_1^k & \text{for } k = 1, 2, \dots; \\ \alpha_1 &= \phi_1 & \alpha_k = 0 & \text{for } k = 2, 3, \dots \end{aligned}$$

So we have an AR(1) model when

- autocorrelation exponentially decay
- there is a single significant partial autocorrelation.

AR(1): Simulation Example



Assignment Project Exam Help


AR(p)

- ACF dies out in an exponential or damped sine-wave manner
- PACF has all zero spikes beyond the p th spike

So we have an AR(p) model when

- the ACF is exponentially decaying or sinusoidal
- there is a significant spike at lag p in PACF, but none beyond p

AR(2): Simulation Example



Assignment Project Exam Help

$\rho_1 = \theta_1$ $\rho_k = 0$ for $k = 2, 3, \dots$;
 $\alpha_k = -(-\theta_1)^k$


<https://powcoder.com>

So we have an MA(1) model when

- the PACF is exponentially decaying and
- there is a single significant spike in ACF

Add WeChat powcoder

MA(1): Simulation Experiment



Assignment Project Exam Help


MA(q)

- PACF dies out in an exponential or damped sine-wave manner
- ACF has all zero spikes beyond the q th spike

So we have an MA(q) model when

- the PACF is exponentially decaying or sinusoidal
- there is a significant spike at lag q in ACF, but none beyond q


MA(2): Simulation Example




Model Identification: Some caveats

- If the data are from an $ARIMA(p, d, 0)$ or $ARIMA(0, d, q)$ model, then the ACF and PACF plots can be helpful in determining the value of p or q .
- If both p and q are positive, then the plots do not help in finding suitable values of p and q .
- It is common practice to identify two sets of models: 1 set of models consider $p = 0$ and $q > 0$ while the other set of models consider $p > 0$ and $q = 0$. The models can then be modified subsequently to have both p and $q > 0$.
- This step of model identification is largely a trial and error process. Known as Box-Jenkins methodology.

Example: US consumption



Example: US consumption



1 Introduction

2 Stationarity and differencing

3 Non-seasonal ARIMA models

4 Model identification

5 Estimation and order selection

6 ARIMA modeling in R

7 Forecasting

8 Seasonal ARIMA models

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Maximum likelihood estimation

Having identified the model order, we need to estimate the parameters $c, \phi_1, \dots, \phi_p, \theta_1, \dots, \theta_q$.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Maximum likelihood estimation

Having identified the model order, we need to estimate the parameters $c, \phi_1, \dots, \phi_p, \theta_1, \dots, \theta_q$.

- Uses maximum likelihood estimation (MLE) to estimate the ARIMA model. This technique finds the values of the parameters which **maximize the probability** of obtaining the data we have observed.
- For ARIMA models, MLE is very similar to least squares estimation obtained by minimizing

$$\sum_{t=1}^T e_t^2$$

- The `Arma()` command allows CLS or MLE estimation.
- Different software will give different estimates due to different optimization algorithms.
- In practice, R will try to maximise the *log likelihood* of the

- The AIC is useful for determining the order of an ARIMA model.

Alaika's Information Criterion (AIC):

$$AIC = -2 \log(L) + 2(p + q + k + 1),$$

where L is the likelihood of the data,

$k = 1$ if $c > 0$ and $k = 0$ if $c = 0$.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

- The AIC is useful for determining the order of an ARIMA model.

Alkaike's information criterion (AIC):

$$AIC = -2 \log(L) + 2(p + q + k + 1),$$

where L is the likelihood of the data,

$k = 1$ if $c \neq 0$ and $k = 0$ if $c = 0$.

Corrected AIC:

$$AIC_c = AIC + \frac{2(p+q+k+1)(p+q+k+2)}{n-p-q-2}.$$

- The AIC is useful for determining the order of an ARIMA model.

Alaika's Information Criterion (AIC):

$$AIC = -2 \log(L) + 2(p + q + k + 1),$$

where L is the likelihood of the data,

$k = 1$ if $c \neq 0$ and $k = 0$ if $c = 0$.

Corrected AIC:

$$AIC_c = AIC + \frac{2(p+q+k+1)(p+q+k+2)}{T-p-q-2}.$$

Bayesian Information Criterion:

$$BIC = AIC + [\log(T) - 2](p + q + k - 1).$$

Information criteria

- The AIC is useful for determining the order of an ARIMA model.

Alaika's Information Criterion (AIC):

$$\text{AIC} = -2 \log(L) + 2(p + q + k + 1),$$

where L is the likelihood of the data,

$k = 1$ if $c \neq 0$ and $k = 0$ if $c = 0$.

Corrected AIC:

$$\text{AICc} = \text{AIC} + \frac{2(p+q+k+1)(p+q+k+2)}{T-p-q-2}.$$

Bayesian Information Criterion:

$$\text{BIC} = \text{AIC} + [\log(T) - 2](p + q + k - 1).$$

Good models are obtained by minimizing either the AIC, AICc or BIC. Our preference is to use the AICc.

1 Introduction

2 Stationarity and differencing

3 Non-seasonal ARIMA models

4 Model identification

5 Estimation and order selection

6 ARIMA modeling in R

7 Forecasting

8 Seasonal ARIMA models

Assignment Project Exam Help

<https://powcoder.com>


Add WeChat powcoder

Modelling procedure with Arima

- 1 Plot the data. Identify any unusual observations.
- 2 If necessary, transform the data (using a Box-Cox transformation) to stabilize the variance.
- 3 If the data are non-stationary: take first differences of the data until the data are stationary.
- 4 Examine the ACF/PACF: Is an AR(p) or MA(q) model appropriate?
- 5 Try your chosen model(s), and use the AICc to search for a better model.
- 6 Check the residuals from your chosen model by plotting the ACF of the residuals, and doing a portmanteau test of the residuals. If they do not look like white noise, try a modified model.
- 7 Once the residuals look like white noise, calculate forecasts.

Seasonally adjusted electrical equipment

```
eadj <- seasadj(stl(elecequip, s.window="periodic"))  
autoplot(eadj) + xlab("Year") +  
  ylab("Seasonally adjusted new orders index")
```



Assignment Project Exam Help

1 Time plot shows sudden changes, particularly big drop in 2008/2009 due to global economic environment. Otherwise nothing unusual and no need for data adjustments.

2 No evidence of changing variance, so no Box-Cox transformation.




3 Data are clearly non-stationary, so we take first differences.

<https://powcoder.com>

Add WeChat powcoder

Seasonally adjusted electrical equipment

```
ggtsdisplay(diff(eeadj))
```



Assignment Project Exam Help

- 4 PACF is suggestive of AR(3). So initial candidate model is ARIMA(3,1,0). No other obvious candidates.
- 5 Fit ARIMA(3,1,0) model along with variations: ARIMA(4,1,0), ARIMA(2,1,0), ARIMA(3,1,1), etc. ARIMA(3,1,1) has smallest AIC value.

<https://powcoder.com>
Add WeChat powcoder

Assignment Project Exam Help

```
(fit <- Arima(eadj, order=c(3,1,0)))
```

```
## Series: eadj
```

```
## ARIMA(3,1,1)
```

```
## https://powcoder.com
```

```
## Coefficients:
```

```
##          ar1      ar2      ar3      ma1
```

```
##      0.0044  0.0916  0.3698  -0.3921
```

```
## s.e. 0.1201  0.0984  0.0569  0.2426
```

```
## Add WeChat powcoder
```

```
## sigma^2 estimated as 9.577:  log likelihood=-492.69
```


```
## AIC=995.38   AICc=995.7   BIC=1011.72
```

Seasonally adjusted electrical equipment

- 6 ACF plot of residuals from ARIMA(3,1,1) model look like white noise.

Assignment Project Exam Help

check residuals (fit)



Assignment Project Exam Help

##

Ljung-Box test

<https://powcoder.com>

data: Residuals from ARIMA(3,1,1)

$Q^* = 24.034$, $df = 20$, $p\text{-value} = 0.2409$


Add WeChat powcoder

Model df: 4. Total lags used: 24

Seasonally adjusted electrical equipment

```
fit %>% forecast %>% autoplot
```

Forecasts from ARIMA(3,1,1)



Modelling procedure with `auto.arima`

1 Plot the data. Identify any unusual observations.

2 If necessary, transform the data (using a Box-Cox transformation) to stabilize the variance.

3 Use `auto.arima` to select a model.

6 Check the residuals from your chosen model by plotting the ACF of the residuals, and doing a portmanteau test of the residuals. If they do not look like white noise, try a modified model.

7 Once the residuals look like white noise, calculate forecasts.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

How does auto.arima() work?

A non-seasonal ARIMA process

$$\phi(B)(1 - B)^d y_t = c + \theta(B)\varepsilon_t$$

Need to select appropriate orders: p, q, d

Hyndman and Khandakar (JSS, 2008) algorithm:

- Select no. differences d and D via KPSS test and seasonal strength measure.
- Select p, q by minimising AICc.
- Use stepwise search to traverse model space.

How does auto.arima() work?

$$AICc = -2 \log(L) + 2(p + q + k + 1) \left[1 + \frac{(p+q+k+2)}{T-p-q-k-2} \right].$$

where L is the maximised likelihood fitted to the *differenced* data,
 $k = 1$ if $c \neq 0$ and $k = 0$ otherwise.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

How does auto.arima() work?

$$AICc = -2 \log(L) + 2(p + q + k + 1) \left[1 + \frac{(p+q+k+2)}{T-p-q-k-2} \right]$$

where L is the maximised likelihood fitted to the *differenced* data,
 $k = 1$ if $c \neq 0$ and $k = 0$ otherwise.

Step1: Select current model (with smallest AICc) from:

ARIMA(2, d, 2)

ARIMA(0, d, 0)

ARIMA(1, d, 0)

ARIMA(0, d, 1)

Add WeChat powcoder

How does auto.arima() work?

$$AICc = -2 \log(L) + 2(p + q + k + 1) \left[1 + \frac{(p+q+k+2)}{T-p-q-k-2} \right]$$

where L is the maximised likelihood fitted to the *differenced* data,
 $k = 1$ if $c \neq 0$ and $k = 0$ otherwise.

Step1: Select current model (with smallest AICc) from:

ARIMA(2, d, 2)

ARIMA(0, d, 0)

ARIMA(1, d, 0)

ARIMA(0, d, 1)

Step 2: Consider variations of current model:

- vary one of p, q , from current model by ± 1 ;
- p, q both vary from current model by ± 1 ;
- Include/exclude c from current model.

Model with lowest AICc becomes current model.


Repeat Step 2 until no lower AICc can be found.

Choosing your own model

```
ggtdisplay(internet)
```

Assignment Project Exam Help




<https://powcoder.com>



Choosing your own model

```
ggtdisplay(diff(internet))
```

diff(internet)



Choosing your own model

```
(fit <- Arima(internet,order=c(3,1,0)))
```

```
## Series: internet
```

```
## ARIMA(3,1,0)
```

```
##
```

```
## Coefficients:
```

```
##          ar1          ar2          ar3
```

```
##          1.1513    -0.6612    0.3407
```

```
## s.e         (0.0956)    (0.1353)    (0.0341)
```

```
##
```

```
## sigma^2 estimated as 9.656:  log likelihood=-  
252
```

```
## AIC=511.99    AICc=512.42    BIC=522.37
```

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Choosing your own model

```
auto.arima(internet)
```

```
## Series: internet
```

```
## ARIMA(1,1,1)
```

```
##
```

```
## Coefficients:
```

```
##          ar1          ma1
```

```
##          0.6504    0.5256
```

```
## s.e      (0.0842) (0.0896)
```

```
##
```

```
## sigma^2 estimated as 9.995:  log likelihood=-  
254.15
```

```
## AIC=514.3    AICc=514.55    BIC=522.08
```

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Choosing your own model

```
auto.arima(internet, stepwise=FALSE,  
approximation=FALSE)
```

```
## Series: internet
```

```
## ARIMA(3,1,0)
```

```
## https://powcoder.com
```

```
## Coefficients:
```

```
##          ar1          ar2          ar3
```

```
##          1.1513      -0.6612      0.3407
```

```
## s.e.    0.0950      0.1353      0.0941
```

```
##
```




```
## sigma^2 estimated as 9.656:  log likelihood=-  
252
```

```
## AIC=511.99    AICc=512.42    BIC=522.37
```

Choosing your own model

```
checkresiduals(fit)
```

Residuals from ARIMA(3,1,0)



Assignment Project Exam Help

##

Ljung-Box test

<https://powcoder.com>

data: Residuals from ARIMA(3,1,0)

$Q^* = 4.4913$, $df = 7$, $p\text{-value} = 0.7218$


Add WeChat powcoder

Model df: 3. Total lags used: 10


Choosing your own model

```
fit %>% forecast %>% autoplot
```

Forecasts from $APIM(3, 0)$



Modelling procedure



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

1 Introduction

2 Stationarity and differencing

3 Non-seasonal ARIMA models

4 Model identification

5 Estimation and order selection

6 ARIMA modeling in R

7 Forecasting

8 Seasonal ARIMA models

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

- 1 Rearrange ARIMA equation so y_t is on LHS.
- 2 Rewrite equation by replacing t by $T+h$.
- 3 On RHS, replace future observations by their forecasts, future errors by zero, and past errors by corresponding residuals.

Start with $h=1$. Repeat for $h=2,3,\dots$

ARIMA(3,1,1) forecasts: Step 1

Assignment Project Exam Help

$$(1 - \phi_1 B - \phi_2 B^2 - \phi_3 B^3)(1 - B)y_t = (1 + \theta_1 B)\varepsilon_t,$$

<https://powcoder.com>

Add WeChat powcoder

ARIMA(3,1,1) forecasts: Step 1

Assignment Project Exam Help

$$(1 - \phi_1 B - \phi_2 B^2 - \phi_3 B^3)(1 - B)y_t = (1 + \theta_1 B)\varepsilon_t,$$
$$\left[1 - (1 + \phi_1)B + (\phi_1 - \phi_2)B^2 + (\phi_2 - \phi_3)B^3 + \phi_3 B^4\right] y_t = (1 + \theta_1 B)\varepsilon_t,$$

Add WeChat powcoder

ARIMA(3,1,1) forecasts: Step 1

Assignment Project Exam Help

$$(1 - \phi_1 B - \phi_2 B^2 - \phi_3 B^3)(1 - B)y_t = (1 + \theta_1 B)\varepsilon_t,$$

$$\left[1 - (1 + \phi_1)B + (\phi_1 - \phi_2)B^2 + (\phi_2 - \phi_3)B^3 + \phi_3 B^4\right] y_t = (1 + \theta_1 B)\varepsilon_t,$$

$$y_t - (1 + \phi_1)y_{t-1} + (\phi_1 - \phi_2)y_{t-2} + (\phi_2 - \phi_3)y_{t-3} + \phi_3 y_{t-4} = \varepsilon_t + \theta_1 \varepsilon_{t-1}.$$

Add WeChat powcoder

ARIMA(3,1,1) forecasts: Step 1

Assignment Project Exam Help

$$(1 - \phi_1 B - \phi_2 B^2 - \phi_3 B^3)(1 - B)y_t = (1 + \theta_1 B)\varepsilon_t,$$

$$\left[1 - (1 + \phi_1)B + (\phi_1 - \phi_2)B^2 + (\phi_2 - \phi_3)B^3 + \phi_3 B^4\right] y_t = (1 + \theta_1 B)\varepsilon_t,$$

$$y_t - (1 + \phi_1)y_{t-1} + (\phi_1 - \phi_2)y_{t-2} + (\phi_2 - \phi_3)y_{t-3} + \phi_3 y_{t-4} = \varepsilon_t + \theta_1 \varepsilon_{t-1}.$$

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} - \phi_3 y_{t-4} + \varepsilon_t + \theta_1 \varepsilon_{t-1}.$$

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} - \phi_3y_{t-4} + \varepsilon_t + \phi_1\varepsilon_{t-1}$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Point forecasts (h=1)

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} - \phi_3y_{t-4} + \varepsilon_{t+1} + \theta_1\varepsilon_{t-1}$$

Assignment Project Exam Help

ARIMA(3,1,1) forecasts: Step 2

$$y_{T+1} = (1 + \phi_1)y_T - (\phi_1 - \phi_2)y_{T-1} - (\phi_2 - \phi_3)y_{T-2} - \phi_3y_{T-3} + \varepsilon_{T+1} + \theta_1\varepsilon_T.$$

Add WeChat powcoder

Point forecasts (h=1)

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} - \phi_3y_{t-4} + \varepsilon_t + \theta_1\varepsilon_{t-1}$$

Assignment Project Exam Help

ARIMA(3,1,1) forecasts: Step 2

$$y_{T+1} = (1 + \phi_1)y_T - (\phi_1 - \phi_2)y_{T-1} - (\phi_2 - \phi_3)y_{T-2} - \phi_3y_{T-3} + \varepsilon_{T+1} + \theta_1\varepsilon_T.$$

ARIMA(3,1,1) forecasts: Step 3

$$\hat{y}_{T+1|T} = (1 + \phi_1)y_T - (\phi_1 - \phi_2)y_{T-1} - (\phi_2 - \phi_3)y_{T-2} - \phi_3y_{T-3} + \theta_1e_T.$$

Add WeChat powcoder

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} - \phi_3y_{t-4} + \varepsilon_t + \phi_1\varepsilon_{t-1}$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Point forecasts (h=2)

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} - \phi_3y_{t-4} + \varepsilon_t + \theta_1\varepsilon_{t-1}$$

Assignment Project Exam Help

ARIMA(3,1,1) forecasts: Step 2

$$y_{T+2} = (1 + \phi_1)y_{T+1} - (\phi_1 - \phi_2)y_T - (\phi_2 - \phi_3)y_{T-1} - \phi_3y_{T-2} + \varepsilon_{T+2} + \theta_1\varepsilon_{T+1}$$

Add WeChat powcoder

Point forecasts (h=2)

$$y_t = (1 + \phi_1)y_{t-1} - (\phi_1 - \phi_2)y_{t-2} - (\phi_2 - \phi_3)y_{t-3} - \phi_3y_{t-4} + \varepsilon_t + \theta_1\varepsilon_{t-1}$$

Assignment Project Exam Help

ARIMA(3,1,1) forecasts: Step 2

$$y_{T+2} = (1 + \phi_1)y_{T+1} - (\phi_1 - \phi_2)y_T - (\phi_2 - \phi_3)y_{T-1} - \phi_3y_{T-2} + \varepsilon_{T+2} + \theta_1\varepsilon_{T+1}.$$

ARIMA(3,1,1) forecasts: Step 3

$$\hat{y}_{T+2|T} = (1 + \phi_1)\hat{y}_{T+1|T} - (\phi_1 - \phi_2)y_T - (\phi_2 - \phi_3)y_{T-1} - \phi_3y_{T-2}.$$

<https://powcoder.com>
Add WeChat powcoder

95% prediction interval

Assignment Project Exam Help

$$\hat{y}_{T+h|T} \pm 1.96 \sqrt{v_{T+h|T}}$$

where $v_{T+h|T}$ is estimated forecast variance.

<https://powcoder.com>

Add WeChat powcoder

95% prediction interval

Assignment Project Exam Help

$\hat{y}_{T+h|T} \pm 1.96 \sqrt{v_{T+h|T}}$
where $v_{T+h|T}$ is estimated forecast variance.

- $v_{t+1|t} = \sigma^2$ for all ARIMA models regardless of parameters and orders.
- Multi-step prediction intervals for ARIMA(0,0,q):

Add WeChat powcoder

$$y_t = a_t + \sum_{i=1}^q \theta_i a_{t-i}$$

$$v_{T+h|T} = \hat{\sigma}^2 \left[1 + \sum_{i=1}^{h-1} \theta_i^2 \right], \quad \text{for } h = 2, 3, \dots$$

95% Prediction interval

Assignment Project Exam Help

$$\hat{y}_{T+h|T} \pm 1.96 \sqrt{v_{T+h|T}}$$

where $v_{T+h|T}$ is estimated forecast variance.

- Multi-step prediction intervals for ARIMA(0,0,q):

$$y_t = \varepsilon_t + \sum_{i=1}^q \theta_i \varepsilon_{t-i}.$$

Add WeChat powcoder

$$v_{T+h|T} = \sigma^2 \left[1 + \sum_{i=1}^{h-1} \theta_i^2 \right] \text{ for } h = 2, 3, \dots$$

Prediction intervals

95% Prediction interval

Assignment $\hat{y}_{T+h|T} \pm 1.96\sqrt{v_{T+h|T}}$ Exam Help
where $v_{T+h|T}$ is estimated forecast variance.

- Multi-step prediction intervals for ARIMA(0,0,q):

$$y_t = \varepsilon_t + \sum_{i=1}^q \theta_i \varepsilon_{t-i}$$

$$v_{T+h|T} = \sigma^2 \left[1 + \sum_{i=1}^{h-1} \theta_i^2 \right] \text{ for } h = 2, 3, \dots$$

- AR(1): Rewrite as MA(∞) and use above result.
- Other models beyond scope of this subject.

- Prediction intervals **increase in size with forecast horizon.**
- Prediction intervals can be difficult to calculate by hand
- Calculations assume residuals are **uncorrelated** and **normally distributed.**
- Prediction intervals tend to be too narrow.
 - the uncertainty in the parameter estimates has not been accounted for.
 - the ARIMA model assumes historical patterns will not change during the forecast period.
 - the ARIMA model assumes uncorrelated future errors

Assignment Project Exam Help

For the us.gdp data

- if necessary, find a suitable Box-Cox transformation for the data;
- fit a suitable ARIMA model to the transformed data using `auto.arima()`;
- check the residual diagnostics;
- produce forecasts of your fitted model. Do the forecasts look reasonable?

<https://powcoder.com>

Add WeChat powcoder

1 Introduction

2 Stationarity and differencing

3 Non-seasonal ARIMA models

4 Model identification

5 Estimation and order selection

6 ARIMA modeling in R

7 Forecasting

8 Seasonal ARIMA models

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

ARIMA

(p, d, q)

$(P, D, Q)_m$

<https://powcoder.com>

Non-seasonal part

Seasonal part of

of the model

of the model

Add WeChat powcoder

where m = number of observations per year.

E.g., ARIMA(1, 1, 1)(1, 1, 1)₄ model (without constant)

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Seasonal ARIMA models

E.g., ARIMA(1, 1, 1)(1, 1, 1)₄ model (without constant)

$$(1 - \phi_1 B)(1 - \Phi_1 B^4)(1 - B)(1 - B^4)y_t = (1 + \theta_1 B)(1 + \Theta_1 B^4)\varepsilon_t.$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Seasonal ARIMA models

E.g., ARIMA(1, 1, 1)(1, 1, 1)₄ model (without constant)

$$(1 - \phi_1 B)(1 - \Phi_1 B^4)(1 - B)(1 - B^4)y_t = (1 + \theta_1 B)(1 + \Theta_1 B^4)\varepsilon_t.$$

(Non-seasonal)
AR(1)

(Non-seasonal)
difference

(Non-seasonal)
MA(1)

(Seasonal)
AR(1)

(Seasonal)
difference

(Seasonal)
MA(1)

Add WeChat powcoder

Seasonal ARIMA models

E.g., ARIMA(1, 1, 1)(1, 1, 1)₄ model (without constant)

$$(1 - \phi_1 B)(1 - \Phi_1 B^4)(1 - B)(1 - B^4)y_t = (1 + \theta_1 B)(1 + \Theta_1 B^4)\varepsilon_t.$$

All the factors can be multiplied out and the general model written as follows:

$$y_t = (1 + \phi_1)y_{t-1} - \phi_1 y_{t-2} + (1 + \Phi_1)y_{t-4}$$

$$- (1 + \phi_1 + \Phi_1 + \phi_1 \Phi_1)y_{t-5} + (\phi_1 + \phi_1 \Phi_1)y_{t-6}$$

$$- (\Phi_1)y_{t-8} + (\Phi_1 + \phi_1 \Phi_1)y_{t-9} - (\phi_1 \Phi_1)y_{t-10}$$

$$+ \varepsilon_t + \theta_1 \varepsilon_{t-1} + \Theta_1 \varepsilon_{t-4} + \theta_1 \Theta_1 \varepsilon_{t-5}.$$

Assignment Project Exam Help

The US Census Bureau uses the following models most often:

ARIMA(0,1,1)(0,1,1)_m with log transformation

ARIMA(0,1,2)(0,1,1)_m with log transformation

ARIMA(2,1,0)(0,1,1)_m with log transformation

ARIMA(0,2,2)(0,1,1)_m with log transformation

ARIMA(2,1,2)(0,1,1)_m with no transformation

The seasonal part of an AR or MA model will be seen in the seasonal lags of the PACF and ACF.

ARIMA(0,0,0)(0,0,1)₁₂ will show:


- a spike at lag 12 in the ACF but no other significant spikes.
- The PACF will show exponential decay in the seasonal lags; that is, at lags 12, 24, 36,

ARIMA(0,0,0)(1,0,0)₁₂ will show:

- exponential decay in the seasonal lags of the ACF
- a single significant spike at lag 12 in the PACF.

European quarterly retail trade

```
autoplot(euretail) +  
  xlab("Year") + ylab("Retail index")
```





European quarterly retail trade

```
euretail %>% diff(lag=4) %>% ggtsdisplay()
```

Assignment Project Exam Help

<https://powcoder.com>





Add WeChat powcoder

European quarterly retail trade

```
euretail %>% diff(lag=4) %>% diff() %>%
```

```
ggtsdisplay()
```



Assignment Project Exam Help

- $d = 1$ and $D = 1$ seems necessary.
- Significant spike at lag 1 in ACF suggests non-seasonal MA(1) component.
- Significant spike at lag 4 in ACF suggests seasonal MA(1) component.
- Initial candidate model: $ARIMA(0,1,1)(0,1,1)_4$
- We could also have started with $ARIMA(1,1,0)(1,1,0)_4$.

<https://powcoder.com>

Add WeChat powcoder

European quarterly retail trade



```
fit <- Arima(euretail, order=c(0,1,1),  
seasonal=c(0,1,1))  
checkresiduals(fit)
```

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Residuals from ARIMA(0,1,1)(0,1,1)[4]



Assignment Project Exam Help

##

Ljung-Box test

<https://powcoder.com>

data: Residuals from ARIMA(0,1,3)(0,1,1)[4]

$Q^* = 0.51128$, $df = 4$, $p\text{-value} = 0.9724$

Add WeChat powcoder

Model df: 4. Total lags used: 8

- ACF and PACE of residuals show significant spikes at lag 2 and maybe lag 3.
- AICc of $\text{ARIMA}(0,1,2)(0,1,1)_4$ model is 74.27.
- AICc of $\text{ARIMA}(0,1,3)(0,1,1)_4$ model is 68.39.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

- ACF and PACF of residuals show significant spikes at lag 2 and maybe lag 3.
- AICc of $\text{ARIMA}(0,1,2)(0,1,1)_4$ model is 74.27.
- AICc of $\text{ARIMA}(0,1,3)(0,1,1)_4$ model is 68.39.

```
fit <- Arima(eu_retail, order=c(0,1,3),  
            seasonal=c(0,1,1))  
checkresiduals(fit)
```



```
## Series: euretail
```

```
## ARIMA(0,1,3)(0,1,1)[4]
```

```
##
```

```
## Coefficients:
```

```
##          ma1          ma2          ma3          sma1
```

```
##          0.2630    0.3694    0.4200    -0.6636
```

```
## s.e.      0.1237    0.1253    0.1294    0.1545
```

```
##
```

```
## sigma^2 estimated as 0.156:  log likelihood=-28.63
```

```
## AIC=67.26    AICc=68.39    BIC=77.65
```

Assignment Project Exam Help



<https://powcoder.com>

Add WeChat powcoder

European quarterly retail trade

```
checkresiduals(fit)
```

Residuals from ARIMA(0, 2, 0)/(0, 1, 1)[4]



Assignment Project Exam Help

##

Ljung-Box test

<https://powcoder.com>

data: Residuals from ARIMA(0,1,3)(0,1,1)[4]

$Q^* = 0.51128$, $df = 4$, $p\text{-value} = 0.9724$


Add WeChat powcoder

Model df: 4. Total lags used: 8

European quarterly retail trade

```
autoplot(forecast(fit, h=12))
```

Forecasts from ARIMA(0, 3)(0, 1)[4]



```
auto.arima(euretail)
```

```
## Series: euretail
```

```
## ARIMA(1,1,2)(0,1,1)[4]
```

```
##
```

```
## Coefficients:
```

```
##          ar1          ma1          ma2          sma1
```

```
##          0.7362    -0.4663    0.2163    -0.8433
```

```
## s.e.      0.2243     0.1990     0.2101     0.1876
```

```
##
```

```
## sigma^2 estimated as 0.1587: log likelihood=-29.62
```

```
## AIC=69.24    AICc=70.38    BIC=79.63
```

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

```
auto.arima(euretail,  
stepwise=FALSE, approximation=FALSE)
```

```
## Series: euretail
```

```
## ARIMA(0,1,3)(0,1,1)[4]
```

```
## https://powcoder.com
```

```
## Coefficients:
```

```
##          ma1          ma2          ma3          sma1
```

```
##          0.2636 0.3694 0.4200 -0.6636
```

```
## s.e.    0.1237 0.1255 0.1294 0.1545
```


```
##
```

```
## sigma^2 estimated as 0.156: log likelihood=-28.63
```

```
## AIC=67.26 AICc=68.39 BIC=77.65
```


Add WeChat powcoder

Corticosteroid drug sales



Corticosteroid drug sales

Seasonally differenced H02 scripts



Assignment Project Exam Help

- Choose $D = 1$ and $d = 0$.
- Spikes in PACF at lags 12 and 24 suggest seasonal AR(2) term.
- Spikes in PACF suggests possible non-seasonal AR(3) term.
- Initial candidate model: $ARIMA(3,0,0)(2,1,0)_{12}$.

<https://powcoder.com>
Add WeChat powcoder

Model	AICc
ARIMA(3,0,1)(0,1,2) ₁₂	-485.48
ARIMA(3,0,1)(1,1,1) ₁₂	-484.25
ARIMA(3,0,1)(0,1,1) ₁₂	-483.67
ARIMA(3,0,1)(2,1,0) ₁₂	-476.31
ARIMA(3,0,0)(2,1,0) ₁₂	-475.12
ARIMA(3,0,2)(2,1,0) ₁₂	-474.88
ARIMA(3,0,1)(1,1,0) ₁₂	-463.40

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

```
(fit <- Arima(h02, order=c(3,0,1), seasonal=c(0,1,2),  
lambda=0))
```

```
## Series: h02
```

```
## ARIMA(3,0,1)(0,1,2)[12]
```

```
## Box-Cox transformation: lambda= 0
```

```
##
```

```
## Coefficients:
```

```
##          ar1          ar2          ar3          ma1          sma1          sma2
```

```
## -0.1603  0.5431  0.5578  0.3827 -0.5222  0.1768
```

```
## s.e.      0.1636  0.0878  0.0942  0.1895  0.0861  0.0872
```

```
##
```

```
## sigma^2 estimated as 0.004278: log likelihood=250.04
```

```
## AIC=-486.08   AICc=-485.48   BIC=-463.28
```

Assignment Project Exam Help




<https://powcoder.com>

Add WeChat powcoder

Corticosteroid drug sales

```
checkresiduals(fit, lag=36)
```

Residuals from ARIMA(3,1,2)(0,1,2)[12]



Assignment Project Exam Help

##

Ljung-Box test

<https://powcoder.com>

data: Residuals from ARIMA(3,0,1)(0,1,2)[12]

$Q^* = 50.712$, $df = 30$, $p\text{-value} = 0.01045$

Add WeChat powcoder

Model df: 6. Total lags used: 36

Assignment Project Exam Help

```
(fit <- auto.arima(h02, lambda=0))
```

```
## Series: h02
```

```
## ARIMA(2,1,3)(0,1,1)[12]
```

```
## Box-Cox transformation: lambda= 0
```

```
##
```

```
## Coefficients:
```

```
##          ar1          ar2          ma1          ma2          ma3          sma1
```

```
##      -1.0194    -0.9351    0.1717    0.2578    -0.4206    -0.6528
```

```
## s.e.      0.1648     0.1297     0.2179     0.1177     0.1140     0.0657
```

```
##
```

```
## sigma^2 estimated as 0.004203:  log likelihood=250.8
```

```
## AIC=-487.6   AICc=-486.99   BIC=-464.83
```




<https://powcoder.com>

Add WeChat powcoder

Corticosteroid drug sales

```
checkresiduals(fit, lag=36)
```

Residuals from ARIMA(2, ,3)(0,1,1)/121



Assignment Project Exam Help

##

Ljung-Box test

<https://powcoder.com>

data: Residuals from ARIMA(2,1,3)(0,1,1)[12]

$Q^* = 46.149$, $df = 30$, $p\text{-value} = 0.03007$

Add WeChat powcoder

Model df: 6. Total lags used: 36


```
(fit <- auto.arima(h02, lambda=0, max.order=9,  
  stepwise=FALSE, approximation=FALSE))
```

```
## Series: h02  
## ARIMA(4,1,1)(2,1,2)[12]  
## Box Cox transformation: lambda= 0  
##  
## Coefficients:  
##          ar1      ar2      ar3      ar4      ma1      sar1      sar2      sma1  
##      -0.0425  0.2098  0.2017  -0.2273  -0.7424   0.6213  -0.3832  -1.2019  
## s.e.   0.2167  0.1813  0.1144   0.0810   0.2074   0.2421   0.1185   0.2491  
##      sma1  
##      0.4959  
## s.e.   0.2135  
##  
## sigma^2 estimated as 0.004049:  log likelihood=254.31  
## AIC=-488.63   AICc=-487.4   BIC=-456.1
```

Assignment Project Exam Help




<https://powcoder.com>

Add WeChat powcoder

Corticosteroid drug sales

```
checkresiduals(fit, lag=36)
```

Residuals from ARIMA(4,1,1)(2,1,2)[12]



Assignment Project Exam Help

##

Ljung-Box test

<https://powcoder.com>

data: Residuals from ARIMA(4,1,1)(2,1,2)[12]

$Q^* = 36.456$, $df = 27$, $p\text{-value} = 0.1057$

Add WeChat powcoder

Model df: 9. Total lags used: 36

Corticosteroid drug sales

Training data: July 1991 to June 2006

Test data: July 2006–June 2008

```
getrmse <- function(x,h,...)
```

```
{
```

```
  train.end <- time(x)[length(x)-h]
```

```
  test.start <- time(x)[length(x)-h+1]
```

```
  train <- window(x,end=train.end)
```

```
  test <- window(x,start=test.start)
```

```
  fit <- Arima(train,...)
```

```
  fc <- forecast(fit,h=h)
```

```
  return(accuracy(fc,test)[2,"RMSE"])
```

```
}
```

```
getrmse(h02,h=24,order=c(3,0,0),seasonal=c(2,1,0),lambda=0)
```

```
getrmse(h02,h=24,order=c(3,0,1),seasonal=c(2,1,0),lambda=0)
```

```
getrmse(h02,h=24,order=c(3,0,2),seasonal=c(2,1,0),lambda=0)
```

```
getrmse(h02,h=24,order=c(3,0,1),seasonal=c(1,1,0),lambda=0)
```

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Corticosteroid drug sales

Model	RMSE
ARIMA(4,1,1)(2,1,2)[12]	0.0615
ARIMA(3,0,1)(0,1,2)[12]	0.0622
ARIMA(3,0,1)(1,1,1)[12]	0.0630
ARIMA(2,1,4)(0,1,1)[12]	0.0632
ARIMA(2,1,3)(0,1,1)[12]	0.0634
ARIMA(3,0,3)(0,1,1)[12]	0.0638
ARIMA(2,1,5)(0,1,1)[12]	0.0640
ARIMA(3,0,1)(0,1,1)[12]	0.0644
ARIMA(3,0,2)(0,1,1)[12]	0.0644
ARIMA(3,0,2)(2,1,0)[12]	0.0645
ARIMA(3,0,1)(2,1,0)[12]	0.0646
ARIMA(3,0,0)(2,1,0)[12]	0.0661

Assignment Project Exam Help


- Models with lowest AICc values tend to give slightly better results than the other models
- AICc comparisons must have the same orders of differencing. But RMSE test set comparisons can involve any models.
- Use the best model available, even if it does not pass all tests.

<https://powcoder.com>

Add WeChat powcoder

Corticosteroid drug sales

```
fit <- Arima(h02, order=c(3,0,1), seasonal=c(0,1,2),  
            lambda=0)  
autoplot(forecast(fit)) +  
  ylab("H02 sales (million scripts)") + xlab("Year")
```



Understanding Seasonal ARIMA models

- If $c = 0$ and $d + D = 0$, the long-term forecasts will go to zero.
- If $c = 0$ and $d + D = 1$, the long-term forecasts will go to a non-zero constant determined by the last few observations.
- If $c = 0$ and $d + D = 2$, the long-term forecasts will follow a straight line with intercept and slope determined by the last few observations.
- If $c \neq 0$ and $d + D = 0$, the long-term forecasts will go to the mean of the data.
- If $c \neq 0$ and $d + D = 1$, the long-term forecasts will follow a straight line with slope equal to the mean of the differenced data.
- If $c \neq 0$ and $d + D = 2$, the long-term forecasts will follow a quadratic trend.