

Choice under uncertainty**Part 1**

1. Introduction to choice under uncertainty	2
2. Risk aversion	15
3. Acceptable gambles	19

Part 2

4. Measures of risk aversion	24
5. Insurance	30
6. Efficient risk sharing	35
7. Portfolio choice	47

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

57 slides

1. Introduction to choice under uncertainty (two states)

Let X be a set of possible outcomes (“states of the world”).

An element of X might be a consumption vector, health status, inches of rainfall etc.

Initially, simply think of each element of X as a consumption bundle. Let \bar{x} be the most preferred element of X and let \underline{x} be the least preferred element.

**

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

1. Introduction to choice under uncertainty (two states)

Let X be a set of possible outcomes (“states of the world”).

An element of X might be a consumption vector, health status, inches of rainfall etc.

Initially, simply think of each element of X as a consumption bundle. Let \bar{x} be the most preferred element of X and let \underline{x} be the least preferred element.

Consumption prospects

Suppose that there are only two states of the world, $X = \{x_1, x_2\}$. Let π_1 be the probability that the state is x_1 so that $\pi_2 = 1 - \pi_1$ is the probability that the state is x_2 .

<https://powcoder.com>

We write this “consumption prospect” as follows:

$$(x; \pi) = (x_1, x_2; \pi_1, \pi_2)$$

Add WeChat powcoder

*

1. Introduction to choice under uncertainty (two states)

Let X be a set of possible outcomes (“states of the world”).

An element of X might be a consumption vector, health status, inches of rainfall etc.

Initially, simply think of each element of X as a consumption bundle. Let \bar{x} be the most preferred element of X and let \underline{x} be the least preferred element.

Consumption prospects

Suppose that there are only two states of the world, $X = \{x_1, x_2\}$. Let π_1 be the probability that the state is x_1 so that $\pi_2 = 1 - \pi_1$ is the probability that the state is x_2 .

<https://powcoder.com>

We write this “consumption prospect” as follows:

$$(x; \pi) = (x_1, x_2; \pi_1, \pi_2)$$

Add WeChat powcoder

If we make the usual assumptions about preferences, but now on prospects, it follows that preferences over prospects can be represented by a continuous utility function

$$U(x_1, x_2, \pi_1, \pi_2) .$$

Prospect or “Lottery”

$$L = (x_1, x_2, \dots, x_S; \pi_1, \dots, \pi_S)$$

(outcomes; probabilities)

Consider two prospects or “lotteries”, L_A and L_B

$$L_A = (x_1, x_2, \dots, x_S; \pi_1^A, \dots, \pi_S^A) \quad L_B = (c_1, c_2, \dots, c_S; \pi_1^B, \dots, \pi_S^B)$$

Independence Axiom (axiom of complex gambles).

Assignment Project Exam Help

Suppose that a consumer is indifferent between these two prospects (we write $L_A \sim L_B$).


Then for any probabilities π_1 and π_2 summing to 1 and any other lottery L_C

<https://powcoder.com>

$$(L_A, L_C; \pi_1, \pi_2) \sim (L_B, L_C; \pi_1, \pi_2)$$

Add WeChat powcoder

Tree representation



This axiom can be generalized as follows:

Suppose that a consumer is indifferent between the prospects L_A and L_B

and is also indifferent between the two prospects L_C and L_D ,

i.e. $L_A \sim L_B$ and $L_C \sim L_D$

Then for any probabilities π_1 and π_2 summing to 1,

$(L_A, L_C; \pi_1, \pi_2) \sim (L_B, L_D; \pi_1, \pi_2)$


Assignment Project Exam Help

Tree representation

We wish to show that if $L_A \sim L_B$ and $L_C \sim L_D$ then


<https://powcoder.com>

Add WeChat powcoder



Proof: $L_A \sim L_B$ and $L_C \sim L_D$

Step 1: By the Independence Axiom, since $L_A \sim L_B$



Assignment Project Exam Help


*

<https://powcoder.com>

Add WeChat powcoder

Proof: $L_A \sim L_B$ and $L_C \sim L_D$

Step 1: By the Independence Axiom, since $L_A \sim L_B$




Assignment Project Exam Help

Step 2: By the Independence Axiom, since $L_C \sim L_D$

<https://powcoder.com>

Add WeChat powcoder



Expected utility

Consider some very good outcome \bar{x} and very bad outcome \underline{x} and outcomes x_1 and x_2 satisfying

$$\underline{x} \prec x_1 \prec \bar{x} \quad \text{and} \quad \underline{x} \prec x_2 \prec \bar{x}$$

Reference lottery

$L_R(v) = (\bar{w}, \underline{w}, v, 1-v)$ so v is the probability of the very good outcome.


$$L_R(0) \prec x_1 \prec L_R(1) \quad \text{and} \quad L_R(0) \prec x_2 \prec L_R(1)$$

Then for some probabilities $v(x_1)$ and $v(x_2)$

$$x_1 \sim L_R(v(x_1)) = (\bar{x}, \underline{x}; v(x_1), 1-v(x_1)) \quad \text{and} \quad x_2 \sim L_R(v(x_2)) = (\bar{x}, \underline{x}; v(x_2), 1-v(x_2))$$


Then by the independence axiom

$$(x_1, x_2; \pi_1, \pi_2) \sim (L_R(v(x_1)), L_R(v(x_2)); \pi_1, \pi_2)$$




Definition: Expectation of $v(x)$


$$\mathbb{E}[v(x)] \equiv \pi_1 v(x_1) + \pi_2 v(x_2)$$



Note that in the big tree there are only two outcomes, \bar{x} and \underline{x} . The probability of the very good outcome is $\pi_1 v(x_1) + \pi_2 v(x_2) = \mathbb{E}[v(x)]$. The probability of the very bad outcome is $1 - \mathbb{E}[v(x)]$. Therefore



We showed that



Assignment Project Exam Help

i.e.

$$(x_1, x_2; \pi_1, \pi_2) \sim (\bar{x}, \underline{x}; \mathbb{E}[v], 1 - \mathbb{E}[v])$$

<https://powcoder.com>

Add WeChat powcoder

Thus the expected win probability in the reference lottery is a representation of preferences over prospects.

An example:

A consumer with wealth \hat{w} is offered a “fair gamble” . With probability $\frac{1}{2}$ his wealth will be $\hat{w} + x$ and with probability $\frac{1}{2}$ his wealth will be $\hat{w} - x$. If he rejects the gamble his wealth remains \hat{w} . Note that this is equivalent to a prospect with $x=0$

In prospect notation the two alternatives are

$$(w_1, w_2; \pi_1, \pi_2) = (\hat{w}, \hat{w}; \frac{1}{2}, \frac{1}{2})$$

and

$$(w_1, w_2; \pi_1, \pi_2) = (\hat{w} + x, \hat{w} - x; \frac{1}{2}, \frac{1}{2}).$$


These are depicted in the figure assuming $x > 0$.

Expected utility

$$U(w_1, w_2, \pi_1, \pi_2) = \mathbb{E}[v] = \pi_1 v(w_1) + \pi_2 v(w_2)$$

Class discussion

MRS if $v(w)$ is a concave function



Convex preferences


The two prospects are depicted opposite.

The level set for $U(w_1, w_2; \frac{1}{2}, \frac{1}{2})$ through the riskless prospect N is depicted.

Note that the superlevel set

$$U(w_1, w_2; \frac{1}{2}, \frac{1}{2}) \geq U(\hat{w}, \hat{w}, \frac{1}{2}, \frac{1}{2})$$

is a convex set.



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

*

Convex preferences


The two prospects are depicted opposite.

The level set for $U(w_1, w_2; \frac{1}{2}, \frac{1}{2})$ through the riskless prospect N is depicted.

Note that the superlevel set

$$U(w_1, w_2; \frac{1}{2}, \frac{1}{2}) \geq U(\hat{w}, \hat{w}; \frac{1}{2}, \frac{1}{2})$$

is a convex set.



<https://powcoder.com>

Add WeChat powcoder

This is the set of acceptable gambles for the consumer.

As depicted the consumer strictly prefers the riskless prospect N to the risky prospect R .

Most individuals, when offered such a gamble (say over \$5) will not take this gamble.

2. Risk aversion

Class Discussion: Which alternative would you choose?

$$N: (w_1, w_2; \pi_1, \pi_2) = (\hat{w}, \hat{w}; \pi_1, \pi_2) \quad R: (w_1, w_2; \pi_1, \pi_2) = (\hat{w} + x, \hat{w} - x; \pi_1, \pi_2) \text{ where } \pi_1 = \frac{50}{100}$$

What if the gamble were “favorable” rather than “fair”

$$R: (w_1, w_2; \pi_1, \pi_2) = (\hat{w} + x, \hat{w} - x; \pi_1, \pi_2) \text{ where (i) } \pi_1 = \frac{55}{100} \text{ (ii) } \pi_1 = \frac{60}{100} \text{ (iii) } \pi_1 = \frac{75}{100}$$

Assignment Project Exam Help

<https://powcoder.com>

*

Add WeChat powcoder

Class Discussion: Which alternative would you choose?

$$N: (w_1, w_2; \pi_1, \pi_2) = (\hat{w}, \hat{w}; \pi_1, \pi_2) \quad R: (w_1, w_2; \pi_1, \pi_2) = (\hat{w} + x, \hat{w} - x; \pi_1, \pi_2) \text{ where } \pi_1 = \frac{50}{100}$$

What if the gamble were “favorable” rather than “fair”

$$R: (w_1, w_2; \pi_1, \pi_2) = (\hat{w} + x, \hat{w} - x; \pi_1, \pi_2) \text{ where (i) } \pi_1 = \frac{55}{100} \text{ (ii) } \pi_1 = \frac{60}{100} \text{ (iii) } \pi_1 = \frac{75}{100}$$

Assignment Project Exam Help

What is the smallest integer n such that you would gamble if $\pi_1 = \frac{n}{100}$? <https://powcoder.com>

Preference elicitation

Add WeChat powcoder

In an attempt to elicit your preferences write down your number n (and your first name) on a piece of paper. The two participants with the lowest number n will be given the riskless opportunity.

Let the three lowest integers be n_1, n_2, n_3 . The win probability will not be $\frac{n_1}{100}$ or $\frac{n_2}{100}$. Both will get

the higher win probability $\frac{n_3}{100}$.

2. Risk preferences

$$U(x, \pi) = \pi_1 v(x_1) + \pi_2 v(x_2) \quad \text{or} \quad U(x, \pi) = \mathbb{E}[v]$$


Risk preferring consumer

Consider the two wealth levels x_1 and $x_2 > x_1$.

$$v(\pi_1 x_1 + \pi_2 x_2) < \pi_1 v(x_1) + \pi_2 v(x_2)$$

If $v(x)$ is convex, then the slope of $v(x)$

is strictly increasing as shown in the top figure.



Add WeChat powcoder

Consumer prefers risk

$$U(x, \pi) = \pi_1 v(x_1) + \pi_2 v(x_2)$$

Risk averse consumer

$$v(\pi_1 x_1 + \pi_2 x_2) > \pi_1 v(x_1) + \pi_2 v(x_2) .$$

In the lower figure $u(x)$ is strictly concave so that


$$v(\pi_1 x_1 + \pi_2 x_2) > \pi_1 v(x_1) + \pi_2 v(x_2) = \mathbb{E}[v].$$

In practice consumers exhibit aversion to such a risk.

Thus we will (almost) always assume that the


expected utility function $v(x)$ is a strictly increasing

strictly concave function.



Consumer prefers risk

Add WeChat powcoder



Risk averse consumer

Class Discussion:

If consumers are risk averse why do they go to Las Vegas?

3. Acceptable gambles: Improving the odds to make the gamble just acceptable.

New risky alternative: $(w_1, w_2; \pi_1, \pi_2) = (\hat{w} + x, \hat{w} - x; \frac{1}{2} + \alpha, \frac{1}{2} - \alpha)$.

Choose α so that the consumer is indifferent between gambling and not gambling.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

3. Acceptable gamble: Improving the odds to make the gamble just acceptable.

New risky alternative: $(w_1, w_2; \pi_1, \pi_2) = (\hat{w} + x, \hat{w} - x; \frac{1}{2} + \alpha, \frac{1}{2} - \alpha)$.

Choose α so that the consumer is indifferent between gambling and not gambling.

For small x we can use the quadratic approximation of the utility function

Quadratic approximation of his utility

As long as x is small we can approximate his utility

as a quadratic. Suppose $u(w+x) = \ln(w+x)$.


Define $\bar{u}(x) = \ln(w+x)$.

Then (i) $\bar{u}(0) = \ln w$ (ii) $\bar{u}'(0) = \frac{1}{w}$ and (iii) $\bar{u}''(0) = -\frac{1}{w^2}$

Consider the quadratic function

$$q(x) = \ln w + \left(\frac{1}{w}\right)x - \frac{1}{2}\left(\frac{1}{w^2}\right)x^2. \quad (3.1)$$

If you check you will find that $\bar{u}(x)$ and $q(x)$ have the same, value, first derivative and second derivative at $x=0$. We then use this quadratic approximation to compute the gambler's (approximated) expected gain.



With probability $\frac{1}{2} + \alpha$ his payoff is $q(x)$ and with probability $\frac{1}{2} - \alpha$ his payoff is $q(-x)$. Therefore his expected payoff is

$$\mathbb{E}[q(x)] = (\tfrac{1}{2} + \alpha)q(x) + (\tfrac{1}{2} - \alpha)q(-x)$$

Substituting from (3.1)

$$\mathbb{E}[q(x)] = (\tfrac{1}{2} + \alpha) \left[\ln w + \left(\frac{1}{w}\right)x - \frac{1}{2} \left(\frac{1}{w^2}\right)x^2 \right]$$

$$+ (\tfrac{1}{2} - \alpha) \left[\ln w + \left(\frac{1}{w}\right)(-x) - \frac{1}{2} \left(\frac{1}{w^2}\right)(-x)^2 \right]$$

Assignment Project Exam Help

*

<https://powcoder.com>

Add WeChat powcoder

With probability $\frac{1}{2} + \alpha$ his payoff is $q(x)$ and with probability $\frac{1}{2} - \alpha$ his payoff is $q(-x)$. Therefore his expected payoff is

$$\mathbb{E}[q(x)] = (\tfrac{1}{2} + \alpha)q(x) + (\tfrac{1}{2} - \alpha)q(-x)$$

Substituting from (3.1)

$$\begin{aligned} \mathbb{E}[q(x)] &= (\tfrac{1}{2} + \alpha) \left[\ln w + \left(\tfrac{1}{w}\right)x - \tfrac{1}{2} \left(\tfrac{1}{w^2}\right)x^2 \right] \\ &\quad + (\tfrac{1}{2} - \alpha) \left[\ln w + \left(\tfrac{1}{w}\right)(-x) - \tfrac{1}{2} \left(\tfrac{1}{w^2}\right)(-x)^2 \right]. \end{aligned}$$

Collecting terms,

$$\mathbb{E}[q(x)] = \ln w + 2\alpha \left(\tfrac{1}{w}\right)x - \tfrac{1}{2} \left(\tfrac{1}{w^2}\right)x^2.$$

If the gambler rejects the opportunity his utility is $\ln w$. Thus his expected gain is

$$\mathbb{E}[q(x)] - \ln w = 2\alpha \left(\tfrac{1}{w}\right)x - \tfrac{1}{2} \left(\tfrac{1}{w^2}\right)x^2 = \frac{2x}{w} \left[\alpha - \tfrac{1}{4} \left(\tfrac{1}{w}\right)x \right].$$

Thus the gambler should take the small gamble if and only if $\alpha > \frac{1}{4} \left(\tfrac{1}{w}\right)x$.

The general case: quadratic approximation of his utility

$$q(x) = v(\hat{w}) + v'(\hat{w})x + \frac{1}{2}v''(\hat{w})x^2$$

Class Exercise: Confirm that the value and the first two derivatives of $\bar{u}(x) \equiv v(\hat{w} + x)$ and $q(x)$ are equal at $x=0$.

The expected value utility of the risky alternative is

$$\mathbb{E}[u(\hat{w} + x)] \approx \mathbb{E}[q(x)] = (\frac{1}{2} + \alpha)q(x) + (\frac{1}{2} - \alpha)q(-x)$$

**

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

The general case: quadratic approximation of his utility

$$q(x) = v(\hat{w}) + v'(\hat{w})x + \frac{1}{2}v''(\hat{w})x^2$$

Class Exercise: Confirm that the value and the first two derivatives of $v(\hat{w} + x)$ and $q(x)$ are equal at $x = 0$.

The expected value utility of the risky alternative is

$$\mathbb{E}[u(\hat{w} + x)] \approx \mathbb{E}[q(x)] = \left(\frac{1}{2} + \alpha\right)q(x) + \left(\frac{1}{2} - \alpha\right)q(-x)$$

$$= \left(\frac{1}{2} + \alpha\right)\left[v(\hat{w}) + v'(\hat{w})x + \frac{1}{2}v''(\hat{w})x^2\right]$$

$$+ \left(\frac{1}{2} - \alpha\right)\left[v(\hat{w}) + v'(\hat{w})(-x) + \frac{1}{2}v''(\hat{w})(-x)^2\right]$$

Collecting terms,

$$\mathbb{E}[q(x)] = v(\hat{w}) + 2\alpha v'(\hat{w})x - \frac{1}{2}v''(\hat{w})x^2.$$

*

The general case: quadratic approximation of his utility

$$q(x) = v(\hat{w}) + v'(\hat{w})x + \frac{1}{2}v''(\hat{w})x^2$$

Class Exercise: Confirm that the value and the first two derivatives of $v(\hat{w} + x)$ and $q(x)$ are equal at $x = 0$.

The expected value utility of the risky alternative is

$$\mathbb{E}[u(\hat{w} + x)] \approx \mathbb{E}[q(x)] = (\frac{1}{2} + \alpha)q(x) + (\frac{1}{2} - \alpha)q(-x)$$

$$= (\frac{1}{2} + \alpha)[v(\hat{w}) + v'(\hat{w})x + \frac{1}{2}v''(\hat{w})x^2]$$

$$+ (\frac{1}{2} - \alpha)[v(\hat{w}) + v'(\hat{w})(-x) + \frac{1}{2}v''(\hat{w})(-x)^2]$$

Collecting terms,

$$\mathbb{E}[q(x)] = v(\hat{w}) + 2\alpha v'(\hat{w})x + \frac{1}{2}v''(\hat{w})x^2.$$

The gain in expected utility is therefore

$$\begin{aligned}\mathbb{E}[q(x)] - v(\hat{w}) &= 2\alpha v'(\hat{w})x + \frac{1}{2}v''(\hat{w})x^2 \\ &= 2v'(\hat{w})x[\alpha - \frac{1}{4}(-\frac{v''(\hat{w})}{v'(\hat{w})})x]\end{aligned}$$

Thus the probability of the good outcome must be increased by $\alpha = \frac{1}{4}(-\frac{v''(\hat{w})}{v'(\hat{w})})x$.