

FIT2014 Theory of Computation

# Assignment Project Exam Help

Lecture 11

(A) Closure properties;

<https://powcoder.com>  
(B) Pumping Lemma for Regular Languages

Add WeChat [powcoder](https://powcoder.com)  
slides by Graham Farr  
based in part on previous slides by David Albrecht

COMMONWEALTH OF AUSTRALIA

Copyright Regulations 1969

Warning

This material has been reproduced and communicated to you by or on behalf of Monash University  
in accordance with s113P of the Copyright Act 1968 (the Act).

The material in this communication may be subject to copyright under the Act.

Any further reproduction or communication of this material by you may be the subject of copyright protection under the Act.

Do not remove this notice.

# Assignment Project Exam Help

- ▶ Closure properties of regular languages
- ▶ Circuits in FAs
- ▶ Pumping Lemma
- ▶ Non-regular Languages

<https://powcoder.com>

Add WeChat powcoder

## Closure properties of regular languages

### Definition

If doing some operation on regular languages always produces another regular language, then we say that the class of regular languages is **closed** under that operation.

Assignment Project Exam Help

We will see that regular languages are closed under:

- ▶ complement
- ▶ union
- ▶ intersection
- ▶ concatenation

<https://powcoder.com>

Add WeChat powcoder

### Theorem.

The complement of a regular language is regular.

We prove this using Kleene's Theorem.

## Closure properties of regular languages

### Theorem

The complement of a regular language is regular.

### Proof. (outline)

Suppose we have a Regular Language.

There must be a regular expression that defines it.

So, by Kleene's Theorem, there is a Finite Automaton (FA) that defines this language.

We can convert this FA into one that defines the complement of the language. (See Lecture 7.)

So, by Kleene's Theorem, there is a regular expression that defines the complement.  $\square$

## Closure properties of regular languages

### Theorem.

The union of two regular languages is regular.

### Proof.

Suppose  $L_1$  and  $L_2$  are regular.

By definition of 'regular language',

there exist regular expressions  $R_1$  and  $R_2$

that describe  $L_1$  and  $L_2$ , respectively.

Then  $R_1 \cup R_2$  is a regular expression that describes  $L_1 \cup L_2$ .

- This uses part 3(iii) of the inductive definition of regular expressions in Lecture 6.

So  $L_1 \cup L_2$  is regular.  $\square$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

# Assignment Project Exam Help

## **Theorem.**

The intersection of two regular languages is regular.

<https://powcoder.com>

We can't just mimic the proof that regular languages are closed under union, since there is no  $\cap$  operation on regular expressions.

Add WeChat powcoder

## Closure properties of regular languages

### Theorem.

The intersection of two regular languages is regular.

### Proof.

Suppose  $L_1$  and  $L_2$  are regular.

We know that their complements  $\overline{L_1}$  and  $\overline{L_2}$  are regular.

So the union of these,  $\overline{L_1} \cup \overline{L_2}$ , is therefore regular, by the previous Theorem.

Its complement  $\overline{\overline{L_1} \cup \overline{L_2}}$  must also be regular.

$$\text{But } \overline{\overline{L_1} \cup \overline{L_2}} = \overline{\overline{L_1}} \cap \overline{\overline{L_2}} = L_1 \cap L_2.$$

So  $L_1 \cap L_2$ , must also be regular.  $\square$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

## Closure properties of regular languages

### Exercises

- ▶ Prove that the class of regular languages is closed under **concatenation**.

**Assignment Project Exam Help**  
 $L_1 L_2 := \{xy : x \in L_1, y \in L_2\}$

- ▶ Prove that the class of regular languages is closed under **symmetric difference**.  
(You can use the closure results we've already proved.)

$L_1 \triangle L_2 := \{\text{strings in } L_1 \text{ but not in } L_2, \text{ or in } L_2 \text{ but not in } L_1\}$

**Add WeChat powcoder**

- ▶ Is the class of regular languages closed under taking **subsets**?  
i.e., is a subset of a regular language necessarily regular?
- ▶ Is the class of regular languages closed under taking **supersets**?  
i.e., is a superset of a regular language necessarily regular?



## Circuits in Finite Automata

### Definition

A **circuit** is a directed path which starts and ends at the same state.

The **length** of a circuit is the number of edges in the path.

### Observation

Take any Finite Automaton.

Take any string  $w$  with has at least as many letters as there are states in that Finite Automaton.

Then the path taken for input  $w$  must contain a circuit.

We can divide  $w$  up naturally into three parts,  $w = xyz$ , where:

$x$  := the part *before* the circuit;

$y$  := the part that *goes around* the circuit;

$z$  := the part *after* the circuit;

$w = \underline{a} \underline{baaa} \underline{abb}$


If  $w$  is accepted,  
then so are ...

$\underline{a} \underline{abb}$   
 $\underline{x} \quad \underline{z}$

$\underline{a} \underline{baaa} \underline{abb}$   
 $\underline{x} \quad \underline{y} \quad \underline{z}$

$\underline{a} \underline{baaa} \underline{baaa} \underline{abb}$   
 $\underline{x} \quad \underline{y} \quad \underline{y} \quad \underline{z}$

...  
...



$w = \underline{ba} \underline{bb}$   
           $y$    $z$

$x = \epsilon$

If  $w$  is accepted,  
then so are ...

$\underline{bb}$   
   $z$

$\underline{ba} \underline{bb}$   
   $y$    $z$


$\underline{ba} \underline{ba} \underline{bb}$   
   $y$    $y$    $z$

...

...

<https://powcoder.com>

Add WeChat powcoder



## Pumping Lemma

### Theorem. (Pumping Lemma)

Let  $L$  be an infinite regular language, accepted by a FA with  $N$  states.

Then for all words  $w \in L$  with at least  $N$  letters  
there exist strings  $x, y, z$ , with  $y \neq \varepsilon$ , such that

►  $w = xyz$

►  $\text{length}(x) + \text{length}(y) \leq N$

► for all  $i \geq 0$ ,  $xy^iz \in L$ ,

i.e.,

$$xz, xyz, xyyz, \dots, xy^nz, \dots \in L.$$

Add WeChat powcoder

Symbolically:

$$\forall w \in L : |w| \geq N \Rightarrow (\exists x, y, z : (w = xyz) \wedge (y \neq \varepsilon) \wedge (|x| + |y| \leq N) \wedge (\forall i \geq 0 : xy^iz \in L))$$

## Pumping Lemma

### Proof.

Take any word  $w \in L$  with  $\geq N$  letters.

By our earlier Observation on circuits in FAs, the path taken by  $w$  must include a circuit.

Let

- $x$  be the letters of  $w$  up to the first circuit.
- $y$  be the letters corresponding to the circuit.
- $z$  be the remaining letters of  $w$ .

We have:

- ▶  $w = xyz$  by construction.
- ▶ Since the circuit exists,  $y \neq \epsilon$ .
- ▶  $\text{length}(x) + \text{length}(y) \leq N$ , since the FA reads  $xy$  without repeating any state.
- ▶ Since  $w = xyz \in L$ , and  $y$  starts and finishes at  $\text{endState}(x)$ , and  $z$  goes from  $\text{endState}(x)$  to a Final State, we can repeat  $y$  any number of times (or none) and still we end up at the same Final State. □

## Pumping Lemma: application

### Consequence

Using the Pumping Lemma we can show there are non-regular languages

### Method

Assume  $L$  is regular.

Then, by Kleene's Theorem, it is recognised by some FA.

Let  $N$  be the number of states in this FA.

Choose a suitable word  $w \in L$ , of length  $\geq N$ .

Show that, for any  $x, y \neq \varepsilon$ , and  $z$  such that  $w = xyz$  and  $|xy| \leq N \dots$

$\dots$  there exists  $i \geq 0$  s.t.  $xy^iz \in L$

*Contradiction.*

Compare quantifiers above with those in Pumping Lemma

## Non-regular languages

HALF-AND-HALF:

$$L := \{a^n b^n : n \geq 0\} = \{\varepsilon, ab, aabb, aaabbb, \dots\}.$$

**Theorem.**

$L$  is not regular.

**Proof.** (by contradiction)

Assume that  $L$  is regular.

Let  $N = \#$  states in an FA for it.

Choose  $w := a^{\lceil N/2 \rceil} b^{\lceil N/2 \rceil}$ .



Observe that  $|w| \geq N$ .

Consider any  $x, y \neq \varepsilon$ , and  $z$  such that  $w = xyz$  and  $|xy| \leq N$ .

*Think:* are  $xz, xyz, xyyz, \dots, xy^N z, \dots$  all in  $L$ ?

## Non-regular languages

**Case 1:**  $y$  is all a's.

$\overbrace{aaa \dots aa}^{\lceil N/2 \rceil \text{ letters}} \underbrace{\hspace{1cm}}_y \overbrace{bbb \dots bb}^{\lceil N/2 \rceil \text{ letters}}$

Then  $xyyz$  has more a's than b's, since  $y \neq \epsilon$ .

So  $xy^2z \notin L$ .

**Case 2:**  $y$  is all b's.

$aaa \dots aa \underbrace{bbb \dots bb}_y$

Then  $xyyz$  has more b's than a's, since  $y \neq \epsilon$ .

So  $xy^2z \notin L$ .

**Case 3:**  $y$  contains  $ab$ .

$aaa \dots aa \underbrace{abbb \dots bb}_y$

Then  $xyyz$  has two occurrences of  $ab$ . This cannot happen for strings in  $L$ . So  $xy^2z \notin L$ .

In every possible case, we have found an  $i$  such that  $xy^iz \notin L$ .

This violates the conclusion of the Pumping Lemma.

*Contradiction.*  $\square$



## Non-regular languages

HALF-AND-HALF:

$L := \{a^n b^n : n \geq 0\} = \{\epsilon, ab, aabb, aaabbb, \dots\}.$

**Theorem.**

$L$  is not regular.

**Proof.** (by contradiction)

Assume that  $L$  is regular. Let  $N = \#$  states in an FA for it.

Choose  $w = a^N b^N$ .

*[No need for  $w$  to be of minimum length.]*

Consider any  $x, y \neq \epsilon$ , and  $z$  such that  $w = xyz \dots$

$\dots$  and  $|xy| \leq N$ . *[Previous proof didn't use  $|xy| \leq N$ . Can it help?]*

Think: are  $xz, xyx, xy^2x, \dots, xy^N z, \dots$  all in  $L$ ?

How many cases now?

## Non-regular languages

Just one case:  $y$  is all a's.

$\overbrace{aaa \dots aa}^{N \text{ letters}} \overbrace{bbb \dots bb}^{N \text{ letters}}$

# Assignment Project Exam Help

Consider  $xyyz$ .

$\overbrace{aaa \dots aa}^{N+|y| \text{ letters}} \overbrace{bbb \dots bb}^{N \text{ letters}}$

# <https://powcoder.com>

It has more a's than b's, since  $y \neq \epsilon$ .

So  $xy^2z \notin L$ .

This violates the conclusion of the Pumping Lemma.

Contradiction.  $\square$

# Add WeChat powcoder

## Non-regular languages

$\text{EQUAL} := \{ \text{all words which have an equal number of a's and b's} \}$

$= \{ \epsilon, ab, ba, aabb, abba, abab, abba, baab, \dots \}$

**Theorem.**

EQUAL is not regular.

**Proof.** Assume EQUAL is regular.

Observe:

$\text{HALF-AND-HALF} := \{ a^n b^n : n \geq 0 \} = \text{EQUAL} \cap a^* b^*.$

This implies that HALF-AND-HALF is also regular, since the language defined by  $a^* b^*$  is regular, and regular languages are closed under intersection.

But we have just seen that HALF-AND-HALF is non-regular.

This is a contradiction.

So our initial assumption, that EQUAL is regular, is wrong.

Therefore EQUAL is non-regular.  $\square$

## Non-regular languages

PALINDROME  $:=$  { all the strings which are the same if they are spelt backwards }  
 $=$  {  $\epsilon$ ,  $a$ ,  $b$ ,  $aa$ ,  $bb$ ,  $aaa$ ,  $abba$ ,  $baab$ ,  $bbbb$ , ... }

**Theorem.**

PALINDROME is non-regular.

**Proof.** (by contradiction)

Assume PALINDROME is regular.

Then there exists a FA with  $N$  states which accepts PALINDROME.

Choose  $w = a^N b a^N$ .

$\overbrace{aaa \cdots \cdots aa}^{N \text{ letters}} b \overbrace{aaa \cdots \cdots aa}^{N \text{ letters}}$

## Non-regular languages

Consider all strings  $x$ ,  $y \neq \varepsilon$ , and  $z$  such that

►  $w = xyz$ ,

►  $\text{length}(x) + \text{length}(y) \leq N$ .



Consider  $xyyz$ .



Since  $y \neq \varepsilon$ , the solitary  $b$  in  $w$  is more than half-way along  $xy^2z$ .

So  $xy^2z$  is not a palindrome.

This contradicts the conclusion of the Pumping Lemma applied to PALINDROME.

So our initial assumption, that PALINDROME is regular, is wrong.

Therefore PALINDROME is not regular.



$\{ \text{all languages} \}$   
 $\{ \text{regular languages} \}$   
**Assignment Project Exam Help**  
**<https://powcoder.com>**  
**Add WeChat powcoder**

Reading: Sipser, Ch. 1.

- ▶ closure properties: pp. 58–63.
- ▶ Pumping Lemma, non-regular languages: pp. 77–82.