

FIT2014 Theory of Computation

Assignment Project Exam Help

Lecture 5

Proofs: the Good, the Bad and the Ugly

<https://powcoder.com>

slides by Graham Farr

Add WeChat powcoder

COMMONWEALTH OF AUSTRALIA

Copyright Regulations 1969

Warning

This material has been reproduced and communicated to you by or on behalf of Monash University
in accordance with s113P of the Copyright Act 1968 (the Act).

The material in this communication may be subject to copyright under the Act.

Any further reproduction or communication of this material by you may be the subject of copyright protection under the Act.

Do not remove this notice.

Assignment Project Exam Help

- ▶ Good proofs
 - ▶ three proofs from The Book
- ▶ Bad proofs
- ▶ Ugly proofs

<https://powcoder.com>

Add WeChat powcoder

Good proofs

Theorem. (Euclid)

There are infinitely many prime numbers.

Proof.

Suppose, by way of contradiction, that there are only finitely many primes.

Let n be the number of primes.

Let p_1, p_2, \dots, p_n be all the primes.

Define: $q := p_1 \cdot p_2 \cdot \dots \cdot p_n + 1$.

This is bigger than every prime p_i . Therefore q must be composite.

Therefore q is a multiple of some prime.

But, for each prime p_i , if you divide q by p_i , you get a remainder of 1.

So q cannot be a multiple of p_i .

So q cannot be a multiple of any prime. **This is a contradiction.**

So our initial assumption was wrong.

So there are infinitely many primes.



Good proofs

Theorem. (Pythagoras)

$\sqrt{2}$ is irrational.

Proof.

Suppose, by way of contradiction, that $\sqrt{2}$ is rational.

Then, by definition, there exist positive integers m, n such that $\sqrt{2} = \frac{m}{n}$.

Among all such pairs m, n , choose a pair that have no common factors.

Squaring each side of our equation gives: $2 = \frac{m^2}{n^2}$.

Rewrite slightly: $2n^2 = m^2$.

This tells us that m^2 is even. Therefore m is even. Therefore $m = 2k$ for some k .

Substituting this back in gives $2n^2 = (2k)^2$, i.e., $2n^2 = 4k^2$, i.e., $n^2 = 2k^2$.

This tells us that n^2 is even. Therefore n is even.

Since m and n are both even, they both have a common factor, namely 2.

But we chose them so that they have no common factors. **This is a contradiction.**

Therefore our initial assumption, that $\sqrt{2}$ is rational, must be wrong.

Therefore $\sqrt{2}$ is *irrational*.



Good proofs

Definition: A set is **countable** if *either*

- ▶ it is finite, or
- ▶ it can be put in one-to-one correspondence (i.e., bijection) with \mathbb{N} .

Assignment Project Exam Help

FIT2014
students

{ Dakṣiṣputra, Muḥammad, Ramon,
Blaise, Gottfried, Nicole-Reine,
Maria, Charles, Ada, George,
Augustus, Annie, Henrietta,
Radhanath, Williamina, Kurt,
Rózsa, Alonzo, Alan, Konrad,
Bill, Tommy, Hedy, Trevor,
Maston, Noam, Winsome,
Stephen, Grace, John, Péisù,
Katherine, Margaret, Richard,
... }

finite

<https://powcoder.com>

Add WeChat powcoder

	\mathbb{N}	\mathbb{Z}	Σ^*
1	\longleftrightarrow 1	\longleftrightarrow 0	\longleftrightarrow ϵ
2	\longleftrightarrow 2	\longleftrightarrow 1	\longleftrightarrow a
3	\longleftrightarrow 3	\longleftrightarrow -1	\longleftrightarrow b
4	\longleftrightarrow 4	\longleftrightarrow 2	\longleftrightarrow aa
5	\longleftrightarrow 5	\longleftrightarrow -2	\longleftrightarrow ab
:	:	:	:
:	:	:	:

Good proofs

Theorem. (Cantor)

The set of *all languages* is *uncountable*.

Idea of proof: If {all languages} were countable ...

		ϵ	a	b	aa	ab	ba	bb	aaa	aab	...
1	\longleftrightarrow	L_1	✓	✓	✓	✓	✓	✓	✓	✓	...
2	\longleftrightarrow	L_2	✗	✓	✓	✓	✓	✓	✓	✓	...
3	\longleftrightarrow	L_3	✓	✓	✗	✗	✓	✗	✗	✓	...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots
m	\longleftrightarrow	L_m	✓	✗	✓	✗	✗	✓	✗	✗	...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\ddots

\hat{L} : ✗ ✓ ✓ ...

Good proofs

Theorem. (Cantor)

The set of *all languages* is *uncountable*.

Proof. Suppose, by way of contradiction, that the set of all languages is countable. Since we know it's not finite, there must be a bijection between \mathbb{N} and {all languages}. Let the members of the set of all languages be L_m , $m \in \mathbb{N}$. Recall that the set of all finite strings is countable, so we can list them as x_n , $n \in \mathbb{N}$. Define the language \hat{L} as follows

$$\forall n \in \mathbb{N} : x_n \in \hat{L} \Leftrightarrow x_n \notin L_n.$$

We have constructed \hat{L} so that, for each n , it differs from L_n in whether or not it contains x_n .

So it differs from all languages. Yet it is a language! **This is a contradiction.**

So our initial assumption was wrong.

So the set of languages is uncountable.



Bad proofs

From a falsehood, you can prove *anything*.

Recall the truth table of $P \Rightarrow Q$: always True when P is false, regardless of Q .

$$2+2 = 5$$

Therefore

$$4 = 5$$

Therefore

$$1 = 1$$

Now,

$$|\{ \text{McTaggart, The Pope} \}| = 2.$$

Therefore

$$|\{ \text{McTaggart, The Pope} \}| = 1.$$

Therefore

McTaggart is the Pope

<https://powcoder.com>

Add WeChat powcoder

attributed to G. H. Hardy in:
Harold Jeffreys, *Scientific Inference*,
Cambridge University Press, 1931/1957/1973.

Bad proofs

“Theorem”: Every graph has a cycle.

For all n : every graph on n vertices has a cycle.

This implies that trees do not exist.

“Proof”. We prove this by induction on the number of vertices.

1. Assume that **every graph on n vertices has a cycle**.
2. Let G be any graph on $n + 1$ vertices.
3. Let v be a vertex of G . Obtain the graph $G - v$ by removing v , and all its incident edges, from G .
4. Now, the graph $G - v$ has n vertices.
5. By the **Inductive Hypothesis**, $G - v$ has a cycle.
6. But, since $G - v$ is a subgraph of G , any cycle in $G - v$ is also a cycle in G .
7. Therefore G has a cycle.
8. Therefore, by Mathematical Induction, the result is true for all n .
So every graph has a cycle.

Assignment Project Exam Help

Definition: A string is *uniform* if all its letters are identical.

- ▶ i.e., it consists entirely of as or entirely of bs

- ▶ i.e., it's either $aa \dots a$ or $bb \dots b$

<https://powcoder.com>

Now, it is commonly thought that not all strings are uniform.

But we will now try to “prove”, by induction, that *all* strings are uniform!

Add WeChat powcoder

Bad proofs

“Theorem”: Every string over the alphabet $\{a,b\}$ is uniform.

“Proof”. We prove this by induction on the string length n .

1. Inductive basis: when $n = 1$, the string can only be “a” or “b”, and these are each of the required form, so the ‘theorem’ is true in this case.
2. Now assume $n \geq 2$, and suppose every string of length n is uniform.
3. Let w be any string of length $n + 1$.
4. Let w_1 be the string obtained from w by deleting the *first* letter of w , and let w_2 be the string obtained from w by deleting the *last* letter of w .
5. Both w_1 and w_2 are of length n .
6. By the Inductive Hypothesis, both w_1 and w_2 must be uniform.
7. w_1 and w_2 overlap in $n - 1$ letters. Since $n - 1 > 0$, this means that the number of letters shared by w_1 and w_2 is nonzero. So w_1 and w_2 must each consist entirely of the *same letter*, i.e., either they both consist entirely of as or they both consist entirely of bs.
8. It follows that w also consists entirely of as or entirely of bs, so it is uniform too.
9. The result follows for all n , by Mathematical Induction.

Ugly proofs

Theorem.

$\text{DOUBLEWORD} \subseteq \text{EVEN-EVEN}$

Assignment Project Exam Help

Proof. Let $w \in \text{DOUBLEWORD}$.

Assume w is not in EVEN-EVEN .

Then $w = xx$ for some word x .

So, $\# \text{ a's in } w = 2 \times (\# \text{ a's in } x)$, so it's even.

Also, $\# \text{ b's in } w = 2 \times (\# \text{ b's in } x)$, so it's even too.

This contradicts our assumption that w is not in EVEN-EVEN .

Therefore that assumption was wrong.

Therefore $w \in \text{EVEN-EVEN}$. \square

<https://powcoder.com>

Add WeChat powcoder

When you have a *direct* proof of your theorem, there's no need to dress it up as a proof by contradiction!

Ugly proofs?

A **colouring** of a graph G is a function that assigns a colour to each vertex of G such that *adjacent* vertices receive *different* colours.

- ▶ i.e., a function $f : V(G) \rightarrow \{\text{colours}\}$ such that $\forall u, v \in V(G) : u \sim v \Rightarrow f(u) \neq f(v)$.

A colouring is a **k -colouring** if the number of colours used is $\leq k$.

Applications:

- ▶ scheduling (timetabling)
- ▶ compilers (register allocation)
- ▶ communications (frequency assignment)

Theorem.

If G is planar then it has a 4-colouring.

Ugly proofs?

Theorem.

If G is planar then it has a 4-colouring.

Proofs:

- ▶ very long proof using computer to check 1476 configurations spanning 400 pages.
 - ▶ K. Appel and W. Haken, Every planar map is four colorable. I. Discharging, *Illinois Journal of Mathematics* **21** (3) (1977) 429–490.
 - ▶ K. Appel, W. Haken and J. Koch, Every planar map is four colorable. II. Reducibility, *Illinois Journal of Mathematics* **21** (3) (1977) 491–567.
- ▶ long proof using computer to check 633 configurations
 - ▶ N. Robertson, D. Sanders, P. Seymour and R. Thomas, The four-colour theorem, *Journal of Combinatorial Theory, Series B* **70** (1997) 2–44.
- ▶ proof by Robertson *et al.* (1997) formalised and formally verified by computer
 - ▶ G. Gonthier, Formal proof — the Four Color Theorem, *Notices of the American Mathematical Society*, **55** (11) (Dec. 2008) 1382–1393.

Ugly proofs?

Recall: to solve quadratic equations, $ax^2 + bx + c = 0$: $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$.

Formulas also exist for cubic and quartic equations. *But...*

Abel-Ruffini Theorem

There is no general algebraic formula (using arithmetic operations, powers & roots) for the roots of polynomials of degree ≥ 5 .

Incomplete proof, > 500 pages:

- ▶ Paolo Ruffini, *Teoria generale delle equazioni, in cui si dimostra impossibile la soluzione algebrica delle equazioni generali di grado superiore al quarto*, Stamperia di S. Tommaso d'Aquino, Bologna, 1799.

Complete proof, six pages:

- ▶ Niels Henrik Abel, *Mémoire sur les équations algébriques, ou l'on démontre l'impossibilité de la résolution de l'équation générale du cinquième degré*, Groendahl, Christiania (Oslo), 1824.