

Assignment Project Exam Help

Lecture 5: Regression Part 1

Spatial Data Science II

<https://powcoder.com>

Dr. Adams

Add WeChat powcoder

Assignment Project Exam Help

```
library(tidyverse)
```

```
> library(tidyverse)
— Attaching packages — tidyverse 1.2.1 —
✓ ggplot2 2.2.1 ✓ purrr 0.2.4
✓ tibble 1.3.4 ✓ dplyr 0.7.4
✓ tidyr 0.7.2 ✓ stringr 1.2.0
✓ readr 1.1.1 ✓ forcats 0.2.0
— Conflicts — tidyverse_conflicts() —
✖ dplyr::filter() masks stats::filter()
✖ dplyr::lag() masks stats::lag()
> |
```

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

1. Examining data distributions
2. Data Visualization
3. Data Management

- ▶ Tidy data

<https://powcoder.com>

Add WeChat powcoder

Big Bang

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

A measure of the dependence between two variables.

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help


- ▶ Measure of the strength of a linear relationship between two variables.
- ▶ Coefficient is represented with r
- ▶ Ranges from -1 to +1
- ▶ $H_0 : r = 0$
- ▶ $H_A : r \neq 0$

<https://powcoder.com>


Add WeChat powcoder

Examples


$r = -0.5$




$r = 0$



$r = 0.33$



$r = 0.85$



Assignment Project Exam Help
<https://powcoder.com>


Add WeChat powcoder

Pearson's r Assumptions

- ▶ Normally distributed
- ▶ Linear relationship

Assignment Project Exam Help

Linear assumption not met.



<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

- ▶ Spearman's ρ or r_s
- ▶ Non-parametric (distribution-free) rank statistic
- ▶ Relationship does not need to be linear
- ▶ Accepts non-interval data

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

It is possible for r to be positive while ρ is negative.


“Make sure not to overinterpret Spearman's rank correlation coefficient as a significant measure of the strength of the associations between two variables” (Hauke and Kossowski 2011)

Add WeChat powcoder

Spurious Correlations I

Assignment Project Exam Help

Per capita cheese consumption
correlates with
Number of people who died by becoming tangled in their bedsheets




Spurious Correlations II

Assignment Project Exam Help


Number of people who drowned by falling into a pool
correlates with

Films Nicolas Cage appeared in




Spurious Correlations III

Assignment Project Exam Help



Assignment Project Exam Help



<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

How much does a house price increase when we increase its square footage?

- ▶ Take a minute and think about how you could answer this question?

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

Model a continuous variable as a linear function of one or more independent variables.

- ▶ This allows us to understand if and how an attribute contributes to an outcome.

<https://powcoder.com>

Add WeChat powcoder

Model Use In Data Science




Fig. 1: Top Data Science, Machine Learning Methods Used, 2018/2019

Assignment Project Exam Help

A statistical model that

- ▶ Predicts a continuous variable: Y
- ▶ Using one or more independent variables: x_n
- ▶ Calculates a set of multipliers: β_i
 - ▶ Regression coefficients
- ▶ Includes an intercept: β_0

The linear regression formula is:

$$Y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 \dots \beta_n x_n + \epsilon$$

ϵ is our error or noise.

<https://powcoder.com>


Add WeChat powcoder

Linear Regression Visualized

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



Assignment Project Exam Help

- ▶ House Price (\$)
- ▶ Number of bedrooms
- ▶ Square Footage
- ▶ Number of bathrooms

<https://powcoder.com>

Add WeChat powcoder

House Sales Data

```
library(tidyverse)
house = read_delim(
  "http://www.rossmanchance.com/iscam2/data/housing.txt",
  delim = "\t") # Tab delimited
```

```
## Parsed with column specification:
## cols(
##   sqft = col_integer(),
##   price = col_integer(),
##   City = col_character(),
##   bedrooms = col_integer(),
##   baths = col_double()
## )
```

Take a look at our data

```
## # A tibble: 83 x 5
```

```
##   sqft    price City bedrooms baths  
##   <int>  <int> <chr>      <int> <dbl>
```

```
## 1   3392 339000 Dublin           3  2.10
```

```
## 2   4100 899900 pleasanton       4  3.00
```

```
## 3   3200 448641 Grayton         5  4.00
```

```
## 4   1436 239999 Moraga           4  3.00
```

```
## 5   1944 377500 Antioch          3  2.00
```

```
## 6   1500 299900 Danville          3  2.50
```

```
## 7   1700 265000 El Dorado Hills       4  3.00
```

```
## 8   2507 449000 Shingle Springs       4  3.00
```

```
## 9   1580 439950 McKinleyville          3  2.00
```

```
## 10  1500 699888 Marina           4  2.00
```

```
## # ... with 73 more rows
```

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

- ▶ House Price (price)
- ▶ Number of bedrooms (bedrooms)
- ▶ Square Footage (sqft)
- ▶ Number of bathrooms (baths)

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

Before we fit our model we need to ensure the data fits the linear regression model's assumptions


1. Linearity between independent variables (x_n) and dependent variable (Y)
2. No outliers in x_n
3. Normally Distributed x_n & Y

<https://powcoder.com>
Add WeChat powcoder


Linearity: Price and Square Footage

Assignment Project Exam Help


```
ggplot(data = house, mapping = aes(y = price, x = sqft)) +  
  geom_point() +  
  geom_smooth(method = "lm", se = F)
```



Linearity: Price and Baths



Linearity: Price and Bedrooms



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

- ▶ Data values outside of $1.5 * \text{interquartile-range}$ may be considered outliers
- ▶ The IQR is the distance from the 25th percentile to the 75th percentile.
- ▶ We often visualize this with the box and whisker plot

<https://powcoder.com>
Add WeChat powcoder

Assignment Project Exam Help


Box and whisker plots:

- ▶ bottom and top of the box are the first and third quartiles
- ▶ band inside the box is the second quartile (the median)
- ▶ `geom_boxplot`:
 - ▶ Whiskers are largest or smallest value within $1.5 * IQR$
 - ▶ Points are outside of $1.5 * IQR$

Add WeChat powcoder

Box Plot Visualized

Assignment Project Exam Help




<https://powcoder.com>

Add WeChat powcoder

Check for outliers: Square Footage

Assignment Project Exam Help


```
ggplot(data = mtcars) +  
  geom_boxplot(mapping = aes(y = sqft, x = ""))
```



Distribution: Price

Assignment Project Exam Help


```
ggplot(data = mtcars) +  
  geom_histogram(mapping = aes(x = price))
```



Distribution: $\log(\text{Price})$

Assignment Project Exam Help

```
ggplot(data = house)
  geom_histogram(mapping = aes(x = price)) +
  scale_x_log10()
```






Distribution: Independent Variables

```
par(mfrow=c(1, 3))  
hist(house$sqft); hist(house$bedrooms); hist(house$baths)
```

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



Transform Variables

```
house%>%  
  mutate(price, log_price = log(price))%>%  
  mutate(sqft, log_sqft = log(sqft))%>%  
  mutate(baths, log_baths = log(baths)) -> house  
house%>%  
  select(log_baths)
```

```
## # A tibble: 83 x 1
```

```
##   log_baths
```

```
##   <dbl>
```

```
## 1 0.742
```

```
## 2 1.10
```

```
## 3 1.39
```

```
## 4 1.10
```

```
## 5 0.693
```

```
## 6 0.916
```

```
## 7 1.10
```

Assignment Project Exam Help



<https://powcoder.com>

Add WeChat powcoder

Transformed

Assignment Project Exam Help

```
par(mfrow=c(1, 2))  
hist(house$baths); hist(house$log_baths)
```



Assignment Project Exam Help

- ▶ Scatter plot: Check for linear relationships between x_n and Y
- ▶ Box plot: Outlier check
- ▶ Histogram: Check variables for normal distributions

Add WeChat powcoder

Assignment Project Exam Help

- ▶ It is good if a x_n is correlated with Y
- ▶ Problematic when multiple x_n are correlated
 - ▶ Multicollinearity
 - ▶ We will address later on in the lecture

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

```
cor_mat <- cor(house%>%  
  select(bedrooms, log_price, sqft, log_baths))  
cor_mat
```

<https://powcoder.com>

```
##           bedrooms log_price      sqft log_baths  
## bedrooms  1.0000000 0.3188020 0.5869470 0.7657905  
## log_price 0.3188020 1.0000000 0.6638824 0.3554938  
## sqft      0.5869470 0.6638824 1.0000000 0.6068155  
## log_baths 0.7657905 0.3554938 0.6068155 1.0000000
```

Add WeChat powcoder


Visualize Correlation: Plots

Assignment Project Exam Help

```
corrplot::corrplot(cor_mat, cli_pos = "bl", tl_pos = "d", add
```

<https://powcoder.com>

Add WeChat powcoder



Assignment Project Exam Help

- ▶ The tidyverse has yet to really address statistical modelling
 - ▶ “You can see some of the pieces in the recipes and rsample packages but we do not yet have a cohesive system that solves a wide range of challenges. This work will largely replace the modelr package used in R4DS.”- Tidyverse site
- ▶ Broom is package that may be helpful.

Add WeChat powcoder

Fit the linear regression model

Assignment Project Exam Help

- ▶ The function we use in R is `stats::lm()`

- ▶ The formula is:

- ▶ $\text{dep} \sim \text{indep}_1 + \text{indep}_2 + \dots + \text{indep}_n$

$$Y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_n x_n + \epsilon$$

?lm: lm is used to fit linear models. It can be used to carry out regression, single stratum analysis of variance and analysis of co-variance (although aov may provide a more convenient interface for these).

Assignment Project Exam Help

Formulas use the tilde (by) and + (plus) characters:

`Y~var1+var2+var3+...varN`

Example

```
lm(hwy~displ+year+cyl, data = mpg)
```

<https://powcoder.com>
Add WeChat powcoder

Assignment Project Exam Help

```
house_reg <- lm(log_price ~ sqft, data = house)
house_reg
```

<https://powcoder.com>

Call:

lm(formula = log_price ~ sqft, data = house)

##

Coefficients:

(Intercept) sqft

1.204e+01 4.274e-04

Add WeChat powcoder

summary(lm())

```
summary(house_reg)
```

```
Call:
lm(formula = log_price ~ sqft, data = house)
```

```
Residuals:
```

```
      10% Median      30%      90%      Max
-1.08588 -0.19591 -0.05899  0.28717  1.20206
```

```
Coefficients:
```

```
              Estimate Std. Error t value Pr(>|t|)
(Intercept) 1.24e+11  1.26e-01  97.36  7.7e-16 ***
sqft        4.274e-04  5.549e-05   7.99  7.87e-12 ***
```

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Residual standard error: 0.4502 on 81 degrees of freedom
```

```
Multiple R-squared:  0.4407,    Adjusted R-squared:  0.4338
```

```
F-statistic: 63.83 on 1 and 81 DF,  p-value: 7.874e-12
```

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

An object of class "lm" is a list containing at least the following components:

- ▶ coefficients: a named vector of coefficients
- ▶ residuals: the residuals, that is response minus fitted values.
- ▶ fitted.values: the fitted mean values.

```
house$residuals[1:5]
```

Add WeChat powcoder

Assignment Project Exam Help

1. Are coefficients statistically significant?

- ▶ Check with the coefficient p -value

- ▶ Uses the t -value

2. Is the model statistically significant, overall p -value

- ▶ Check with the coefficient p -value

- ▶ Uses the F -test

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

The statistical significance of each coefficient is tested with the t-value

$$t = \frac{\text{coefficient}}{\text{std. error}}$$

<https://powcoder.com>

- ▶ Should be greater 1.96 for p-value to be less than 0.05
- ▶ We reject the null hypothesis when $p < 0.05$
- ▶ When $p > 0.05$ we remove this variable from the model.

Add WeChat powcoder

Assignment Project Exam Help

- ▶ The F-statistic assess the overall model
- ▶ Null hypothesis:
 - ▶ An equal fit of the model with a model with zero predictors.
- ▶ Alternative hypothesis:
 - ▶ This model perform better than an intercept only model.
- ▶ If the p-value associated to the F-statistic is < 0.05 we reject

H_0

Add WeChat powcoder

Refresher on p-values if you need it

<https://www.youtube.com/watch?v=128yz0OCG-I>

Type I Error:

- ▶ Incorrect rejection of a true null hypothesis
 - ▶ False positive

Type II Error

- ▶ Incorrectly retaining a false alternative hypothesis
 - ▶ False negative

Alpha

As we decrease our chance of a Type I error, we increase our risk of Type II

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

R^2 tells us is the proportion of variation in the dependent (response) variable that has been explained by this model.

The adjusted R^2 accounts for the effect that occurs when you add more independent variables that your R^2 increases.

- ▶ Increases only if a new term improves the model more than expected by chance.
- ▶ Decreases when a predictor improves the model by less than expected by chance.

Assignment Project Exam Help

Checking your model:

1. The mean of the residuals is zero
2. Homoscedasticity of residuals or equal variance
3. Multicollinearity
4. The x_n variables and residuals are uncorrelated
5. The variability in X values is positive
6. The number of observations must be greater than number of x_n
7. Normality of residuals
8. No auto-correlation of residuals

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

The mean of the residuals is zero. Residuals difference in model estimates and actual values.

```
mean(house_reg$residuals)
```

```
## [1] 2.70306e-17
```

Add WeChat powcoder

Homoscedasticity: Equal variance across values

Homoscedasticity:

Requires variance of residuals to be the same across the fitted values.





Heteroscedasticity:

- ▶ When the size of the error term differs across values of an independent variable.
 - ▶ Violation of homoscedasticity
- ▶ Linear Regression (Ordinary Least Squares), seeks to minimize residuals
- ▶ OLS equally weights all observations
 - ▶ Cases with larger errors have more effect on the model estimation.

Checking with `plot(lm(y~x, data = data))`

```
par(mfrow=c(2,2)) # set 2 rows and 2 column plot layout
plot(house_reg)
```

Assignment Project Exam Help



```
# Breusch-Pagan test  
bptest(house_reg)  
car::ncvTest(house_reg)
```

1. Try different predictors
2. Variable transformation
 - ▶ Box-Cox

3. Select a different regression model (last case)

```
library(caret)  
bc <- BoxCoxTrans('values')  
predict(bc, 'values')
```


No autocorrelation of residuals

Assignment Project Exam Help

- ▶ Temporal Autocorrelation
 - ▶ The value at one point is not dependent on the previous value
- ▶ Spatial Autocorrelation
 - ▶ Values at one location are not dependent on near values
 - ▶ Moran's I

<https://powcoder.com>
Add WeChat powcoder

The x_n variables and residuals are uncorrelated

```
cor.test(house$sqft, house_reg$residuals)
```

Assignment Project Exam Help

```
##  
## Pearson's product-moment correlation  
## data: house$sqft and house_reg$residuals  
## t = -4.7743e-16, df = 81, p-value = 1  
## alternative hypothesis: true correlation is not equal to 0  
## 95 percent confidence interval:  
## -0.2156893 0.2156893  
## sample estimates:  
## cor  
## -5.304794e-17
```

<https://powcoder.com>

Add WeChat powcoder

The variability in x_n values is positive

Assignment Project Exam Help

```
var(house$sqft)
```

```
## [1] 863896.6
```

This is much greater than 0.

We cannot use a x_n variable with a single value.

<https://powcoder.com>

Add WeChat powcoder

The number of observations must be greater than number of Xs


Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Normality of residuals

Assignment Project Exam Help



No perfect multicollinearity

Assignment Project Exam Help

Occurs when we have 2 or more predictor variables.

- ▶ Assessed using VIF
 - ▶ Variance inflation factors
- ▶ Rule of thumb $VIF > 4$
 - ▶ You should revise your variable selection

<https://powcoder.com>

```
cars.vif()
```

Add WeChat powcoder

```
car::mtcars
```

- ▶ mpg: Miles/(US) gallon
- ▶ cyl: Number of cylinders
- ▶ disp: Displacement (cu.in)
- ▶ hp: Gross horsepower
- ▶ drat: Rear axle ratio
- ▶ wt: Weight (1000 lbs)
- ▶ qsec: 1/4 mile time
- ▶ vs: V/S
- ▶ am: Transmission (0 = automatic, 1 = manual)
- ▶ gear: Number of forward gears
- ▶ carb: Number of carburetors

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

```
library(car)
# We will include all variables in the model
mpg_lm <- lm(mpg ~ ., data=mtcars)
vif(mpg_lm)
```

<https://powcoder.com>

```
##           cyl           disp           hp           drat           wt           c
## 15.373333 21.620241 9.832037 3.374620 15.164887 7.527
##           am           gear           carb
## 4.648487 5.357452 7.908747
```

Add WeChat powcoder

Assignment Project Exam Help

```
mpg_lm_2 <- lm(mpg ~ cyl + gear + am, data=mtcars)  
vif(mpg_lm_2)
```

<https://powcoder.com>

```
##          cyl          gear          am  
## 1.407382 2.768828 2.884543
```

Add WeChat powcoder

Assignment Project Exam Help

- ▶ Increase the variance of the coefficient estimates
- ▶ Estimates may be very sensitive to minor changes in the model
- ▶ Statistical power is reduced
- ▶ Coefficient sign switching

<https://powcoder.com>

“A caution must be taken when more than two predictors in the model have even weak pairwise correlation coefficients ($r=0.25$) as they can result in a significant multicollinearity effect” (P. Vatcheva and Lee 2016 ,pg. 13)

Add WeChat powcoder

Assignment Project Exam Help

Hauke, Jan, and Tomasz Kossowski. 2011. "Comparison of values of pearson's and spearman's correlation coefficients on the same sets of data." *Quaestiones Geographicae* 30 (2): 87–93.

doi:10.2478/v18117-011-0021-1.
<https://powcoder.com>

P. Vatcheva, Kristina, and MinJae Lee. 2016. "Multicollinearity in Regression Analyses Conducted in Epidemiologic Studies."

Epidemiology: Open Access 06 (02): 1–20.
[Add WeChat powcoder](https://powcoder.com)
doi:10.4172/2161-1165.1000227.