

Propositional Logic
Assignment Project Exam Help

cntd
<https://powcoder.com>

Add WeChat powcoder
Fariba Sadri

What we have done so far on Propositional Logic

- Syntax of wffs
- Practice on how to formalise English sentences in propositional logic
- Truth tables for the semantics of the connectives $\neg, \wedge, \vee, \rightarrow, \leftrightarrow$
- Tautologies, inconsistencies, contingencies
- Equivalences

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

What we will do now in this set of slides

- Semantic consequence
 - Natural deduction proofs
 - Soundness and completeness
- Assignment Project Exam Help
<https://powcoder.com>
Add WeChat powcoder

Recap Exercise

From 2012-13 examination paper:

Assignment Project Exam Help

a. Define a new connective \otimes for *exclusive-or*, using any (combination) of the usual connectives, \wedge , \vee , \neg , \rightarrow , \leftrightarrow . Thus $p \otimes q$ is to mean either p or q but not both.

<https://powcoder.com>

Add WeChat powcoder

b. Use the new connective \otimes together with any of the other usual connectives to express the following sentences in propositional logic, where *either ... or* is to be understood as *exclusive-or*. The propositions to be used are given in the text in *Italics* inside brackets.

Assignment Project Exam Help

Either John will leave the company (*jL*) or Mary will (*mL*).

If John leaves then either the tax department will close (*closeTax*), or Peter will be shared between two departments (*pShare*) and an administrator will be recruited (*recruitA*).

If Mary leaves then either an administrator will be recruited or a secretary will be recruited (*recruitS*), provided John is shared between two departments (*jShare*).

Definition:

Semantic Consequence

Let

S be a set of wffs, and

W be a wff.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

If whenever all the wffs in S are true W is also true, then W is a **semantic consequence** of S.

Semantic Consequence cntd.

Denoted as

Assignment Project Exam Help

" \models " is the semantic turnstile

(a metasyMBOL)

Add WeChat powcoder

We also say W is **semantically entailed** by S .

If W is a tautology then

$\models W$.

Exercise



Show the following:

- a. $A \wedge B \vdash A \vee B$
- b. $\text{snow, mild} \rightarrow \text{snow} \vdash \text{mild}$
- c. Go back to one of the argument at the beginning of the notes you think is valid and show that the conclusion of the argument is semantically entailed by the premises.

Definitions: Valid, Satisfiable

- *Valid* is just another name for tautology.

So a formula is valid if it is true in every interpretation.

$\models A$ if A is valid.

- A formula is *satisfiable* if it is true in at least one interpretation.

	Validity	Satisfiability
$A \models B$?? valid	?? unsatisfiable
$A \models B$ and $B \models A$?? valid	?? unsatisfiable

Inference

Example: Given

(pass_exams \wedge pass_projects) \rightarrow pass_MSc

pass_exams <https://powcoder.com>

pass_projects Add WeChat powcoder

one can infer (conclude)

pass_MSc.

Example: Given

Assignment Project Exam Help
 $\text{thursday} \rightarrow \text{logic_lecture}$

$\neg \text{logic_lecture}$ <https://powcoder.com>

Add WeChat powcoder

Can you infer

$\neg \text{thursday} ?$

The “elections” example

Given

- If there are national elections then the Tory party wins or the Labour party wins.
- If the unions do not support the Labour party then it does not win.
- There are national elections.

Can you infer

If the Tory party does not win then the unions support the Labour party?

The “elections” example: Formalisation in logic

Given

Elections <https://powcoder.com> **Assignment Project Exam Help**
→ Labour_wins ∨ Tory_wins
¬ Unions_support_Labour → Labour_wins
Elections **Add WeChat powcoder**

can you infer

¬ Tory_wins → Unions_support_Labour ?

The “elections” example: Abbreviation

Premise:

Assignment Project Exam Help

1. $E \rightarrow L \vee T$

2. $\neg U \rightarrow \neg L$

3. Add WeChat powcoder

Conclusion:

$\neg T \rightarrow U$

You can try to use truth tables to see if the conclusion is semantically entailed by the premises.

<https://powcoder.com>

How many rows?

Too many!

E	L	T	U	$L \vee T$	$E \rightarrow L \vee T$	$\neg U$	$\neg L$	$\neg U \rightarrow \neg L$	$\neg T$	$\neg T \rightarrow U$
---	---	---	---	------------	--------------------------	----------	----------	-----------------------------	----------	------------------------

T	T	T	T	T	T	F	F	T	F	T
---	---	---	---	---	---	---	---	---	---	---

.....

..

..

..

.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Can you give an informal proof of the
conclusion from the premises without using
truth tables?

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder




Performing inferences is very important in many applications of logic

Argument: Premise  Conclusion

Modelling : Theory  Consequences

Programming: <https://powcoder.com>

Specification  Program

Specification  Properties

Prolog:

Program  Answers to queries

Rules of Inference

Natural Deduction

(Reasoning purely at the syntactic level)

\wedge -elimination ($\wedge E$) Assignment Project Exam Help

$$\frac{X \wedge Y}{X}$$

$$\frac{X \wedge Y}{Y}$$

<https://powcoder.com>
Add WeChat powcoder

\wedge -introduction ($\wedge I$)

$$\frac{X, Y}{X \wedge Y}$$

$$\frac{X, Y}{Y \wedge X}$$

\vee -elimination ($\vee E$)

$$\frac{X \vee Y, \neg X}{Y} \quad \frac{X \vee Y, \neg Y}{X}$$

<https://powcoder.com>

Add WeChat powcoder

\vee -introduction ($\vee I$)

$$\frac{X}{X \vee Y} \quad \frac{X}{Y \vee X}$$

\rightarrow -elimination (\rightarrow E) (Modus Ponens)

$X, X \rightarrow Y$ Assignment Project Exam Help

Y

<https://powcoder.com>

\rightarrow -introduction (\rightarrow I)

X Assume Add WeChat powcoder

.

.

Y

$X \rightarrow Y$

\neg -elimination and \neg -introduction (Reductio Ad Absurdum) (RAA) (Proof by contradiction)

$\neg X$ assume

·

·

·

$\frac{Y, \neg Y}{X}$

X

<https://powcoder.com>

Add WeChat powcoder

X assume

·

·

·

$\frac{Y, \neg Y}{\neg X}$

$\neg X$

Note: X and Y may be the same wff.

$\neg X$ **assume** X **assume**

• <https://powcoder.com> •

• Add WeChat powcoder •

•

$$\frac{\neg X, X}{X}$$

•

$$\frac{X, \neg X}{\neg X}$$

\leftrightarrow -introduction (\leftrightarrow I)

Assignment Project Exam Help

$$\frac{X \rightarrow Y, Y \rightarrow X}{X \leftrightarrow Y}$$

<https://powcoder.com>

Add WeChat powcoder

\leftrightarrow -elimination (\leftrightarrow E)

$$\frac{X \leftrightarrow Y}{X \rightarrow Y}$$

$$\frac{X \leftrightarrow Y}{Y \rightarrow X}$$

Note: In all the inference rules X and Y stand for any wffs. So the following, for example, is an application of the \rightarrow elimination rule:

<https://powcoder.com>

Given $A \wedge (B \vee C)$ and $(A \wedge (B \vee C)) \rightarrow ((A \rightarrow D) \vee (\neg E \wedge F))$

we can infer

$$(A \rightarrow D) \vee (\neg E \wedge F)$$

Example

You may be wondering why we need the $\forall I$ rule. Here is an example that uses it.

<https://powcoder.com>

Example:

Add WeChat powcoder

If there is a shortage of petrol or the tax on petrol is high then people are angry. There is a shortage of petrol.

So people are angry.

Premise

1. $(SP \vee HT) \rightarrow \text{Anger}$

2. SP

we want to conclude

Anger.

1. $(SP \vee HT) \rightarrow \text{Anger}$ **given**
2. SP **Assignment Project Exam Help** **given**

<https://powcoder.com>

?

Add WeChat powcoder

Anger

1. $(SP \vee HT) \rightarrow \text{Anger}$ **given**

2. SP **Assignment Project Exam Help**
given

<https://powcoder.com>

?

Add WeChat powcoder

Anger

$\rightarrow E$

1. $(SP \vee HT) \rightarrow \text{Anger}$ given
2. SP Assignment Project Exam Help
given

<https://powcoder.com>

?

Add WeChat powcoder

$SP \vee HT$

$\text{Anger} \rightarrow E$

Proof

1. $(SP \vee HT) \rightarrow \text{Anger}$ given
2. SP Assignment Project Exam Help given
3. $SP \vee HT$ <https://powcoder.com> 2, \vee I
4. Anger Add WeChat powcoder 1,3, \rightarrow E

Example

Derive $P \vee Q$ from $P \wedge Q$.

Assignment Project Exam Help

1. $P \wedge Q$ <https://powcoder.com> given

?????? Add WeChat powcoder

$P \vee Q$

Derive $P \vee Q$ from $P \wedge Q$.

Assignment Project Exam Help

1. $P \wedge Q$ <https://powcoder.com> given

?????? Add WeChat powcoder

$P \vee Q$

$\vee I$

Example

Derive $P \vee Q$ from $P \wedge Q$.

Assignment Project Exam Help

1. $P \wedge Q$ <https://powcoder.com> given
2. P Add WeChat powcoder 1, $\wedge E$
3. $P \vee Q$ 2, $\vee I$

Example

Derive **R** from **P, Q, $(P \wedge Q) \rightarrow R$** .
Assignment Project Exam Help

1. **P** <https://powcoder.com>
given
2. **Q** Add WeChat powcoder
given
3. **$(P \wedge Q) \rightarrow R$** given

??????

R

1. P **given**

2. Q **Assignment Project Exam Help**
given

3. $(P \wedge Q) \rightarrow R$ **https://powcoder.com**
given

?????? **Add WeChat powcoder**

R $\rightarrow E$

1. P given
2. Q given
3. $(P \wedge Q) \rightarrow R$ given
4. $P \wedge Q$ 1,2, $\wedge I$
5. R 3,4, $\rightarrow E$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Example

Derive $Q \rightarrow R$ from $P, (P \wedge Q) \rightarrow R$.

Assignment Project Exam Help

1. P <https://powcoder.com> given
2. $(P \wedge Q) \rightarrow R$ given
?????
?????
?????
 $Q \rightarrow R$

1. P **given**

2. $(P \wedge Q) \rightarrow R$ **given**

?????

<https://powcoder.com>

?????

Add WeChat powcoder

?????

$Q \rightarrow R$

$\rightarrow I$

1. P

given

2. $(P \wedge Q) \rightarrow R$

given

Assignment Project Exam Help

Q <https://powcoder.com>

assume

?? Add WeChat powcoder

R

$Q \rightarrow R$

$\rightarrow I$

1. P

given

2. $(P \wedge Q) \rightarrow R$

Assignment Project Exam Help

given

Q

<https://powcoder.com>

assume

???

Add WeChat powcoder

R

\rightarrow E

6. $Q \rightarrow R$

3, 5, \rightarrow I

1. P

given

2. $(P \wedge Q) \rightarrow R$

given

3.

Q <https://powcoder.com>

assume

4.

$P \wedge Q$ Add WeChat powcoder

1, 3, \wedge I

5.

R

2, 4, \rightarrow E

6. $Q \rightarrow R$

3, 5, \rightarrow I

Example

Derive $\neg R$ from $P \wedge Q, R \rightarrow \neg P$.

Assignment Project Exam Help

1. $P \wedge Q$ <https://powcoder.com> given

2. $R \rightarrow \neg P$ given


Add WeChat powcoder

????

????

????

$\neg R$



Assignment Project Exam Help

1. $P \wedge Q$

<https://powcoder.com> given

2. $R \rightarrow \neg P$


Add WeChat powcoder given

????

????

????

$\neg R$



RAA

1. $P \wedge Q$

given

2. $R \rightarrow \neg P$

Assignment Project Exam Help
given

<https://powcoder.com>

R

Add WeChat powcoder

assume

????

????

$\neg R$

RAA

1. $P \wedge Q$

given

2. $R \rightarrow \neg P$

Assignment Project Exam Help

given

3. P

<https://powcoder.com>

1, $\wedge E$

4. R Add WeChat powcoder assume

5. $\neg P$

2, 4, $\rightarrow E$

6. $\neg R$

3, 4, 5, RAA



\vdash

$P \vdash W$

denotes W is (syntactically) derivable from P .

\vdash is called the syntactic turnstile. It is a symbol in the metalanguage.

Example:

In the last example:


$P \wedge Q, R \rightarrow \neg P \vdash \neg R$

Definition

A **derivation** or **proof** of a wff W in propositional logic from Assignment Project Exam Help premises, is a finite sequence of wffs such that the last wff is W and each wff in the sequence is one of the following: <https://powcoder.com> Add WeChat powcoder

- a premise, i.e. a wff in P
- an immediate consequence of one or more wffs preceding it in the sequence, as determined by one of the inference rules of propositional logic.
- An assumption (that is later discharged by an application of $\rightarrow I$ or RAA).

Exercise



Show

$A \rightarrow B, B \rightarrow C \vdash A \rightarrow C$


Assignment Project Exam Help

<https://powcoder.com>

(Transitivity of the implication.)

Add WeChat powcoder

Exercise



Show

$Q \rightarrow R \vdash (P \rightarrow Q) \rightarrow (P \rightarrow R)$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Exercise

Consider the following derivation:

1. A Assignment Project Exam Help
assume
2. $A \rightarrow A$ https://powcoder.com
1, 1, \rightarrow I
3. A Add WeChat powcoder
1, 2, \rightarrow E

Seemingly this proves $\vdash A$.

Is there anything wrong with it?

If so, what?

Exercise

Show

Assignment Project Exam Help
 $\text{snow, mild} \rightarrow \neg \text{snow} \vdash \neg \text{mild}$
<https://powcoder.com>

It is enough to show: Add WeChat powcoder

$p, q \rightarrow \neg p \vdash \neg q$

Some useful derived inference rules

Double negation elimination ($\neg\neg E$)

Assignment Project Exam Help

$$\frac{\neg\neg X}{X}$$

<https://powcoder.com>

Add WeChat powcoder

Double negation introduction ($\neg\neg I$)

$$\frac{X}{\neg\neg X}$$

Law of excluded middle

Assignment Project Exam Help

$$\overline{X \vee \neg X}$$

<https://powcoder.com>

Add WeChat powcoder

Proof by cases

$$\frac{X \vee Y, X \rightarrow Z, Y \rightarrow Z}{Z}$$

Modus Tollens

$$\frac{X \rightarrow Y, \neg Y}{\neg X}$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Contraposition

$$\frac{X \rightarrow Y}{\neg Y \rightarrow \neg X}$$

Dilemma


$$\frac{X \rightarrow Y, \neg X \rightarrow Y}{Y}$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Exercises



- Give a formal proof for the “Exercises” example.

<https://powcoder.com>

- Using the basic inference rules ($\wedge I$, $\wedge E$, $\vee I$, $\vee E$, $\rightarrow I$, $\rightarrow E$, $\leftrightarrow I$, $\leftrightarrow E$, $\neg I$, $\neg E$, $\neg E$, $\neg E$), show that the derived inference rules hold.

Be careful when you use assumptions in a derivation

Show $\vdash \neg(\neg A \wedge \neg B) \rightarrow (A \vee B)$

1. $\neg(\neg A \wedge \neg B)$ Assignment Project Exam Help

2. $\neg(A \vee B)$

assume

3. $\neg A$

assume

4. $\neg B$ assume
Add WeChat powcoder

5. $\neg A \wedge \neg B$ 3,4, \wedge I

6. B

4,5,1,RAA

7. $A \vee B$

6, \vee I

8. A

3,2,7,RAA

9. $A \vee B$

8, \vee I

10. $A \vee B$

2,9,RAA


11. $\neg(\neg A \wedge \neg B) \rightarrow (A \vee B)$

1,10, \rightarrow I

Notes

- The only inference rules that make use of assumptions are \rightarrow and \rightarrow Elimination. Assignment Project Exam Help
- It is very important to be clear about the scope of assumptions. <https://powcoder.com>
- Any assumption made during a derivation will remain in force, and ultimately count as one of the premises for the conclusion, unless it gets discharged before the conclusion is reached in the proof. Add WeChat powcoder

Exercise



Show Assignment Project Exam Help

$\vdash ((P \wedge Q) \vee (\neg P \wedge R)) \rightarrow (Q \vee R)$

Add WeChat powcoder

Exercise

Show

$P, \neg P \vdash Q$

Assignment Project Exam Help

<https://powcoder.com>

Note:

Add WeChat powcoder

This exercise shows that anything can be derived from an inconsistent set of premises.

Notes

➤ $\vdash W$ denotes W is derivable from an empty set of premises.

➤ Let A, B be wffs.

If $A \vdash B$ then $\vdash A \rightarrow B$.

If $\vdash A \rightarrow B$ then $A \vdash B$.

In general if

P is a set of wffs, and

P' is a conjunction of the wffs in P , and

W is a wff then $P \vdash W$ iff $\vdash P' \rightarrow W$

Notes cntd.

- Proofs (derivations) are independent of the "meaning" of the propositional symbol.

Assignment Project Exam Help

<https://powcoder.com>

So a proof is still valid if the symbols are replaced consistently.

Add WeChat powcoder

Example: If we have a proof for

$P, Q \rightarrow \neg P \vdash \neg Q$

Then the following also holds (replacing P with snow and Q with mild)

$\text{snow}, \text{mild} \rightarrow \neg \text{snow} \vdash \neg \text{mild}$

$\text{winter} \rightarrow \text{cold}, \text{globalWarming} \rightarrow \neg(\text{winter} \rightarrow \text{cold})$
 $\vdash \neg \text{globalWarming}$

Notes cntd.

- For convenience, in a derivation we can use instances of previous derivations. That is, if we have previously shown

$S \vdash W$,

<https://powcoder.com>

Add WeChat powcoder

and we are now attempting a new proof for another wff, but we have so far shown

an instance of S , then we can write down the same
instance of W in the derivation without reproducing its entire proof.



Exercise A

Show [Assignment Project Exam Help](https://powcoder.com)

$P \rightarrow Q, R \rightarrow S \vdash (P \vee R) \rightarrow (Q \vee S).$

[Add WeChat powcoder](https://powcoder.com)

Exercise B

Show

$(P \vee Q) \vee R$	\vdash	$P \vee (Q \vee R)$	and
$P \vee (Q \vee R)$	\vdash	$(P \vee Q) \vee R.$	



Exercise C

Formalise the following argument and show that it is valid.
You may use the theorems in A and B, above.

<https://powcoder.com>
In Britain one of the three parties, Tory, Labour or Liberal Democrat, is in power.

[Add WeChat powcoder](#)
If the Tories are in power the government may support cuts in public spending.

If Labour is in power the government may support tax increases.

If the Liberal Democrats are in power the government may support proportional representation.

So in Britain the government may support cuts in public spending or tax increases or proportional representation.

Soundness and Completeness

Propositional logic is both **sound** and **complete**.

Let S be any set of wffs, and let W be a wff.

<https://powcoder.com>

Soundness means the following:

If $S \vdash W$ then $S \models W$.

Completeness means the following:

If $S \models W$ then $S \vdash W$.

So in propositional logic we are justified in switching between syntactic proofs and semantic consequences.

Assignment Project Exam Help

<https://powcoder.com>

Example

Add WeChat powcoder

$A \equiv B$ iff

$A \models B$ and $B \models A$ iff

$A \vdash B$ and $B \vdash A$ iff

$A \vDash B$ and $B \vDash A$


It also means in proofs we can use
equivalences.

Assignment Project Exam Help

<https://powcoder.com>

But note in assessments you need to check the
specifications in the questions carefully.

Exercise



We know

$$P \rightarrow Q \equiv \neg P \vee Q$$

Assignment Project Exam Help

So

<https://powcoder.com>

$$P \rightarrow Q \vdash \neg P \vee Q$$

$$\neg P \vee Q \vdash P \rightarrow Q$$

Add WeChat powcoder

Also

$$\vdash (P \rightarrow Q) \leftrightarrow (\neg P \vee Q)$$

As an exercise show the last using inference rules.

Exercises



Given the equivalence

$$A \wedge (B \wedge C) \equiv (A \wedge B) \wedge C$$

$$(P \wedge Q) \rightarrow R, \neg(R \wedge S) \vdash \neg(P \wedge (Q \wedge S)).$$

<https://powcoder.com>

Using the equivalences

$$A \rightarrow B \equiv \neg A \vee B \quad \text{and}$$

$$\neg(A \vee B) \equiv \neg A \wedge \neg B$$

or otherwise show

$$\vdash (P \rightarrow Q) \vee (Q \rightarrow P).$$