

Assignment Project Exam Help

Models for Nonstationary Time Series


https://powcoder.com

MAS 640 - Time Series Analysis and Forecasting

Add WeChat powcoder


1/29/2018

Determining Model Order



	ACF	PACF
[1]	0.87	0.87



Determining Model Order



	ACF	PACF
[1, 1]	0.50	0.50

Determining Model Order


Series: sim2



Add WeChat powcoder

ACF PACF
[1,] 0.82 0.82

Determining Model Order



ACF PACF
[1] 0.11 0.11

Assignment Project Exam Help

- ▶ Models for Nonstationary Time Series

<https://powcoder.com>

- ▶ Chapter 5 from text

Add WeChat powcoder

Assignment Project Exam Help

Any time series without a constant mean over time is nonstationary.

<https://powcoder.com>

$$Y_t = \mu_t + \epsilon_t$$

If μ varies across t , the series is nonstationary

Add WeChat powcoder

Detrending and Stationarity

Assignment Project Exam Help

- ▶ We fixed this issue by building a model for μ_t and studying the residuals

- ▶ “Detrending”

- ▶ We called this modeling “deterministic” trends
- ▶ Only reasonable if we assume this trend is an intrinsic property of the time series
- ▶ Implicitly assumes that the trend is “forever”
 - ▶ Which is often difficult to believe

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

- ▶ Consider the random walk process


$Y_t = Y_{t-1} + e_t$
<https://powcoder.com>

- ▶ By definition, it has constant mean $\mu_t = 0$
- ▶ Mistaking a trend seen from a random walk as deterministic ("forever") wouldn't be appropriate

Add WeChat powcoder

Random Walk - 30 Days of Data

30 realizations from a random walk process




Assignment Project Exam Help

<https://powcoder.com>


Add WeChat powcoder

Random Walk - 180 Days of Data

180 realizations from a random walk process



Price of Gold




Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Price of Gold


Assignment Project Exam Help



<https://powcoder.com>

Add WeChat powcoder

Oil Price




Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Oil Price




Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

CO2 Levels



Assignment Project Exam Help

- ▶ An alternative approach is to study the differenced time series


$\nabla Y_t = Y_t - Y_{t-1}$
<https://powcoder.com>

- ▶ No assumptions on trend through time
- ▶ No model to fit or parameters to estimate
- ▶ Tends to work well in practice


Add WeChat powcoder

- ▶ Good, quick, simple for forecasts



CREF Time Series



CREF Differenced





Autocorrelations for Differenced CREF Data





	ACF	PACF
[1,]	0.06	0.06

Air Passenger Miles





Autocorrelations for Differenced airmiles Data





ACF PACF

[1] 0.31 0.31

Electricity



Autocorrelations for Differenced electricity Data



	ACF	PACF
[1,]	0.00	0.00

Differencing

Assignment Project Exam Help

- ▶ Moral of the story: differencing can effectively remove the nonconstant mean in a time series
- ▶ Rather than build a model for the trend and study residuals, study the differenced data
- ▶ Note: For exponential trends, you may need to take two differences!

`diff(diff(DATA))`

Implementation of Differencing

Assignment Project Exam Help

- ▶ Since differencing is so standard for time series analyses and forecasting, most software will allow you to specify it directly.
 - ▶ So rather than building a model and studying the residuals, we can simply pass the original data and say “difference it”

<https://powcoder.com>

```
sarima(x, p, d, q)
```

- Add WeChat powcoder
- ▶ p = AR order
 - ▶ d = Number of differences for stationarity
 - ▶ q = MA order

ARIMA(p , d , q)

Assignment Project Exam Help

ARMA models that need to be differenced are referred to as ARIMA models

- ▶ <https://powcoder.com>
- ▶ Autoregressive Integrated Moving Average
 - ▶ Denoted ARIMA(p , d , q)

Add WeChat powcoder

ARIMA(p, d, q)

ARIMA(p, d, q) models encompass every class of models we have encountered up to this point

► ARIMA(p, 0, q) =

► <https://powcoder.com>

► ARIMA(0, 0, q) =

► Add WeChat powcoder

► ARIMA(0, 0, 0) =

Stationarity

Assignment Project Exam Help

- ▶ Stationarity is a very important concept in time series and one that you will often hear. Broadly speaking, a time series is called **stationary** if...

1. No systematic change in the mean (no trend),
2. No systematic change in the variance,
3. No noticeable seasonal patterns exist

In other words, the properties of one section of the data are the same as any other section.

Add WeChat powcoder


<https://powcoder.com>

Assignment Project Exam Help

- ▶ If we have clear evidence of nonconstant variance over time, a suitable transformation might fix (or lessen the impact of) the nonconstant variance pattern.
- ▶ Any transformations we apply to the data should be a **first** step.
 - ▶ Transform the data before looking at differences or modeling the trend.

<https://powcoder.com>
Add WeChat powcoder

Monthly Electricity Usage in the US



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder


What do we learn from this time series plot?

Assignment Project Exam Help

- ▶ Variance is increasing over time
- ▶ Time series that exhibit a “fanning-out” shape are not stationary because the variance changes over time.
- ▶ Before modeling, we should transform the data to *stabilize* the variance.


<https://powcoder.com>
Add WeChat powcoder

$\log(\text{Electricity})$ in the US




Variance looks ok! Let's plot the differences next.

$\text{diff}(\log(\text{Electricity}))$ in the US



Remember: transform first, then difference

Box-Cox for Power Transformations



Assignment Project Exam Help

λ	$T(Y)$	Description
-2.0	$1/Y^2$	Inverse Square
-1.0	$1/Y$	Inverse or Reciprocal
-0.5	$1/\sqrt{Y}$	Inverse square root
0.0	$\ln(Y)$	Logarithm
0.5	\sqrt{Y}	Square root
1.0	Y	Identity (No transformation)
2.0	Y^2	Square

<https://powcoder.com>

Add WeChat powcoder

Comments on using BoxCox procedure

Assignment Project Exam Help

- ▶ In the electricity example, the optimal λ was found to be about -0.1. However, this transformation makes little sense and isn't interpretable.
- ▶ Interval suggested values between about -0.4 and 0.2.
- ▶ So log transformation would be appropriate.

Want to find a reasonable transformation, not necessarily an optimal..

Add WeChat powcoder

Variance Stabilizing Transformations

Assignment Project Exam Help

- ▶ Can only perform variance stabilizing transformations on positive time series

- ▶ All values > 0

- ▶ However, if some or all Y are negative, we can simply add the same positive constant c to every observation so that every value becomes positive


- ▶ This does not affect anything

<https://powcoder.com>

Add WeChat powcoder

Adding Constant to Obtain a Positive Time Series

Consider the hypothetical time series below. Note that observations 2, 4, and 8 are negative, reach as low as -2.



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder


Adding Constant to Obtain a Positive Time Series

Simply add 3 to it. Shape is maintained, but now it's entirely positive.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



Assignment Project Exam Help

- ▶ Suppose that Y_t has relatively stable percent changes from period to period. That is, suppose that

$$\text{https://powcoder.com}$$
$$Y_t = (1 + r_t) Y_{t-1}$$

- ▶ where $100r_t$ represents the percent change from Y_{t-1} to Y_t .

Add WeChat powcoder

Relationship with Returns

- ▶ Suppose now that we take the log of this time series, and then the difference.

Assignment Project Exam Help

$$\log(Y_t) - \log(Y_{t-1}) = \log\left(\frac{Y_t}{Y_{t-1}}\right) = \log(1 + r_t)$$

<https://powcoder.com>

- ▶ If r_t is relatively low (<20% returns), then $\log(1 + r_t) \approx r_t$
- ▶ Consequently,

Add WeChat powcoder

$$\nabla \log(Y_t) \approx r_t$$

- ▶ Common in time series studies of financial data where returns are important and meaningful