

Assignment Project Exam Help

Chapter 3: Poisson Process

<https://powcoder.com>

Add WeChat powcoder

The Poisson distribution

Assignment Project Exam Help

Reminder: A random variable N taking values in $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$ has a *Poisson distribution* with a parameter $\lambda > 0$ (and we write $N \sim \text{Pois}(\lambda)$), if its probability mass function is given by

<https://powcoder.com>

$$\mathbb{P}(N = n) = \frac{e^{-\lambda} \lambda^n}{n!}, \quad \text{for } n \in \mathbb{Z}_+.$$

The mean and variance of N are both equal to λ .

Add WeChat powcoder

The exponential distribution

Reminder: A random variable T has an exponential distribution with parameter $\lambda > 0$ (called the **rate**), denoted by $T \sim \exp(\lambda)$, if its distribution function is

$$F_T(t) = \begin{cases} 1 - e^{-\lambda t}, & \text{for } t \geq 0, \\ 0, & \text{for } t < 0. \end{cases}$$

It follows that the probability density function of T is

$$f_T(t) = \begin{cases} \lambda e^{-\lambda t}, & \text{for } t \geq 0, \\ 0, & \text{for } t < 0. \end{cases}$$

The mean of T is $1/\lambda$ and the variance of T is $1/\lambda^2$.

The law of rare events

The Poisson distribution arises as the limit of the binomial distribution: Fix $\lambda > 0$, $k \in \mathbb{Z}_+$.

- ▶ If $X_n \sim \text{Bin}(n, \lambda/n)$ and $N \sim \text{Pois}(\lambda)$, then

$$\lim_{n \rightarrow \infty} \mathbb{P}(X_n = k) = \frac{e^{-\lambda} \lambda^k}{k!} = \mathbb{P}(N = k).$$

The exponential distribution arises as the limit of the geometric distribution: Fix $\lambda, t > 0$.

- ▶ If $Y_n \sim \text{Geo}(\lambda/n)$ (for $n > \lambda$) and $T \sim \text{Exp}(\lambda)$, then

$$\lim_{n \rightarrow \infty} \mathbb{P}(Y_n/n \leq t) = 1 - e^{-\lambda t} = \mathbb{P}(T \leq t).$$

The Poisson Process

Definition:

A nonnegative integer-valued process $(N_t)_{t \geq 0}$ is a **Poisson process** with a rate λ if

- (i) it has independent increments on disjoint intervals: for $k \geq 2$ and $0 \leq s_1 < t_1 \leq s_2 < t_2 \leq \dots < t_k$,

$$N_{t_1} - N_{s_1}, \dots, N_{t_k} - N_{s_k}$$

are independent variables, and

- (ii) For each $t > s \geq 0$, $N_t - N_s \sim \text{Pois}(\lambda(t - s))$.

Assignment Project Exam Help


If $(N_t)_{t \geq 0}$ is a Poisson process with rate λ , show that for fixed $r > 0$, the process $(N_t^*)_{t \geq 0}$ defined by $N_t^* = N_{t+r} - N_r$, is also Poisson process with rate λ

<https://powcoder.com>

[This is true even if r is a stopping time for the process.]

Add WeChat powcoder

A trajectory



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

- ▶ Earthquakes
- ▶ Grazing animal's head raises
- ▶ Horse kick deaths

<https://powcoder.com>

Add WeChat powcoder

The Poisson Process

Let $T_0 = 0$ and $T_j = \min\{t : N_t = j\}$ (the time of j^{th} jump) and define $\tau_j = T_j - T_{j-1}$ (time between the $(j-1)^{\text{st}}$ and j^{th} jumps).

Theorem: $(N_t)_{t \geq 0}$ is a Poisson process with rate λ if and only if $(\tau_j)_{j \in \mathbb{N}}$ are independent $\text{Exp}(\lambda)$ random variables.

Proof: The key to the proof is to observe that the event $\{T_j \leq t\}$ is the same as $\{N_t \geq j\}$. That is the waiting time until the j th jump is less than or equal to t if and only if there are j or more jumps up to (and including) time t .

Assume that $(N_t)_{t \geq 0}$ is a Poisson process. Then

$$\mathbb{P}(T_1 \leq t) = \mathbb{P}(N_t \geq 1) = 1 - \mathbb{P}(N_t = 0) = 1 - e^{-\lambda t}, \text{ so}$$

$$T_1 \sim \text{Exp}(\lambda).$$

The Poisson Process

Furthermore, we have

$$\begin{aligned}\mathbb{P}(T_j \leq t) &= \mathbb{P}(N_t \geq j) \\ &= \sum_{k=j}^{\infty} e^{-\lambda t} \frac{(\lambda t)^k}{k!} \\ &= 1 - \sum_{k=0}^{j-1} e^{-\lambda t} \frac{(\lambda t)^k}{k!}.\end{aligned}$$

This final expression is the distribution function for gamma distribution with parameter k and rate λ . You can check this by differentiating to get the density function

$$f_{T_j}(t) = e^{-\lambda t} \lambda^j t^{j-1} / (j-1)!.$$

So the waiting time until the j th event is the sum of j independent exponentially-distributed inter-event times with parameter λ .

The Poisson Process

This argument also holds in reverse.
Assuming that $\tau_1 \sim \text{Exp}(\lambda)$, we know that $\mathbb{P}(T_1 \leq t) = 1 - e^{-\lambda t}$,

which tells us that $\mathbb{P}(N_t = 0) = e^{-\lambda t}$.

Furthermore, for $j > 1$, if $\{\tau_1, \dots, \tau_j\}$ are i.i.d. $\text{Exp}(\lambda)$, then T_j has a Gamma distribution with parameters λ and j . So

$$\mathbb{P}(N_t \geq j) = \mathbb{P}(T_j \leq t) = 1 - \sum_{k=0}^{j-1} e^{-\lambda t} \frac{(\lambda t)^k}{k!},$$

Add WeChat powcoder

which tells us that N_t has a Poisson distribution with parameter λt .

Assignment Project Exam Help

We'd still need to show that $N_{t_i} - N_{s_i}$ are independent $\sim \text{Pois}(\lambda(t_i - s_i))$ over sets $[s_i, t_i)$ of disjoint intervals.

This follows from the memoryless property of the exponential distribution (so the remaining time from s_i until the next jump doesn't depend on the time $s_i - T_{N_{s_i}}$ since the previous jump) and the independence of the τ_j .

Add WeChat powcoder

Order statistics

For random variables $\xi_1, \xi_2, \dots, \xi_k$, denote by $\xi_{(i)}$ the i^{th} smallest of them. Then $\xi_{(1)}, \xi_{(2)}, \dots, \xi_{(k)}$ are called the **order statistics** associated with $\xi_1, \xi_2, \dots, \xi_k$.

For example, if we sample these random variables and find that $\xi_1 = 1.3, \xi_2 = 0.9, \xi_3 = 0.7, \xi_4 = 1.1$ and $\xi_5 = 1.5$, then $\xi_{(1)} = 0.7, \xi_{(2)} = 0.9, \dots, \xi_{(5)} = 1.5$.

Order statistics play a very important role in applications. For example, the maximum likelihood estimator of θ for a sample $\xi_1, \xi_2, \dots, \xi_k$ from the uniform $[0, \theta]$ distribution is $\xi_{(k)}$.

Order Statistics: examples

- ▶ If X_1, X_2 and X_3 are independent and identically-distributed random variables taking values 1, 2 and 3 each with probability $1/3$, find the joint probability mass function of $(X_{(1)}, X_{(2)}, X_{(3)})$.

- ▶ If Y_1, Y_2, \dots, Y_k are i.i.d. random variables with distribution function F , the distribution function of $Y_{(i)}$ is

$$F_{Y_{(i)}}(x) = \sum_{\ell=i}^k \binom{k}{\ell} F(x)^\ell (1 - F(x))^{k-\ell}.$$

- ▶ So, for the special case where Y_1, Y_2, \dots, Y_k are i.i.d. $\sim U[0, t]$, for $i \leq k$ and $x \leq t$ the distribution function of the order statistic $Y_{(i)}$ is given by

$$F_{Y_{(i)}}(x) = \sum_{\ell=i}^k \binom{k}{\ell} (x/t)^\ell (1 - x/t)^{k-\ell}.$$

Order Statistics

- Above we saw that if Y_1, Y_2, \dots, Y_k are i.i.d. with distribution function F the distribution function of $Y_{(i)}$ is

$$F_{Y_{(i)}}(x) = \sum_{\ell=i}^k \binom{k}{\ell} F(x)^\ell (1 - F(x))^{k-\ell}.$$

If they are also absolutely continuous, with density f , then the density of the order statistic $Y_{(i)}$ is

$$\begin{aligned} f_{Y_{(i)}}(x) &= \binom{k}{i-1} (k-i+1) F(x)^{i-1} f(x) (1 - F(x))^{k-i} \\ &= \binom{k}{i} i F(x)^{i-1} f(x) (1 - F(x))^{k-i}. \end{aligned}$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Similarly the joint densities for $1 \leq r \leq k$ and $x_1 < \dots < x_r$ are

Assignment Project Exam Help

$$f_{Y_{(i_1)}, \dots, Y_{(i_r)}}(x_1, \dots, x_r) = \binom{k}{i_1 - 1, 1, i_2 - i_1 - 1, 1, \dots, 1, k - i_r} \times \prod_{j=1}^r f(x_j) \prod_{j=0}^r (F(x_{j+1}) - F(x_j))^{i_{j+1} - i_j - 1},$$

where $\binom{\ell}{a_1, \dots, a_j}$ is the number of ways to choose subsets of sizes a_1, \dots, a_j from a set of size ℓ and for the sake of brevity we set $x_0 = -\infty$ and $x_{r+1} = \infty$ so $F(x_0) = 0$ and $F(x_{r+1}) = 1$.

Add WeChat powcoder

Assignment Project Exam Help

In particular for $r = k$, $x_1 < \cdots < x_r$,

$$f_{Y_{(1)}, \dots, Y_{(k)}}(x_1, \dots, x_k) = c \prod_{j=1}^k f(x_j)$$

<https://powcoder.com>

Add WeChat powcoder

The Poisson Process

Theorem: The conditional distribution of (T_1, \dots, T_k) given that $N_t = k$ is the same as the distribution of order statistics of a sample of k independent and identically-distributed random variables uniformly distributed on $[0, t]$.

I.e. $(T_1, \dots, T_k) | \{N_t = k\} \stackrel{d}{=} (U_{(1)}, \dots, U_{(k)})$

where U_1, \dots, U_k are independent $\sim U(0, t)$.

The same representation holds for the conditional distribution of (T_1, \dots, T_k) given that $T_{k+1} = t$.

"Proof" of theorem

According to our derivation for order statistics, $(U_{(1)}, \dots, U_{(k)})$

has density $k!t^{-k}$ for $0 = x_0 < x_1 < \dots < x_k < t$.

So we show the LHS has the same density:

$$\begin{aligned} & \mathbb{P}(T_1 \in dx_1, \dots, T_k \in dx_k | N_t = k) \\ &= \frac{\mathbb{P}(T_1 \in dx_1, T_2 \in d(x_2 - x_1), \dots, T_k \in d(x_k - x_{k-1}), T_{k+1} > t - x_k)}{\mathbb{P}(N_t = k)} \\ &= \frac{(\prod_{i=1}^k \lambda e^{-\lambda(x_i - x_{i-1})}) e^{-\lambda(t - x_k)}}{(\lambda t)^k e^{-\lambda t} / k!} dx_1 \dots dx_k \\ &= k! t^{-k} dx_1 \dots dx_k \end{aligned}$$

The proof of the second claim is similar.

Assignment Project Exam Help

The theorem implies that if τ_1, \dots, τ_n are i.i.d. exponential variables, then

$$\left(\frac{\tau_1}{\sum_{j=1}^{n+1} \tau_j}, \frac{\tau_1 + \tau_2}{\sum_{j=1}^{n+1} \tau_j}, \dots, \frac{\sum_{j=1}^n \tau_j}{\sum_{j=1}^{n+1} \tau_j} \right)$$

have the same distribution as $U(0, 1)$ order statistics.

Add WeChat powcoder

Superposition of Poisson processes

Theorem: Let $(N_t)_{t \geq 0}$ and $(M_t)_{t \geq 0}$ be two independent Poisson processes with rates λ_1 and λ_2 respectively and $L_t = N_t + M_t$.

Then $(L_t)_{t \geq 0}$ is a Poisson process with rate $\lambda_1 + \lambda_2$.

Proof:

- ▶ By independence, $L_t - L_s \sim \text{Pois}(\lambda_1(t-s) + \lambda_2(t-s))$.
- ▶ For disjoint $[s_1, t_1]$ and $[s_2, t_2]$,

$$L_{t_1} - L_{s_1} = (N_{t_1} - N_{s_1}) + (M_{t_1} - M_{s_1})$$

$$L_{t_2} - L_{s_2} = (N_{t_2} - N_{s_2}) + (M_{t_2} - M_{s_2})$$

which are independent because of the same property of $(N_t)_{t \geq 0}$ and $(M_t)_{t \geq 0}$. This argument extends to all finite collections of disjoint intervals.

Superposition example

Assignment Project Exam Help

A shop has two entrances, one from East St. the other from West St. Flows of customers through the two entrances are independent Poisson processes with rates 0.5 and 1.5 per minute, respectively.

- ▶ What is the probability that no new customers enter the shop in a fixed three minute time interval?
- ▶ What is the mean time between arrivals of new customers?
- ▶ What is the probability that a given customer entered from West St?

<https://powcoder.com>

Add WeChat powcoder

Thinning of a Poisson process

Assignment Project Exam Help

Suppose in a Poisson process $(N_t)_{t \geq 0}$ each 'customer' is 'marked' independently with probability p . Let M_t count the number of 'marked customers' that arrive on $[0, t]$.

Theorem. The processes $(M_t)_{t \geq 0}$ and $(N_t - M_t)_{t \geq 0}$ are independent Poisson processes with rates λp and $\lambda(1 - p)$ respectively.

Add WeChat powcoder

Thinning - "proof"

Assignment Project Exam Help

$$\begin{aligned}\mathbb{P}(M_t = j, N_t - M_t = k) &= \mathbb{P}(M_t = j, N_t = k + j) \\ &= \mathbb{P}(M_t = j | N_t = k + j) \mathbb{P}(N_t = k + j) \\ &= \binom{k+j}{j} p^j (1-p)^k \frac{e^{-\lambda t} (\lambda t)^{k+j}}{(k+j)!} \\ &= \frac{e^{-p\lambda t} (p\lambda t)^j e^{-(1-p)\lambda t} ((1-p)\lambda t)^k}{j! k!}.\end{aligned}$$

<https://powcoder.com>

Add WeChat powcoder

Thinning - example

The flow of customers to a shop is a Poisson process with rate 25 customers per hour. Each of the customers independently has a probability $p = 0.8$ of making a purchase.

- ▶ What is the probability that all customers who enter the shop during the time interval from 11.00 am to 11.15 am make a purchase?
- ▶ What is the probability that, conditional on there being two customers that made a purchase during that period, all customers who enter the shop during the time interval make a purchase?

The Compound Poisson Process

Assignment Project Exam Help

Suppose that $(N_t)_{t \geq 0}$ is a Poisson process and $(X_i)_{i \in \mathbb{N}}$ are independent and identically-distributed random variables, which are also independent of $(N_t)_{t \geq 0}$.

For $t \geq 0$, define $Y_t = \sum_{j \leq N_t} X_j$. Then $(Y_t)_{t \geq 0}$ is called a compound Poisson process.

It can be shown that $(Y_t)_{t \geq 0}$ has independent increments and it is possible to compute the distribution of Y_t by conditioning on N_t .

Add WeChat powecoder

Compound Poisson example

Assignment Project Exam Help

Suppose claims made to an insurance company arrive according to a Poisson process with rate λ , and each policy holder carries a policy for an amount X_k . Assume X_1, X_2, \dots are independent and identically distributed, and the number of claims and the size of claims are independent.

Calculate the mean and variance of the total amount of claims on the company up to time t .

Add WeChat powcoder