

Assignment Project Exam Help

Spline Regression

<https://powcoder.com>

MAST90083 Computational Statistics and Data Mining
Karim Seghouane
School of Mathematics & Statistics
The University of Melbourne

Add WeChat powcoder

Outline

Assignment Project Exam Help

§3.1 Introduction

§3.2 Motivation

§3.3 Spline

§3.4 Penalized Spline Regression

<https://powcoder.com>

Add WeChat powcoder

Introduction


Assignment Project Exam Help

- ▶ Some data sets are hard or impossible to model using traditional parametric techniques
- ▶ Many data sets also involve nonlinear effects that are difficult to model parametrically
- ▶ There is a need for flexible techniques to handle complicated nonlinear relationships
- ▶ Here we look at some ways of freeing oneself of the restrictions of parametric regression models

<https://powcoder.com>
Add WeChat powcoder

Introduction

The interest is the discovery of the **underlying trend** in the observed data which are treated as a collection of points on the plane



<https://powcoder.com>

Add WeChat powcoder

Introduction

Assignment Project Exam Help

- ▶ Alternatively, we could think of the vertical axis as a realization of a random variable y conditional on the variable x
- ▶ The underlying trend would then be a function

<https://powcoder.com>

$$f(x) = E(y|x)$$

- ▶ This can also be written as

Add WeChat powcoder

$$y_i = f(x_i) + \epsilon_i, \quad E(\epsilon_i) = 0$$

- ▶ and the problem is referred as nonparametric regression

Introduction

Assignment Project Exam Help

- ▶ Aim Estimate the unspecified smooth function from the pairs (x_i, y_i) , $i = 1, \dots, n$
- ▶ x here will be considered univariate
- ▶ There are several available methods, here we focus first on penalized splines

<https://powcoder.com>

Add WeChat powcoder

Motivation

Assignment Project Exam Help

- ▶ Let's start with the straight line regression model

$y_i = \beta_0 + \beta_1 x_i + \epsilon_i$
<https://powcoder.com>



Add WeChat powcoder


Motivation

Assignment Project Exam Help

- ▶ The corresponding basis for this model are the functions: 1 and x

<https://powcoder.com>

Add WeChat powcoder



- ▶ The model is a linear combination of these functions which is the reason for use of the world basis

Motivation

Assignment Project Exam Help

- ▶ The basis functions correspond to the columns of X for fitting the regression

<https://powcoder.com>

$$\begin{bmatrix} 1 & x_1 \\ \vdots & \vdots \\ 1 & x_n \end{bmatrix}$$

- ▶ Add WeChat powcoder


$$\hat{\mathbf{y}} = X \left(X^T X \right)^{-1} X^T \mathbf{y} = H \mathbf{y}$$

Motivation

Assignment Project Exam Help

- ▶ The quadratic model is a simple extension of the linear model

$y_i = \beta_0 + \beta_1 x_i + \beta_2 x_i^2 + \epsilon_i$
<https://powcoder.com>




Motivation

Assignment Project Exam Help

- There is an extra basis function x^2 corresponding to the addition of the $\beta_2 x_i^2$ term to the model

<https://powcoder.com>

Add WeChat powcoder



- The quadratic model is an example of how the simple linear model might be extended to handle nonlinear structure

Motivation

Assignment Project Exam Help

- ▶ The basis functions correspond to the columns of X for fitting the regression in the case of a quadratic model is given by

<https://powcoder.com>

$$\begin{bmatrix} 1 & x_1 & x_1^2 \\ \vdots & \vdots & \vdots \\ 1 & x_n & x_n^2 \end{bmatrix}$$

- ▶ Add WeChat powcoder

$$\hat{\mathbf{y}} = X \left(X^T X \right)^{-1} X^T \mathbf{y} = H \mathbf{y}$$


Spline basis function

Assignment Project Exam Help

- ▶ We know look at how the model can be extended to accommodate a different type of nonlinear structure

<https://powcoder.com>

Add WeChat powcoder




- ▶ Broken line model: it consists of two differently sloped lines that join together

Spline basis function

Assignment Project Exam Help

- Broken line: A linear combination of three basis functions



<https://powcoder.com>

Add WeChat powcoder

- where we have $(x - 0.6)_+$ with

$$u_+ = \begin{cases} u & u > 0 \\ 0 & u \leq 0 \end{cases}$$

Spline basis function

Assignment Project Exam Help

- Broken line model is

$$y_i = \beta_0 + \beta_1 x_i + \beta_{11} (x_i - 0.6)_+ + \epsilon_i$$

- which can be fit using the least square estimator with

$$\text{Add WeChat powcoder} \begin{bmatrix} 1 & x_1 & (x_1 - 0.6)_+ \\ \vdots & \vdots & \vdots \\ 1 & x_n & (x_n - 0.6)_+ \end{bmatrix}$$


Spline basis function

Assignment Project Exam Help

- Assume a more complicated structure

<https://powcoder.com>

Add WeChat powcoder




Spline basis function

Assignment Project Exam Help

► If we have good reason to believe that our underlying structure is of this basic, we could change the basis ?

<https://powcoder.com>

Add WeChat powcoder



► where the functions: $(x - 0.5)_+, (x - 0.55)_+, \dots, (x - 0.95)_+$

Spline basis function

Assignment Project Exam Help

- ▶ The basis can do a reasonable job with a linear portion between $x = 0$ and $x = 0.5$
- ▶ We can use least square to fit such model with

<https://powcoder.com>

Add WeChat powcoder

$$\begin{bmatrix} 1 & x_1 & (x_1 - 0.5)_+ & (x_1 - 0.55)_+ & \dots & (x_1 - 0.95)_+ \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_n & (x_n - 0.5)_+ & (x_n - 0.55)_+ & \dots & (x_n - 0.95)_+ \end{bmatrix}$$

Spline basis function

Assignment Project Exam Help

- ▶ It is possible to handle any complex type of structure by simply adding functions of the form $(x - k)_+$ to the basis
- ▶ This is equivalent to adding a column of values to the X matrix
- ▶ The value k is usually referred to as knots
- ▶ The function is made up of two lines that are tied together at $x = k$

<https://powcoder.com>

Add WeChat powcoder

Spline basis function

Assignment Project Exam Help

- ▶ The function $(x - 0.6)_+$ is called a linear spline basis function
- ▶ A set of such functions is called a linear spline basis
- ▶ Any linear combination of linear spline basis functions $1, x, (x - k_1)_+, \dots, (x - k_K)_+$ is a piecewise linear function with knots k_1, k_2, \dots, k_K and called spline

<https://powcoder.com>

Add WeChat powcoder

Spline basis function

Assignment Project Exam Help

- ▶ Rather than referring to the spline basis function $(x - k)_+$ it is common to simply refer to it knots k
- ▶ We say the model has a knot at 0.35 if the function $(x - 0.35)_+$ is the basis
- ▶ The spline model for a function f is

Add WeChat powcoder

$$f(x) = \beta_0 + \beta_1 x + \sum_{i=1}^K \beta_{ki} (x - k_i)_+$$


Illustration

▶ The selection of a good basis is usually challenging
▶ Start by trying to choose knots by trial

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder




Illustration

▶ The fit lacking in quality for low values of range
▶ An obvious remedy is to use more knots

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder




Illustration

- ▶ Larger set of knots, the fitting procedure has much more flexibility
- ▶ The plots is heavily overfitted

<https://powcoder.com>

Add WeChat powcoder




Illustration

- ▶ Pruning the knots to overcome the overfitting issue
- ▶ This fits the data well without overfitting
- ▶ This was arrived at, after a lot of time consuming trial and error

<https://powcoder.com>

Add WeChat powcoder



Knot selection

Assignment Project Exam Help

- ▶ A natural attempt at automatic selection of the knots is to use a model selection criterion
- ▶ If there are K candidate knots then there are 2^K possible models assuming the overall intercept and linear term are always present
- ▶ Highly computational intensive

<https://powcoder.com>

Add WeChat powcoder

Penalized spline regression

Assignment Project Exam Help

- ▶ Too many knots in the model induces roughness of the fit
- ▶ <https://powcoder.com> An alternative approach: retain all the knots but constrain their influence
- ▶ Hope: this will result in a less variable fit
- ▶ Consider a general spline model with K knots, K large

Add WeChat powcoder

Penalized spline regression

Assignment Project Exam Help

- ▶ The ordinary least square fit is written as

<https://powcoder.com>
 $\hat{\mathbf{y}} = \mathbf{X}\hat{\boldsymbol{\beta}}$ where $\hat{\boldsymbol{\beta}}$ minimizes $\|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|^2$

- ▶ and $\boldsymbol{\beta} = [\beta_0, \beta_1, \beta_{11}, \dots, \beta_{1K}]$ with β_{1k} the coefficient of the k^{th} knot.
- ▶ Unconstrained estimation of the $\boldsymbol{\beta}$ leads to a wiggly fit

Add WeChat powcoder

Penalized spline regression

Assignment Project Exam Help

Constraints on the β_{1k} that might help avoid this situation are

- ▶ $\max |\beta_{1k}| < C$
- ▶ $\sum |\beta_{1k}| < C$
- ▶ $\sum \beta_{1k}^2 < C$

<https://powcoder.com>

Add WeChat powcoder

Penalized spline regression

Assignment Project Exam Help

Define the matrix \mathbf{D} if size $(K + 2) \times (K + 2)$

$$\mathbf{D} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 1 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \cdots & 1 \end{bmatrix} = \begin{bmatrix} \mathbf{0}_{2 \times 2} & \mathbf{0}_{2 \times K} \\ \mathbf{0}_{K \times 2} & \mathbf{I}_{K \times K} \end{bmatrix}$$

<https://powcoder.com>
Add WeChat: powcoder

Penalized spline regression

Assignment Project Exam Help

- ▶ The third constraint is easier to implement than the first two
- ▶ The minimization problem

Minimize $\|y - X\beta\|^2$ subject to $\beta^T D \beta \leq c$

- ▶ This is equivalent to choosing β to minimize

Add WeChat powcoder

- ▶ with solution

$$\hat{\beta}_\lambda = (X^T X + \lambda D)^{-1} X^T y$$

Penalized spline regression

Assignment Project Exam Help


- ▶ The term $\lambda \beta^T \mathbf{D} \beta$ is called a roughness penalty since it penalizes fits that are too rough, thus yielding smoother result
- ▶ The amount of smoothness is controlled by λ , which is therefore referred to as a smoothing parameter
- ▶ The fitted values for penalized spline regression are

Add WeChat powcoder

$$\hat{\mathbf{y}} = \mathbf{X} \left(\mathbf{X}^T \mathbf{X} + \lambda \mathbf{D} \right)^{-1} \mathbf{X}^T \mathbf{y}$$

Illustration

- Linear penalized spline regression fits for different values of the smoothing parameter



Quadratic spline bases

Assignment Project Exam Help

- ▶ We have discussed linear splines, that is continuous, piecewise linear functions
- ▶ The reason for the piecewise linear nature of the functions ?
- ▶ is that they are a linear combination of piecewise linear functions of the form $(x - k)_+$
- ▶ A simple way of escaping from piecewise linearity ?

<https://powcoder.com>

Add WeChat powcoder

Quadratic spline bases

Assignment Project Exam Help

- ▶ We have discussed linear splines, that is continuous, piecewise linear functions
- ▶ The reason for the piecewise linear nature of the functions ?
- ▶ is that they are a linear combination of piecewise linear functions of the form $(x - k)_+$
- ▶ A simple way of escaping from piecewise linearity ?
- ▶ is to add x^2 to the basis and also to replace each $(x - k)_+$ by its square $(x - k)_+^2$

<https://powcoder.com>


Add WeChat powcoder

Illustration of a quadratic spline basis function

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



Quadratic spline bases

Assignment Project Exam Help

- ▶ The function doesn't have a sharp corner like $(x - 0.6)_+$ does
- ▶ The function $(x - 0.6)_+^2$ has a continuous first derivative
- ▶ Any linear combination of the functions

<https://powcoder.com>

$$1, x, x^2, (x - k_1)_+^2, \dots, (x - k_K)_+^2$$

- ▶ also have a continuous first derivative and not have any sharp corner
- ▶ This result in better fit
- ▶ This is called a quadratic spline basis with knots at k_1, \dots, k_K

Add WeChat powcoder


Illustration of quadratic spline basis functions

Assignment Project Exam Help

Quadratic spline do a better job of fitting peaks and valleys

<https://powcoder.com>

Add WeChat powcoder



Other spline bases

- ▶ We discussed linear and quadratic spline models
- ▶ One reason for considering other models is to achieve smoother fits → important if one plans to differentiate the fit to estimate derivative of the regression function
- ▶ In principle a change of basis does not change the fit but some bases are more stable and allow computation of a fit with better accuracy
- ▶ Besides numerical stability, ease of implementation is another reason for selecting one basis over another
- ▶ An obvious generalization is given by

$$1, x, \dots, x^p, (x - k_1)_+^p, \dots, (x - k_K)_+^p$$

- ▶ known as the truncated power basis of degree p

Other spline bases

Assignment Project Exam Help

▶ since the function $(x - k)_+^p$ has $p - 1$ continuous derivatives, higher values of p lead to smoother spline functions

- ▶ The p^{th} degree spline model is

$$f(x) = \beta_0 + \beta_1 x + \dots + \beta_p x^p + \sum_{i=1}^K \beta_{ki} (x - k_i)_+^p$$

- ▶ The expression for the fitted values is given by

$$\hat{\mathbf{y}} = \mathbf{X} \left(\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{D} \right)^{-1} \mathbf{X}^\top \mathbf{y}$$

$$\mathbf{D} = \text{diag}(\mathbf{0}_{p+1}, \mathbf{1}_K)$$